

acetylcholine receptor compositions containing the beta 4 subunit; Stephen F. Heinemann, et al., 435/6, \*\*69.1\*\*, 252.3, \*\*320.1\*\*, 536/23.5 [IMAGE AVAILABLE]

14. 5,371,188, Dec. 6, 1994, Neuronal nicotinic acetylcholine receptor compositions; Stephen F. Heinemann, et al., 530/350, 435/6, \*\*69.1\*\*, 252.3, \*\*320.1\*\* [IMAGE AVAILABLE]

15. 5,369,028, Nov. 29, 1994, DNA and mRNA encoding human neuronal nicotinic acetylcholine receptor compositions and cells transformed with same; Michael M. Harpold, et al., 435/252.3, \*\*69.1\*\*, 69.7, 70.1, 71.1, 71.2, 536/23.5, 25.3 [IMAGE AVAILABLE]

16. 5,300,436, Apr. 5, 1994, Genetically modified tyrosine hydroxylase and uses thereof; Menek Goldstein, et al., 435/190, 252.3, \*\*320.1\*\*, 536/23.2 [IMAGE AVAILABLE]

=> s neuron? or microglia?

7840 NEURON?  
158 MICROGLIA?  
L12 7896 NEURON? OR MICROGLIA?  
=> s I12 and neurotox?

1641 NEUROTOX?  
L13 869 L12 AND NEUROTOX?  
=> s I13 and (cell death or apopto?)

227674 CELL  
21859 DEATH  
3013 CELL DEATH  
(CELL(W)DEATH)  
849 APOPTO?  
L14 306 L13 AND (CELL DEATH OR APOPTO?)  
=> s I14 and method#

1308657 METHOD#  
L15 306 L14 AND METHOD#  
=> s neurotox?(5a)(inhibit?)

1641 NEUROTOX?  
271147 INHIBIT?  
L16 128 NEUROTOX?(5A)(INHIBIT?)  
=> s I16 and (cell death or apopto?)

227674 CELL

21859 DEATH  
3013 CELL DEATH  
(CELL(W)DEATH)  
849 APOPTO?  
L17 49 L16 AND (CELL DEATH OR APOPTO?)  
=> d 40-

40. 5,464,871, Nov. 7, 1995, Aromatic nitro and nitroso compounds and their metabolites useful as anti-viral and anti-tumor agents; Ernest Kun, et al., 514/617, 564/166 [IMAGE AVAILABLE]

41. 5,444,042, Aug. 22, 1995, Method of treatment of neurodegeneration with calpain inhibitors; Raymond T. Bartus, et al., 514/2, 435/23, 184, 514/16, 17, 18, 457 [IMAGE AVAILABLE]

42. 5,336,689, Aug. 9, 1994, Tri- and tetra-substituted guanidines and their use as excitatory amino acid antagonists; Eckard Weber, et al., 514/634, 183, 457, 549/288, 552/8, 564/230, 237, 238, 239, 240 [IMAGE AVAILABLE]

43. 5,266,594, Nov. 30, 1993, Inhibitors of nitric oxide synthase and use thereof to prevent glutamate neurotoxicity; Valina L. Dawson, et al., 514/560 [IMAGE AVAILABLE]

44. 5,262,568, Nov. 16, 1993, Tri- and tetra-substituted guanidines and their use as excitatory amino acid antagonists; Eckard Weber, et al., 564/238, 549/288, 552/8, 564/230, 237, 239, 240 [IMAGE AVAILABLE]

45. 5,232,911, Aug. 3, 1993, Mixture of a non-covalent heterodimer complex and a basic amphiphatic peptide as cytotoxic agent; Juan C. Vidal, 514/12, 424/94.3, 542, 514/21, 530/300, 324, 350, 856 [IMAGE AVAILABLE]

46. 5,190,976, Mar. 2, 1993, N,N'-disubstituted guanidines and their use as excitatory amino acid antagonists; Eckard Weber, et al., 514/634 [IMAGE AVAILABLE]

47. 5,093,525, Mar. 3, 1992, N,N'-disubstituted guanidines and their use as excitatory amino acid antagonists; Eckard Weber, et al., 564/238, 239 [IMAGE AVAILABLE]

48. 5,011,834, Apr. 30, 1991, PCP receptor ligands and the use thereof; Eckard Weber, et al., 514/212 [IMAGE AVAILABLE]

49. 4,906,779, Mar. 6, 1990, N,N'-disubstituted guanidines and their use as excitatory amino acid antagonists; Eckard Weber, et al., 564/238 [IMAGE AVAILABLE]

=> s advanced glycation end product

156496 ADVANCED  
112 GLYCATION  
1559723 END  
616123 PRODUCT  
L18 1 ADVANCED GLYCATION END PRODUCT  
(ADVANCED(W)GLYCATION(W)END(W)PRODUCT)  
=> d

1. 5,246,971, Sep. 21, 1993, Method of inhibiting nitric oxide formation; Joseph R. Williamson, et al., 514/634, 866 [IMAGE AVAILABLE]

=> d kwic

US PAT NO: 5,246,971 [IMAGE AVAILABLE] L18: 1 of 1

SUMMARY:  
BSUM(10)  
In . . . the prevention by methylguanidine of diabetesinduced vascular dysfunction is attributable to its ability to block NO production rather than blocking \*\*advanced\*\* \*\*glycation\*\* \*\*end\*\*  
\*\*product\*\* formation.  
=> e stern, david/in

E#	FILE	FREQUENCY	TERM
E1	USPAT	1	STERN, CHRISTIAN/IN
E2	USPAT	1	STERN, CHRISTOPHER/IN
E3	USPAT	14	--> STERN, DAVID/IN
E4	USPAT	1	STERN, DAVID F/IN
E5	USPAT	27	STERN, DAVID L/IN
E6	USPAT	14	STERN, DAVID M/IN
E7	USPAT	8	STERN, DAVID R/IN
E8	USPAT	1	STERN, DEREK V/IN
E9	USPAT	23	STERN, DONALD J/IN
E10	USPAT	12	STERN, DONALD S/IN
E11	USPAT	1	STERN, DONOVAN P/IN
E12	USPAT	1	STERN, E GEORGE/IN

=> s e3-e7

14 "STERN, DAVID"/IN  
 1 "STERN, DAVID F"/IN  
 27 "STERN, DAVID L"/IN  
 14 "STERN, DAVID M"/IN  
 8 "STERN, DAVID R"/IN  
 64 ("STERN, DAVID"/IN OR "STERN, DAVID F"/IN OR "STERN, DAVID L"/IN OR "STERN, DAVID M"/IN OR "STERN, DAVID R"/IN)  
 IN OR "STERN, DAVID M"/IN OR "STERN, DAVID R"/IN  
 => s 119 and neurotox?  
 1641 NEUROTOX?  
 L20 0 L19 AND NEUROTOX?  
 => s 119 and presenilin?  
 3 PRESENILIN?  
 L21 0 L19 AND PRESENILIN?  
 => s 119 and (neuron? or microglia?)  
 7840 NEURON?  
 158 MICROGLIA?  
 L22 0 L19 AND (NEURON? OR MICROGLIA?)  
 => e yan, shi du/in  
 E# FILE FREQUENCY TERM  
 E1 USPAT 3 YAN, SAN JYH/IN  
 E2 USPAT 1 YAN, SAU CHI B/IN  
 E3 USPAT 0 -> YAN, SHI DU/IN  
 E4 USPAT 1 YAN, SHUIA/IN  
 E5 USPAT 2 YAN, SHIU HANG/IN  
 E6 USPAT 1 YAN, SUZIE C/IN  
 E7 USPAT 1 YAN, TAK W/IN  
 E8 USPAT 117 YAN, TSOUNG Y/IN  
 E9 USPAT 65 YAN, TSOUNG YUAN/IN  
 E10 USPAT 1 YAN, TSUN YEE/IN  
 E11 USPAT 1 YAN, WANG/IN  
 E12 USPAT 1 YAN, WANG ZHOA/IN  
 => e wolozin, benjamin/in  
 E# FILE FREQUENCY TERM  
 E1 USPAT 9 WOLOWSKI, ECKARD/IN  
 E2 USPAT 1 WOLOWYK, MICHAEL W/IN  
 E3 USPAT 1 -> WOLOZIN, BENJAMIN/IN  
 E4 USPAT 1 WOLOZIN, BENJAMIN L/IN  
 E5 USPAT 1 WOLPA, MARK E/IN  
 E6 USPAT 1 WOLFALKA, KONSTANTIN/IN  
 E7 USPAT 1 WOLPAW, JONATHAN R/IN  
 E8 USPAT 12 WOLPE, STEPHEN D/IN  
 E9 USPAT 2 WOLPENSINGER, WERNER/IN  
 E10 USPAT 1 WOLPER, ANDRE E/IN  
 E11 USPAT 1 WOLPER, ANDRE EBERHARD/IN  
 E12 USPAT 1 WOLPER, DONALD F/IN  
 => s e3-e4  
 1 "WOLOZIN, BENJAMIN"/IN  
 1 "WOLOZIN, BENJAMIN L"/IN  
 L23 2 ("WOLOZIN, BENJAMIN"/IN OR "WOLOZIN, BENJAMIN L"/IN)  
 => d 1-  
 1. 5,869,266, Feb. 9, 1999, Human olfactory neuron cultures to diagnose Alzheimer's disease; \*\*Benjamin L. Wolozin\*\*, et al., 435/7.21, 325, 368; 436/63, 503, 811 [IMAGE AVAILABLE]  
 2. 5,811,310, Sep. 22, 1998, The Alz-50 monoclonal antibody and diagnostic assay for alzheimer's disease; Hossein A. Ghanbari, et al., 436/518; 435/7.1, 7.21, 7.92, 70.21, 326; 436/528, 531, 811; 530/388.1 [IMAGE AVAILABLE]  
 => d 1- ab  
 US PAT NO: 5,869,266 [IMAGE AVAILABLE] L23: 1 of 2  
 ABSTRACT:  
 The present invention relates to a culture of human olfactory neurons. The neurons may display a normal neuronal pathology or a pathology characteristic of a generalized central nervous system disease. The cultured neurons can be used for neurotoxicity tests, screening for therapeutic drugs and anti-viral agents, and diagnosing Alzheimer's disease.  
 US PAT NO: 5,811,310 [IMAGE AVAILABLE] L23: 2 of 2  
 ABSTRACT:  
 The invention relates to an antigen associated with Alzheimer's disease and to antibodies specific for said antigen. This invention further relates to methods for diagnosing Alzheimer's disease utilizing assays containing Alzheimer's associated antigen, antibodies specific for said antigen and samples from an individual suspected of having Alzheimer's disease.  
 => file epoab  
 FILE 'EPOABS' ENTERED AT 14:56:06 ON 11 APR 1999  
 .....  
 \* EUROPEAN PATENT ABSTRACTS \*  
 .....  
 => s 19  
 L24 0 PRESENILIN?  
 => s 118  
 9760 ADVANCED  
 10 GLYCATION  
 348261 END  
 71373 PRODUCT  
 L25 0 ADVANCED GLYCATION END PRODUCT  
 (ADVANCED(W)GLYCATION(W)END(W)PRODUCT)  
 => s 117  
 105 NEUROTOX?  
 34921 INHIBIT?  
 5 NEUROTOX?(SA)(INHIBIT?)  
 44761 CELL  
 362 DEATH  
 75 CELL DEATH  
 (CELL(W)DEATH)  
 70 APOPTO?  
 L26 0 L16 AND (CELL DEATH OR APOPTO?)  
 => s (neuron? or microglia?) and (neurotox? or neurodegen?)  
 831 NEURON?  
 2 MICROGLIA?  
 105 NEUROTOX?  
 228 NEURODEGEN?  
 L27 68 (NEURON? OR MICROGLIA?) AND (NEUROTOX? OR NEURODEGEN?)  
 => s 127 and screen?  
 34918 SCREEN?  
 L28 6 L27 AND SCREEN?  
 => d 1- cit ab  
 1. US 05449609A, Sep. 12, 1995, Methods for \*\*screening\*\* for \*\*neurotoxicity\*\* using a clonal human teratocarcinoma cell line; DONALD P YOUNKIN, et al., G01N 33/567  
 US 05449609A L28: 1 of 6  
 ABSTRACT:  
 Methods for \*\*screening\*\* for excitotoxic effects of an agent on

**\*\*neurons\*\*** of the central nervous system are provided by the present invention.

2. US 05342942A, Aug. 30, 1994, Pyrazoloquinazoline derivatives as neurotrophic agents; JUAN C JAEN, et al., C07D 487/04; C07D 487/14; C07D 491/147; C07D 495/14

US 05342942A L28: 2 of 6

ABSTRACT:

Pyrazolo[5,1-b]quinazoline compounds, salts thereof, methods of production, intermediates in their production, pharmaceutical compositions containing said compounds, and methods for treating **\*\*neurodegenerative\*\*** disorders, tumors of **\*\*neural\*\*** origin, inflammation, allergy, and pain, and methods for **\*\*screening\*\*** compounds that interact with the neurotrophic receptors using said compositions are disclosed.

that interact with the neurotrophic receptors using said compositions are disclosed.

3. US 05334618A, Aug. 2, 1994, Method of preventing NMDA receptor-mediated **\*\*neural\*\*** damage; STUART A LIPTON, A61K 31/13

US 05334618A L28: 3 of 6

ABSTRACT:

Disclosed is a method for reducing non-ischemic NMDA receptor-mediated **\*\*neural\*\*** damage in a mammal by administering to the mammal a compound of the formula shown in FIG. 1 (or a physiologically-acceptable salt thereof), wherein R1 includes an amino group, R2-R17 are independently H or a short chain aliphatic group comprising 1-5 carbons, and R4 and R10 also may (independently) be a halogen or an acyl group. Also disclosed is a **\*\*screen\*\*** for antagonists of NMDA receptor mediated **\*\*neurotoxicity\*\*** which have an enhanced prospect for being clinically tolerated and selective against such **\*\*neurotoxicity\*\***.

which have an enhanced prospect for being clinically tolerated and selective against such **\*\*neurotoxicity\*\***.

4. WO 09422866A1, Oct. 13, 1994, PYRAZOLOQUINAZOLONE DERIVATIVES AS NEUROTROPHIC AGENTS; JUAN CARLOS JAEN, et al., C07D 487/04; A61K 31/505

WO 09422866A1 L28: 4 of 6

ABSTRACT:

Pyrazolo[5,1-b]quinazoline compounds, salts thereof, methods of production, intermediates in their production, pharmaceutical compositions containing said compounds, and methods for treating **\*\*neurodegenerative\*\*** disorders, tumors of **\*\*neural\*\*** origin, inflammation, allergy, and pain, and methods for **\*\*screening\*\*** compounds that interact with the neurotrophic receptors using said compositions are disclosed.

5. WO 09405275A1, Mar. 17, 1994, METHOD OF PREVENTING NMDA RECEPTOR-MEDIATED **\*\*NEURONAL\*\*** DAMAGE; STUART A LIPTON, A61K 31/13

WO 09405275A1 L28: 5 of 6

ABSTRACT:

Disclosed is a method for reducing non-ischemic NMDA receptor-mediated **\*\*neural\*\*** damage in a mammal by administering to the mammal a compound of the formula shown in Fig. 1 (or a physiologically-acceptable salt thereof), wherein R1 includes an amino group, R2-R17 are independently H or a short chain aliphatic group comprising 1-5 carbons, and R4 and R10 also may (independently) be a halogen or an acyl group. Also disclosed is a **\*\*screen\*\*** for antagonists of NMDA receptor mediated **\*\*neurotoxicity\*\*** which have an enhanced prospect for being clinically tolerated and selective against such **\*\*neurotoxicity\*\***.

which have an enhanced prospect for being clinically tolerated and selective against such **\*\*neurotoxicity\*\***.

6. WO 09005138A1, May 17, 1990, CYTOTOXIC AMYLOID PRECURSORS AND **\*\*SCREENING\*\*** ASSAYS USING THEM; RACHAEL L NEVE, et al., C07H 21/04; C12N 5/00; C12N 15/11; C12Q 1/02; G01N 33/48

WO 09005138A1 L28: 6 of 6

ABSTRACT:

**\*\*Neurotoxic\*\*** amyloid precursor proteins (NAPPs) are produced, e.g. using recombinant DNA, and used in an assay to **\*\*screen\*\*** candidate compounds for their ability to antagonize **\*\*neural\*\*** toxicity. Specifically, **\*\*neurons\*\*** are cultured in the presence of an NAPP that has been treated with the candidate compound. The assay is useful to **\*\*screen\*\*** candidate therapeutics for Alzheimer's Disease. Assays for the

presence of NAPP are useful for identifying and monitoring the progression of Alzheimer's Disease.

=> file uspat

FILE 'USPAT' ENTERED AT 15:00:07 ON 11 APR 1999

\*\*\*\*\*  
\* WELCOME TO THE \*  
\* U.S. PATENT TEXT FILE \*  
\*\*\*\*\*

=> s 128

7840 NEURON?  
158 MICROGLIA?  
1641 NEUROTOX?  
1628 NEURODEGEN?  
244698 SCREEN?  
L29 815 L27 AND SCREEN?  
=> s 129 and (peptide or peptidomimetic)

28593 PEPTIDE  
299 PEPTIDOMIMETIC  
L30 515 L29 AND (PEPTIDE OR PEPTIDOMIMETIC)  
=> s 129 and nucleic acid#

22095 NUCLEIC  
469030 ACID#  
21978 NUCLEIC ACID#  
(NUCLEIC(W)ACID#)  
L31 393 L29 AND NUCLEIC ACID#  
=> d 385-

385. 5,212,082, May 18, 1993, Genetically modified tyrosine hydroxylase and uses thereof; Menek Goldstein, et al., 435/190 [IMAGE AVAILABLE]

386. 5,210,026, May 11, 1993, Human MK gene and method of expression; Imre Kovessi, et al., 435/69.1, 252.3, 252.33, 320.1, 488; 536/23.5 [IMAGE AVAILABLE]

387. 5,202,257, Apr. 13, 1993, Isolated **\*\*nucleic\*\*** **\*\*acids\*\*** encoding glutamate receptor protein; Stephen F. Heinemann, et al., 435/252.3, 69.1, 320.1; 536/23.1, 24.3 [IMAGE AVAILABLE]

388. 5,196,333, Mar. 23, 1993, DNA sequences involved in **\*\*neural\*\*** degeneration, multicellular organisms containing same and uses thereof;

Marin Chalfie, et al., 435/369, 29, 69.1, 70.3; 536/23.5 [IMAGE AVAILABLE]

389. 5,180,820, Jan. 19, 1993, Brain-derived neurotrophic factor; Yves-Alain Barde, et al., 536/23.5; 435/69.1, 69.3, 320.1; 530/399, 412 [IMAGE AVAILABLE]

390. 5,141,856, Aug. 25, 1992, Expression of purified ciliary neurotrophic factor; Franklin D. Collins, et al., 435/69.1, 91.41, 235.1, 252.3, 252.33, 254.2, 254.21, 320.1, 360; 530/350; 536/23.51, 24.31 [IMAGE AVAILABLE]

391. 4,997,929, Mar. 5, 1991, Purified ciliary neurotrophic factor; Franklin D. Collins, et al., 435/365.1, 69.1, 69.4, 320.1, 369; 536/23.5, 23.51 [IMAGE AVAILABLE]

392. 4,866,042, Sep. 12, 1989, Method for the delivery of genetic material across the blood brain barrier; Edward A. Neuwelt, 424/93.2; 514/44 [IMAGE AVAILABLE]

393. 4,666,828, May 19, 1987, Test for Huntington's disease; James F. Gusella, 435/6, 91.53; 436/811 [IMAGE AVAILABLE]

=> s 130(10a)(compound or composition)

\*WARNING\* - PROXIMITY OPERATOR PRECEDENCE LEVEL CONFLICTS OR IS NOT CONSISTENT WITH FIELD CODE - 'AND' OPERATOR ASSUMED  
 L30(10a)(COMPOUND)  
 389711 COMPOUND  
 455263 COMPOSITION  
 L32 471 L30(10a)(COMPOUND OR COMPOSITION)

=> s 130(5a)(compound or composition)

\*WARNING\* - PROXIMITY OPERATOR PRECEDENCE LEVEL CONFLICTS OR IS NOT CONSISTENT WITH FIELD CODE - 'AND' OPERATOR ASSUMED  
 L30(5a)(COMPOUND)  
 389711 COMPOUND  
 455263 COMPOSITION  
 L33 471 L30(5a)(COMPOUND OR COMPOSITION)

=> s 129 and peptidomimetic

299 PEPTIDOMIMETIC  
 L34 17 L29 AND PEPTIDOMIMETIC

=> d 1-

1. 5,892,003, Apr. 6, 1999, Ciliary neurotrophic factor receptor antibodies; Samuel Davis, et al., 530/388.22 [IMAGE AVAILABLE]

2. 5,863,795, Jan. 26, 1999, Nucleic acids that encode peptides which modulate apoptosis; Thomas D. Chittenden, et al., 435/325, 243, 320.1, 410; 536/23.5, 24.31 [IMAGE AVAILABLE]

3. 5,856,171, Jan. 5, 1999, Cell death regulators; Stanley J. Korsmeyer, 435/254.2, 6, 252.3, 810; 530/324, 350, 388.1, 389.1, 827; 536/23.5 [IMAGE AVAILABLE]

4. 5,854,215, Dec. 29, 1998, Modulators of beta-amyloid peptide aggregation; Mark A. Findeis, et al., 514/12; 530/324, 345 [IMAGE AVAILABLE]

5. 5,854,204, Dec. 29, 1998, A beta peptides that modulate beta-amyloid aggregation; Mark A. Findeis, et al., 514/2, 12, 14; 530/324, 326 [IMAGE AVAILABLE]

6. 5,849,897, Dec. 15, 1998, Nucleic acid molecule encoding ciliary neurotrophic factor receptor; Samuel Davis, et al., 536/23.5; 435/69.1, 252.3, 254.2, 320.1, 325, 348; 530/350 [IMAGE AVAILABLE]

7. 5,844,079, Dec. 1, 1998, Vertebrate embryonic pattern-inducing proteins, and uses related thereto; Philip W. Ingham, et al., 530/350; 435/7.1, 69.1, 252.3, 320.1; 530/300; 536/23.1, 23.5 [IMAGE AVAILABLE]

8. 5,834,234, Nov. 10, 1998, Apoptosis associated protein Bbk; Gregory J. Gallo, 435/69.1, 70.1, 71.1, 243, 375; 530/324, 325, 326, 327, 350 [IMAGE AVAILABLE]

9. 5,817,626, Oct. 6, 1998, Modulators of beta-amyloid peptide aggregation; Mark A. Findeis, et al., 514/12; 530/324, 326, 345 [IMAGE AVAILABLE]

10. 5,807,708, Sep. 15, 1998, Conserving nucleic acid molecules and compositions; Dean A. Falb, et al., 435/69.1, 252.3, 254.11, 320.1, 325; 536/23.1, 23.5 [IMAGE AVAILABLE]

11. 5,795,734, Aug. 18, 1998, EPH receptor ligands, and uses related thereto; John G. Flanagan, et al., 435/69.1, 7.1, 252.3, 320.1, 325; 530/300, 350; 536/23.1, 23.5 [IMAGE AVAILABLE]

12. 5,767,252, Jun. 16, 1998, \*\*Neuronal\*\* cell growth factor, Narp; Paul Worley, et al., 530/399, 350 [IMAGE AVAILABLE]

13. 5,723,301, Mar. 3, 1998, Method to \*\*screen\*\* compounds that affect GAPDH binding to polyglutamine; James R. Burke, et al., 435/7.1 [IMAGE AVAILABLE]

14. 5,700,638, Dec. 23, 1997, Cell death regulator; Stanley J. Korsmeyer, 435/6, 7.1, 7.2, 7.21, 7.31, 7.8, 69.1, 477; 436/501; 530/350 [IMAGE AVAILABLE]

15. 5,656,725, Aug. 12, 1997, Peptides and compositions which modulate apoptosis; Thomas D. Chittenden, et al., 530/324, 325, 326, 327, 328, 329, 330 [IMAGE AVAILABLE]

16. 5,648,334, Jul. 15, 1997, Methods of treatment using ciliary neurotrophic factor; Samuel Davis, et al., 514/12, 2; 530/350, 399 [IMAGE AVAILABLE]

17. 5,426,177, Jun. 20, 1995, Ciliary neurotrophic factor receptor; Samuel Davis, et al., 530/395, 350, 839 [IMAGE AVAILABLE]

=> d his

(FILE 'USPAT' ENTERED AT 14:37:16 ON 11 APR 1999)

L1 1583 S 435/4/CCLS  
 L2 3908 S 435/69.1/CCLS  
 L3 2 S 435/172.1/CCLS  
 L4 49 S 435/368/CCLS  
 L5 5870 S 435/320.1/CCLS  
 L6 123 S 435/455/CCLS  
 L7 8933 S L1-L6  
 L8 1 S L7 AND PRESENILIN?  
 L9 3 S PRESENILIN?  
 L10 138 S L7 AND NEUROTOX?  
 L11 16 S L10 AND PC12  
 L12 7896 S NEURON? OR MICROGLIA?  
 L13 869 S L12 AND NEUROTOX?  
 L14 306 S L13 AND (CELL DEATH OR APOPTO?)  
 L15 306 S L14 AND METHOD#  
 L16 128 S NEUROTOX?(5A)(INHIBIT?)  
 L17 49 S L16 AND (CELL DEATH OR APOPTO?)  
 L18 1 S ADVANCED GLYCATION END PRODUCT E STERN, DAVID/IN  
 L19 64 S E3-E7  
 L20 0 S L19 AND NEUROTOX?  
 L21 0 S L19 AND PRESENILIN?  
 L22 0 S L19 AND (NEURON? OR MICROGLIA?)  
 E YAN, SHI DU/IN  
 E WOLOZIN, BENJAMIN/IN  
 L23 2 S E3-E4

FILE 'EPOABS' ENTERED AT 14:56:06 ON 11 APR 1999

L24 0 S L9  
 L25 0 S L18  
 L26 0 S L17  
 L27 68 S (NEURON? OR MICROGLIA?) AND (NEUROTOX? OR NEURODEGEN?)  
 L28 6 S L27 AND SCREEN?

FILE "USPAT" ENTERED AT 15:00:07 ON 11 APR 1999

L29 815 S L28  
L30 515 S L29 AND (PEPTIDE OR PEPTIDOMIMETIC)  
L31 393 S L29 AND NUCLEIC ACID#  
L32 471 S L30(10A)COMPOUND OR COMPOSITION)  
L33 471 S L30(5A)COMPOUND OR COMPOSITION)  
L34 17 S L29 AND PEPTIDOMIMETIC

=> log y

U.S. Patent & Trademark Office LOGOFF AT 15:23:29 ON 11 APR  
1999

CS Department of Neuropsychiatry, Kumamoto University School of Medicine,  
Japan. tkimura@kaiju.medic.kumamoto-u.ac.jp  
SO PATHOLOGY INTERNATIONAL, (1998 Aug) 48 (8) 575-9.  
Journal code: BXQ. ISSN: 1320-5463.  
CY Australia  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199901  
EW 19990104  
AB The recent identification of age-related accumulation of advanced glycation end-products (AGE) of the Maillard reaction in neurons\*\*\*  
the  
and vessels of the human brain suggests the involvement of AGE in the aging process. A variety of inclusions such as lipofuscin granules, corpora amylacea, Hirano bodies, granulovacuolar degenerations and ubiquitin-positive granular structures are found in the aged human brain.  
These age-related inclusions contain insoluble and non-degradable proteins. \*\*\*Advanced\*\*\* glycation\*\*\* end\*\*\* product\*\*\* -modified proteins are also known to be insoluble and protease resistant. The similarity between proteins in such inclusions and AGE-modified proteins suggests the presence of AGE in inclusions. To investigate this possibility, the presence of two known AGE structures, N epsilon(carboxymethyl)lysine (CML) and pentosidine, was examined in age-related inclusions. Immunohistochemical examination of the medial temporal area of brain tissues obtained at autopsy from seven non-demented elderly individuals demonstrated positive reactions in lipofuscin and corpora amylacea but not in other inclusions for anti-CML and anti-pentosidine antibodies. As CML and pentosidine are glycoxidation products among AGE, the results suggest that glycation and/or oxidation may be involved in the formation of lipofuscin granules and corpora amylacea.  
L8 ANSWER 2 OF 3 MEDLINE  
AN 1998340856 MEDLINE  
DN 98340856  
TI Accelerated formation of N epsilon-(carboxymethyl) lysine, an \*\*\*advanced\*\*\* glycation\*\*\* end\*\*\* product\*\*\* by glyoxal and 3-deoxyglucosone in cultured rat sensory neurons\*\*\*  
AU Niwa H, Takeda A, Wakai M, Miyata T, Yasuda Y, Mitsuma T, Kurokawa K;

'AB' IS NOT A VALID FIELD CODE  
0 RAGE/AB  
894 RAGE/BI  
L3 0 L1 AND RAGE/AB,BI  
=> s advanced glycation end product/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
0 ADVANCED GLYCATION END PRODUCT/AB  
69312 ADVANCED/BI  
1403 GLYCATION/BI  
161501 END/BI  
100849 PRODUCT/BI  
71 ADVANCED GLYCATION END PRODUCT/BI  
((ADVANCED(W)GLYCATION(W)END(W)PRODUCT/BI)  
L4 71 ADVANCED GLYCATION END PRODUCT/AB,BI  
=> s l1 or l4  
L5 478 L1 OR L4  
=> s l5 and neuron7/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
0 NEURON7/AB  
211506 NEURON7/BI  
L6 114 L5 AND NEURON7/AB,BI  
=> s l1 and neuron7/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
0 NEURON7/AB  
211506 NEURON7/BI  
L7 111 L1 AND NEURON7/AB,BI  
=> s l4 and neuron7/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
0 NEURON7/AB  
211506 NEURON7/BI  
L8 3 L4 AND NEURON7/AB,BI  
=> d l1- bib ab  
YOU HAVE REQUESTED DATA FROM 3 ANSWERS -  
CONTINUE? Y(N):y  
L8 ANSWER 1 OF 3 MEDLINE  
AN 1998405810 MEDLINE  
DN 98405810  
TI Localization of identified \*\*\*advanced\*\*\* glycation\*\*\* end\*\*\* product\*\*\* structures, N epsilon(carboxymethyl)lysine and pentosidine, in age-related inclusions in human brains.  
AU Kimura T, Takamatsu J, Miyata T, Miyakawa T, Horiuchi S

\*\*\*\*\*STN Columbus \*\*\*\*\*  
..  
FILE 'HOME' ENTERED AT 15:56:10 ON 11 APR 1999  
=> medline  
MEDLINE IS NOT A RECOGNIZED COMMAND  
The previous command name entered was not recognized by the system.  
For a list of commands available to you in the current file, enter "HELP COMMANDS" at an arrow prompt (=>).  
=> file medline  
COST IN U.S. DOLLARS ENTRY SESSION TOTAL  
FULL ESTIMATED COST 0.15 0.15  
FILE 'MEDLINE' ENTERED AT 15:56:17 ON 11 APR 1999  
FILE LAST UPDATED: 2 APR 1999 (19990402/UP). FILE COVERS 1966 TO DATE.  
MEDLINE has been reloaded to reflect the annual MeSH changes made by the National Library of Medicine for 1999. Enter HELP RLOAD for details.  
THIS FILE CONTAINS CAS REGISTRY NUMBERS FOR EASY AND ACCURATE SUBSTANCE IDENTIFICATION.  
=> s presentlin7/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
0 PRESENTLIN7/AB  
407 PRESENTLIN7/BI  
L1 407 PRESENTLIN7/AB,BI  
=> s l1 and advanced glycation end product/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
0 ADVANCED GLYCATION END PRODUCT/AB  
69312 ADVANCED/BI  
1403 GLYCATION/BI  
161501 END/BI  
100849 PRODUCT/BI  
71 ADVANCED GLYCATION END PRODUCT/BI  
((ADVANCED(W)GLYCATION(W)END(W)PRODUCT/BI)  
L2 0 L1 AND ADVANCED GLYCATION END PRODUCT/AB,BI  
=> s l1 and rage/ab,bi

Sobue G  
 CS Department of Neurology, Nagoya University School of Medicine, Japan.  
 SO BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1998 Jul 9) 248 (1) 93-7.  
 Journal code: 9Y8. ISSN: 0006-291X.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 19981002  
 AB The formation of advanced glycation end products (AGEs) is associated with pathophysiological changes with aging and disease processes. In the neurodegeneration in Alzheimer's disease and other neurodegenerative diseases. AGEs are speculated to play a role in their pathogenesis. We provide the first evidence for the induction of AGEs in cultured \*\*\*neuronal\*\*\* cells. Glyoxal and 3-deoxyglucosone (3-DG), AGE precursors, induced N epsilon-(carboxymethyl) lysine (CML), a well characterized and major AGE structure, in cultured rat sensory \*\*\*neurons\*\*\* in a time- and dose-dependent manner. CML formation was prevented by addition of aminoguanidine, an inhibitor of AGE formation. This culture system provides a useful model to analyze the role of the glycoxidation reaction in \*\*\*neuronal\*\*\* aging and neurodegenerative disorder.

L8 ANSWER 3 OF 3 MEDLINE  
 AN 96029671 MEDLINE  
 DN 96029671  
 TI The receptor for advanced glycation end products (RAGE) is a cellular binding site for amphoterin. Mediation of neurite outgrowth and co-expression of rage and amphoterin in the developing nervous system.  
 AU Hori O; Brett J; Slattery T; Cao R; Zhang J; Chen J X; Nagashima M; Lundh E R; Vijay S; Nitecki D; et al  
 CS Department of Physiology, Columbia University, College of Physicians and Surgeons, New York, New York 10032, USA..  
 NC AG00602 (NIA)  
 HL21006 (NHLBI)  
 SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1995 Oct 27) 270 (43) 25752-61.  
 Journal code: HIV. ISSN: 0021-9258.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)

LA English  
 FS Priority Journals; Cancer Journals  
 EM 199602  
 AB The receptor for advanced glycation end products (RAGE), a newly-identified member of the immunoglobulin superfamily, mediates interactions of \*\*\*advanced\*\*\* \*\*\*glycation\*\*\*  
 \*\*\*end\*\*\*  
 \*\*\*product\*\*\* (AGE)-modified proteins with endothelium and other cell types. Survey of normal tissues demonstrated RAGE expression in situations in which accumulation of AGEs would be unexpected, leading to the hypothesis that under physiologic circumstances, RAGE might mediate interaction with ligands distinct from AGEs. Sequential chromatography of bovine lung extract identified polypeptides with M(r) values of approximately 12,000 (p12) and approximately 23,000 (p23) which bound RAGE. NH2-terminal and internal protein sequence data for p23 matched that reported previously for amphoterin. Amphoterin purified from rat brain or recombinant rat amphoterin bound to purified sRAGE in a saturable and dose-dependent manner, blocked by anti-RAGE IgG or a soluble form of RAGE (sRAGE). Cultured embryonic rat \*\*\*neurons\*\*\*, which express RAGE, displayed dose-dependent binding of 125I-amphoterin which was prevented by blockade of RAGE using antibody to the receptor or excess soluble receptor (sRAGE). A functional correlate of RAGE-amphoterin interaction was inhibition by anti-RAGE F(ab)2 and sRAGE of neurite formation by cortical \*\*\*neurons\*\*\* specifically on amphoterin-coated substrates. Consistent with a potential role for RAGE-amphoterin interaction in development, amphoterin and RAGE mRNA/antigen were co-localized in developing rat brain. These data indicate that RAGE has physiologically relevant ligands distinct from AGEs which are likely, via their interaction with the receptor, to participate in physiologic processes outside of the context of diabetes and accumulation of AGEs.  
 => d his

(FILE 'HOME' ENTERED AT 15:56:10 ON 11 APR 1999)

FILE 'MEDLINE' ENTERED AT 15:56:17 ON 11 APR 1999  
 L1 407 S PRESENILIN7/AB.BI  
 L2 0 S L1 AND ADVANCED GLYCATION END PRODUCT/AB.BI  
 L3 0 S L1 AND RAGE/AB.BI  
 L4 71 S ADVANCED GLYCATION END PRODUCT/AB.BI  
 L5 478 S L1 OR L4  
 L6 114 S L5 AND NEURON7/AB.BI  
 L7 111 S L1 AND NEURON7/AB.BI  
 L8 3 S L4 AND NEURON7/AB.BI  
 => s (14)(3a)(receptor#)/ab.bi  
 'AB' IS NOT A VALID FIELD CODE  
 0 (RECEPTOR#)/AB  
 427150 (RECEPTOR#)/BI  
 L9 4 (14)(3a)(RECEPTOR#)/AB.BI  
 => d l- bib ab  
 YOU HAVE REQUESTED DATA FROM 4 ANSWERS -  
 CONTINUE? Y/(N);y

L9 ANSWER 1 OF 4 MEDLINE  
 AN 1999009111 MEDLINE  
 DN 99009111  
 TI A redox-triggered ras-effector interaction. Recruitment of phosphatidylinositol 3'-kinase to Ras by redox stress.  
 AU Deora A A; Win T; Vanhaesebroeck B; Lander H M  
 CS Department of Biochemistry, Cornell University Medical College, New York, New York 10021, USA.  
 NC GM55509 (NIGMS)  
 AL37637 (NIH/AD)  
 SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1998 Nov 6) 273 (45) 29923-8  
 Journal code: HIV. ISSN: 0021-9258.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199902  
 EW 19990204  
 AB Reactive free radical species are known to trigger biochemical events culminating in transcription factor activation and modulation of gene expression. The cytosolic signaling events triggered by free radicals result in nuclear responses are largely unknown. Here we identify a signaling cascade triggered immediately upon redox activation of Ras. We examined two physiologically relevant models of redox signaling: 1) nitric oxide in human T cells, and 2) advanced glycation end product in rat

phoeochromocytoma cells. Reactive free radical species generated by nitric oxide donors and the interaction of \*\*\*advanced\*\*\*  
 \*\*\*glycation\*\*\* \*\*\*product\*\*\* with its \*\*\*receptor\*\*\* led to the recruitment of p85/p110 phosphatidylinositol 3-kinase (PI3K) to the plasma membrane, where it associated directly with the effector domain of Ras and became activated. Only the p110beta and p110delta (but not p110alpha) catalytic subunits were recruited by redox-activated Ras.  
 Activation of downstream targets of PI3K such as protein kinase B/Akt and mitogen-activated protein kinase was found to be PI3K dependent. Our study demonstrates that nitrosative and oxidative stressors trigger Ras-dependent and PI3K-regulated events in cells and define a biochemical pathway that is triggered by redox signaling.

L9 ANSWER 2 OF 4 MEDLINE  
 AN 97368045 MEDLINE  
 DN 97368045  
 TI Recombinant \*\*\*advanced\*\*\* \*\*\*glycation\*\*\*  
 \*\*\*end\*\*\*  
 \*\*\*product\*\*\* \*\*\*receptor\*\*\* pharmacokinetics in normal and diabetic rats.  
 AU Renard C; Chappay O; Wautier M P; Nagashima M; Lundh E; Moser J; Zhao L;  
 Schmidt A M; Schermann J M; Wautier J L  
 CS Laboratoire de Recherche en Biologie Vasculaire et Cellulaire, Universite Paris 7, Hopital Lariboisiere, France.  
 SO MOLECULAR PHARMACOLOGY, (1997 Jul) 52 (1) 54-62.  
 Journal code: NGR. ISSN: 0026-895X.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199710  
 EW 19971003  
 AB Vascular dysfunction in patients with diabetes mellitus is related to advanced glycation end product (AGE) formation. We previously showed that AGES produce an increase in vascular permeability and generated an oxidant stress after binding to the receptor (RAGE) present on endothelium. RAGE, a 35-kDa protein that belongs to the immunoglobulin superfamily, has been cloned from a rat lung cDNA library, and recombinant rat soluble RAGE

(rR-RAGE) has been produced in insect cells. The sequence of RAGE is highly conserved between human and rat. We studied the biological effect of rR-RAGE and pharmacokinetics of 125I-rR-RAGE after intravenous or intraperitoneal administration in normal and streptozotocin-induced diabetic rats. rR-RAGE prevented albumin or inulin transfer through a bovine aortic endothelial cell monolayer, restored the hyperpermeability observed in diabetic rats or induced in normal rats by diabetic rat red blood cells, and corrected the reactive oxygen intermediate production after intravenous or intraperitoneal administration. After intravenous injection of 125I-rR-RAGE, the distribution half-life was longer (p < or = 0.01) in diabetic (0.15 and 4.01 hr) than in normal (0.02 and 0.21 hr) rats, as was the case for the elimination half-lives (diabetic, \$7.17 hr; normal, 26.02 hr; p < or = 0.01). Distribution volume was higher in diabetic than in normal rats (6.94 and 3.24 liter/kg, respectively; p = 0.049). Our study showed that rR-RAGE was biologically active in vivo and slowly cleared, which suggests it could be considered as a potential therapy.

L9 ANSWER 3 OF 4 MEDLINE  
 AN 96289840 MEDLINE  
 DN 96289840  
 TI Recent progress in advanced glycation and diabetic vascular disease: role of \*\*\*advanced\*\*\* \*\*\*glycation\*\*\* \*\*\*end\*\*\*  
 \*\*\*product\*\*\*  
 \*\*\*receptors\*\*\*  
 AU Viassara H; Bucala R  
 CS Picower Institute for Medical Research, Manhasset, New York 10030, USA.  
 SO DIABETES, (1996 Jul) 45 Suppl 3 S65-6. Ref: 17  
 Journal code: E8X. ISSN: 0012-1797.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LA English  
 FS Abridged Index Medicus Journals; Priority Journals  
 EM 199610  
 AB Advanced glycosylation end products (AGEs) form principally from the rearrangement of early glycation products, i.e., Amadori products, which produce a class of stable moieties that possess distinctive chemical crosslinking and biological properties. It has been generally believed

that proteins with half-lives of longer than a few weeks are most susceptible to advanced glycosylation and that the highest levels of AGES occur on proteins that comprise the long-lived structural components of connective tissue matrix and basement membrane.

L9 ANSWER 4 OF 4 MEDLINE  
 AN 95096076 MEDLINE  
 DN 95096076  
 TI Receptor-mediated endocytic uptake of methylglyoxal-modified serum albumin. Competition with advanced glycation end product-modified serum albumin at the \*\*\*advanced\*\*\* \*\*\*glycation\*\*\*  
 \*\*\*end\*\*\*  
 \*\*\*product\*\*\* \*\*\*receptor\*\*\*  
 AU Westwood M E; McLellan A C; Thornalley P J  
 CS Department of Chemistry and Biological Chemistry, University of Essex, Colchester, United Kingdom.  
 SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1994 Dec 23) 269 (51) 32293-8.  
 Journal code: HIV. ISSN: 0021-9258.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199503  
 AB Methylglyoxal binds and irreversibly modifies arginine and lysine residues in bovine serum albumin (BSA) under physiological conditions, producing a protein with an increased net negative charge at physiological pH. At 4 degrees C, methylglyoxal-modified BSA (MG-BSA) was bound by cell surface receptors on murine P388D1 macrophages. The apparent dissociation constant KD value was 435 +/- 2 nM, and there were 8.89 +/- 0.02 x 10(5) receptors/cell (n = 6), compare with an apparent KD value of 263 +/- 52 nM and 10.17 +/- 0.93 x 10(5) receptors/cell (n = 11) for advanced glycation end product-modified BSA (AGE-BSA). AGE-BSA competed with MG-BSA for binding to a common receptor; however, a component of AGE-BSA receptor binding could not be displaced by MG-BSA, and a component of receptor binding could not be displaced by AGE-BSA, suggesting that there are binding sites for both AGE-BSA and MG-BSA, competitive and noncompetitive, to MG-BSA and AGE-BSA on P388D1 cells at 4 degrees C. At 37 degrees C, receptor binding of AGE-BSA and MG-BSA was



followed by endocytosis and lysosomal degradation of the modified protein. Methylglyoxal-modified proteins are ligands for the AGE receptor, and their formation and metabolism may be linked to the development of diabetic complications.

=> file medline embase biosis wpids caplus

COST IN U.S. DOLLARS	ENTRY	SINCE FILE	SESSION	TOTAL
4.19	4.19	4.34		

FILE MEDLINE ENTERED AT 16:02:33 ON 11 APR 1999

FILE 'EMBASE' ENTERED AT 16:02:33 ON 11 APR 1999  
COPYRIGHT (C) 1999 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 16:02:33 ON 11 APR 1999  
COPYRIGHT (C) 1999 BIOSIS(R)

FILE 'WPIDS' ENTERED AT 16:02:33 ON 11 APR 1999  
COPYRIGHT (C) 1999 DERWENT INFORMATION LTD

FILE 'CAPLUS' ENTERED AT 16:02:33 ON 11 APR 1999  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 1999 AMERICAN CHEMICAL SOCIETY (ACS)

=> s 11 and 19

'AB' IS NOT A VALID FIELD CODE  
2 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
4 FILES SEARCHED...  
L10 0 L1 AND L9

=> s mutant presenilin-2/ab.bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L11 10 MUTANT PRESENILIN-2/AB.BI

=> dup rem 111

PROCESSING COMPLETED FOR L11  
L12 4 DUP REM L11 (6 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 4 ANSWERS -

CONTINUE? Y(N)y

L12 ANSWER 1 OF 4 MEDLINE DUPLICATE 1  
AN 1998361992 MEDLINE  
DN 98361992

TI Molecular dissection of domains in \*\*\*mutant\*\*\*  
T1 \*\*\*presenilin\*\*\*

\*\*\*\* that mediate overproduction of amyloidogenic forms of amyloid beta peptides. Inability of truncated forms of PS2 with familial Alzheimer's disease mutation to increase secretion of Abeta42.

AU Tomita T; Tokuhito S; Hashimoto T; Aiba K; Saïdo T C; Maruyama K; Iwatsubo T

CS Department of Neuropathology and Neuroscience, Graduate School of Pharmaceutical Sciences, University of Tokyo, Tokyo 113-0033, Japan.  
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1998 Aug 14) 273 (33) 21153-60.

Journal code: HIV. ISSN: 0021-9258.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)

LA English  
FS Priority Journals; Cancer Journals  
EM 199811

EW 19981103  
AB Mutations in presenilin (PS) 1 or PS2 genes account for the majority of early-onset familial Alzheimer's disease, and these mutations have been shown to increase production of species of amyloid beta peptide ending at residue 42, i.e. the most amyloidogenic form of Abeta. To gain insight into the molecular mechanisms whereby mutant PS induces overproduction of Abeta42, we constructed cDNAs encoding mutant and/or truncated forms of PS2 and examined the secretion of Abeta42 from COS or neuro2a cells transfected with these genes. Cells expressing PS2 harboring both N141I and M239V mutations in the same polypeptide induced overproduction of Abeta42, although the levels of Abeta42 were comparable with those in cells engineered to express PS2 with one or the other of these PS2 mutations. In contrast, cells engineered to express partially truncated PS2 (eliminating the COOH-terminal third of PS2 while retaining the endoproteolytic NH2-terminal fragment) and harboring a N141I mutation, as well as cells expressing COOH-terminal fragments of PS2, did

not overproduce Abeta42, and the levels of Abeta42 were comparable with those in cells that expressed full-length, wild-type PS2 or fragments thereof. These data indicate that: (i) the Abeta42-promoting effects of mutant PS2 proteins reach the maximum level with a given single amino acid substitution (i.e. N141I or M239V); and (ii) the expression of full-length mutant PS2 is required for the overproduction of Abeta42. Hence, cooperative interactions of NH2- and COOH-terminal fragments generated from full-length mutant PS2 may be important for the overproduction of Abeta42 that may underlie familial Alzheimer's disease.

L12 ANSWER 2 OF 4 MEDLINE DUPLICATE 2  
AN 1998311214 MEDLINE  
DN 98311214

T1 \*\*\*Mutant\*\*\* \*\*\*presenilin\*\*\* \*\*\*\* transgenic mouse: effect on an age-dependent increase of amyloid beta-protein 42 in the brain.

AU Oyama F; Sawamura N; Kobayashi K; Morishima-Kawashima M; Kuramochi T; Ito M; Tomita T; Maruyama K; Saïdo T C; Iwatsubo T; Capell A; Walter J; Grunberg J; Ueyama Y; Haass C; Ihara Y

CS Department of Neuropathology, Faculty of Medicine, University of Tokyo, Japan.  
SO JOURNAL OF NEUROCHEMISTRY, (1998 Jul) 71 (1) 313-22.

Journal code: JAV. ISSN: 0022-3042.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)

LA English  
FS Priority Journals  
EM 199809

EW 19980902  
AB The N141I missense mutation in presenilin (PS) 2 is tightly linked with a form of autosomal dominant familial Alzheimer's disease (AD) in the Volga German families. We have generated transgenic mouse lines overexpressing human wild-type or mutant PS2 under transcriptional control of the chicken beta-actin promoter. In the brains of transgenic mice, the levels of PS2 mRNA were found to be five- to 15-fold higher than that of endogenous mouse PS2 mRNA. The amyloid beta-protein (Abeta) 42 levels in the brains of mutant PS2 transgenic mice were higher than those in wild-type PS2 transgenic mice at the age of 2, 5, or 8 months. In addition, the

Abeta42 levels appeared to increase steadily in the mutant PS2 transgenic mouse brains from 2 to 8 months of age, whereas there was only a small increase in wild-type transgenic mice between the ages of 5 and 8 months. There was no definite difference in the levels of N-terminal and C-terminal fragments between wild-type and mutant PS2 transgenic mice at the age of 2, 5, or 8 months. These data show a definite effect of the PS2 mutation on an age-dependent increase of Abeta42 content in the brain.	(FAD), the authors expressed cDNAs for wild-type PS2 and PS2 with the Volga German (N141I) mutation in cultured cells and then examined the metabolites of the transfected proteins and their effect on the C-terminal properties of secreted amyloid beta protein (A beta). PS2 was identified as a 50-55-kDa protein, which was cleaved to produce N-terminal fragments of 35-40 kDa and C-terminal fragments of 19-23 kDa. The Volga German (N141I) mutation did not cause any significant change in the metabolites of PS2. COS-1 cells doubly transfected with cDNAs for N141I mutant PS2 and human beta-amyloid precursor protein (beta APP) or a C-terminal fragment thereof, as well as mouse Neuro2a neuroblastoma cells stably transfected with N141I mutant PS2 alone, secreted 1.5-10-fold more A beta. residues 42 (or 43) [A beta 42(43)] compared with those expressing the wild-type PS2. Apparently, the PS2 mutation (N141I) linked to FAD alters the metabolite of A beta/beta APP to foster the production of the form of A beta that most readily deposits in amyloid plaques. Thus, mutant PS2 may lead to AD by altering the metabolite of A beta/beta APP.	=> e yan shi du/au E1 1 YAN SHENTSHAN/AU E2 7 YAN SHI/AU E3 73 -> YAN SHI DU/AU E4 28 YAN SHI FANG/AU E5 1 YAN SHI GU/AU E6 1 YAN SHI KUN/AU E7 1 YAN SHI LEI/AU E8 1 YAN SHI MING/AU E9 1 YAN SHI PIN/AU E10 45 YAN SHI PING/AU E11 1 YAN SHI QI/AU E12 2 YAN SHI QIANG/AU => s e2-e3 L15 80 ("YAN SHI"/AU OR "YAN SHI DU"/AU) => s l15 and presenilin7/ab,bi 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE L16 0 L15 AND PRESENILIN7/AB,BI => e wolozin, benjamin/au E1 3 WOLOZIN M W/AU E2 2 WOLOZIN R/AU E3 0 -> WOLOZIN, BENJAMIN/AU E4 1 WOLOZON B L/AU E5 1 WOLOZSZUK R/AU E6 1 WOLOZYN W/AU E7 2 WOLPA BRENDA/AU E8 1 WOLPA LORI J/AU E9 3 WOLPA M E/AU E10 1 WOLPART S I/AU E11 6 WOLPAW D R/AU E12 2 WOLPAW DANIEL R/AU => e wolozin, b/au E1 3 WOLOZIN M W/AU E2 2 WOLOZIN R/AU E3 0 -> WOLOZIN, B/AU E4 1 WOLOZON B L/AU E5 1 WOLOZSZUK R/AU E6 1 WOLOZYN W/AU E7 2 WOLPA BRENDA/AU E8 1 WOLPA LORI J/AU E9 3 WOLPA M E/AU E10 1 WOLPART S I/AU E11 6 WOLPAW D R/AU E12 2 WOLPAW DANIEL R/AU
Abeta42 levels appeared to increase steadily in the mutant PS2 transgenic mouse brains from 2 to 8 months of age, whereas there was only a small increase in wild-type transgenic mice between the ages of 5 and 8 months. There was no definite difference in the levels of N-terminal and C-terminal fragments between wild-type and mutant PS2 transgenic mice at the age of 2, 5, or 8 months. These data show a definite effect of the PS2 mutation on an age-dependent increase of Abeta42 content in the brain.	(FAD), the authors expressed cDNAs for wild-type PS2 and PS2 with the Volga German (N141I) mutation in cultured cells and then examined the metabolites of the transfected proteins and their effect on the C-terminal properties of secreted amyloid beta protein (A beta). PS2 was identified as a 50-55-kDa protein, which was cleaved to produce N-terminal fragments of 35-40 kDa and C-terminal fragments of 19-23 kDa. The Volga German (N141I) mutation did not cause any significant change in the metabolites of PS2. COS-1 cells doubly transfected with cDNAs for N141I mutant PS2 and human beta-amyloid precursor protein (beta APP) or a C-terminal fragment thereof, as well as mouse Neuro2a neuroblastoma cells stably transfected with N141I mutant PS2 alone, secreted 1.5-10-fold more A beta. residues 42 (or 43) [A beta 42(43)] compared with those expressing the wild-type PS2. Apparently, the PS2 mutation (N141I) linked to FAD alters the metabolite of A beta/beta APP to foster the production of the form of A beta that most readily deposits in amyloid plaques. Thus, mutant PS2 may lead to AD by altering the metabolite of A beta/beta APP.	=> e stern david/au E1 1 STERN DARRYL/AU E2 4 STERN DARRYL A/AU E3 162 -> STERN DAVID/AU E4 9 STERN DAVID A/AU E5 84 STERN DAVID B/AU E6 1 STERN DAVID BENJAMIN/AU E7 2 STERN DAVID E/AU E8 71 STERN DAVID F/AU E9 1 STERN DAVID FREDERICK/AU E10 2 STERN DAVID H/AU E11 4 STERN DAVID I/AU E12 40 STERN DAVID L/AU => s e3 L13 162 "STERN DAVID"/AU => s l13 and presenilin7/ab,bi 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE L14 0 L13 AND PRESENILIN7/AB,BI
Abeta42 levels appeared to increase steadily in the mutant PS2 transgenic mouse brains from 2 to 8 months of age, whereas there was only a small increase in wild-type transgenic mice between the ages of 5 and 8 months. There was no definite difference in the levels of N-terminal and C-terminal fragments between wild-type and mutant PS2 transgenic mice at the age of 2, 5, or 8 months. These data show a definite effect of the PS2 mutation on an age-dependent increase of Abeta42 content in the brain.	(FAD), the authors expressed cDNAs for wild-type PS2 and PS2 with the Volga German (N141I) mutation in cultured cells and then examined the metabolites of the transfected proteins and their effect on the C-terminal properties of secreted amyloid beta protein (A beta). PS2 was identified as a 50-55-kDa protein, which was cleaved to produce N-terminal fragments of 35-40 kDa and C-terminal fragments of 19-23 kDa. The Volga German (N141I) mutation did not cause any significant change in the metabolites of PS2. COS-1 cells doubly transfected with cDNAs for N141I mutant PS2 and human beta-amyloid precursor protein (beta APP) or a C-terminal fragment thereof, as well as mouse Neuro2a neuroblastoma cells stably transfected with N141I mutant PS2 alone, secreted 1.5-10-fold more A beta. residues 42 (or 43) [A beta 42(43)] compared with those expressing the wild-type PS2. Apparently, the PS2 mutation (N141I) linked to FAD alters the metabolite of A beta/beta APP to foster the production of the form of A beta that most readily deposits in amyloid plaques. Thus, mutant PS2 may lead to AD by altering the metabolite of A beta/beta APP.	=> e stern david/au E1 1 STERN DARRYL/AU E2 4 STERN DARRYL A/AU E3 162 -> STERN DAVID/AU E4 9 STERN DAVID A/AU E5 84 STERN DAVID B/AU E6 1 STERN DAVID BENJAMIN/AU E7 2 STERN DAVID E/AU E8 71 STERN DAVID F/AU E9 1 STERN DAVID FREDERICK/AU E10 2 STERN DAVID H/AU E11 4 STERN DAVID I/AU E12 40 STERN DAVID L/AU => s e3 L13 162 "STERN DAVID"/AU => s l13 and presenilin7/ab,bi 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE 'AB' IS NOT A VALID FIELD CODE L14 0 L13 AND PRESENILIN7/AB,BI

=> e wolozin b/au

E1 1 WOLOZEN J J JR/AU  
E2 1 WOLOZIA B/AU  
E3 84 -> WOLOZIN B/AU  
E4 1 WOLOZIN B I/AU  
E5 1 WOLOZIN B I/AU  
E6 70 WOLOZIN B L/AU  
E7 3 WOLOZIN BEN/AU  
E8 1 WOLOZIN BENJAMIN L/AU  
E9 25 WOLOZIN BENJAMIN/AU  
E10 16 WOLOZIN BENJAMIN L/AU  
E11 3 WOLOZIN M W/AU  
E12 2 WOLOZIN R/AU

=> s e3-e10

L17 201 ("WOLOZIN B"/AU OR "WOLOZIN B I"/AU OR  
"WOLOZIN B J"/AU OR "WOLO  
ZIN B L"/AU OR "WOLOZIN BEN"/AU OR "WOLOZIN  
BENJAMIN L"/AU OR  
"WOLOZIN BENJAMIN"/AU OR "WOLOZIN  
BENJAMIN L"/AU)

=> s I17 and presenilin/ab,bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L18 23 L17 AND PRESENILIN/AB,BI

=> dup rem I18

PROCESSING COMPLETED FOR L18  
L19 12 DUP REM L18 (11 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 12 ANSWERS -  
CONTINUE? Y/(N)y

L19 ANSWER 1 OF 12 CAPLUS COPYRIGHT 1999 ACS  
AN 1998:258266 CAPLUS  
DN 129:52570  
TI Regulation of apoptosis by \*\*\*presenilin\*\*\* 1  
AU \*\*\*Wolozin, B \*\*\* ; Alexander, P.; Palacino, J  
CS Department of Pharmacology, Loyola University Medical Center,  
Maywood, IL,  
60153, USA  
SO Neurobiol. Aging (1998), 19(Suppl. 1), Proceedings of the 11th  
Annual Tokyo  
Institute of Psychiatry International Symposium, 1997), S23-S27  
CODEN: NEAGDO; ISSN: 0197-4580  
PB Elsevier Science Inc.  
DT Journal

LA English  
AB Familial Alzheimer's disease is transmitted as an autosomal  
dominant  
disorder and, in 5-10% of the cases, is caused by mutations in the  
coding  
regions of two homologous genes. \*\*\*Presenilin\*\*\* 1 and 2  
(P51 and  
PS2). Previously, we have shown that PS2, a homolog of P51,  
regulates  
apoptosis induced in neurons by trophic withdrawal or A.beta., and  
in  
T-cells by Fas ligand. We now report that PS1 also regulates  
apoptosis.  
Both wild-type and the H115Y mutant form of PS1 enhance  
Fas-mediated  
apoptosis in Jurkat cells. We also obsd. that wild-type and the  
mutant form of PS1 differentially regulate Jun Kinase, an important  
enzyme  
regulating apoptosis.

L19 ANSWER 2 OF 12 MEDLINE DUPLICATE  
1  
AN 1998356210 MEDLINE  
DN 98356210  
TI \*\*\*Presenilin\*\*\* 1 associates with glycogen synthase  
kinase-3beta and  
its substrate tau.  
AU Takashima A; Murayama M; Murayama O; Kohno T; Honda T;  
Yasuike K,  
Nihonmatsu N; Mercken M; Yamaguchi H; Sugihara S;  
\*\*\*Wolozin B\*\*\*  
CS Laboratory for Alzheimer's Disease, Brain Science Institute,  
RIKEN, 2-1  
Hiroawa, Wako-shi, Saitama 350-01, Japan.  
SO PROCEEDINGS OF THE NATIONAL ACADEMY OF  
SCIENCES OF THE UNITED STATES OF  
AMERICA, (1998 Aug 4) 95 (16) 9637-41.  
Journal code: PV3. ISSN: 0027-8424.

CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199811  
EW 19981102  
AB Families bearing mutations in the \*\*\*presenilin\*\*\* 1 (PS1)  
gene  
develop Alzheimer's disease. Previous studies have shown that the  
Alzheimer-associated mutations in PS1 increase production of  
amyloid beta  
protein (Abeta1-42). We now show that PS1 also regulates  
phosphorylation  
of the microtubule-associated protein tau. PS1 directly binds tau  
and a  
tau kinase, glycogen synthase kinase 3beta (GSK-3beta). Deletion  
studies  
show that both tau and GSK-3beta bind to the same region of PS1,

residues  
250-298, whereas the binding domain on tau is the  
microtubule-binding  
repeat region. The ability of PS1 to bring tau and GSK-3beta into  
close  
proximity suggests that PS1 may regulate the interaction of tau with  
GSK-3beta. Mutations in PS1 that cause Alzheimer's disease  
increase the  
ability of PS1 to bind GSK-3beta and, correspondingly, increase its  
tau-directed kinase activity. We propose that the increased  
association of  
GSK-3beta with mutant PS1 leads to increased phosphorylation of  
tau.

L19 ANSWER 3 OF 12 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1999:67066 BIOSIS  
DN PREV19990067066  
TI Association of \*\*\*presenilin\*\*\* 1 with beta-catenin.  
AU Takashima, A. (1); Murayama, M. (1); Murayama, O. (1);  
Honda, T. (1);  
Palacino, James; \*\*\*Wolozin, Benjamin\*\*\*  
CS (1) Lab. Alzheimer's Dis., RIKEN, BSI, 2-1 Hiroawa, Wako-shi,  
Saitama  
350-01 Japan  
SO Society for Neuroscience Abstracts, (1998) Vol. 24, No. 1-2, pp.  
758.  
Meeting Info.: 28th Annual Meeting of the Society for  
Neuroscience, Part 1  
Los Angeles, California, USA November 7-12, 1998 Society for  
Neuroscience  
. ISSN: 0190-5295.  
DT Conference  
LA English

L19 ANSWER 4 OF 12 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1999:48375 BIOSIS  
DN PREV19990048375  
TI \*\*\*Presenilin\*\*\* 1 regulates stimulated cleavage of the  
amyloid  
precursor protein.  
AU Palacino, J. J. (1); Berechid, B.; Alexander, P. (1); Nye, J.;  
\*\*\*Wolozin, B. (1)\*\*\*  
CS (1) Dep. Pharmacol., Loyola Univ. Chicago, Maywood, IL 60153  
USA  
SO Society for Neuroscience Abstracts, (1998) Vol. 24, No. 1-2, pp.  
470.  
Meeting Info.: 28th Annual Meeting of the Society for  
Neuroscience, Part 1  
Los Angeles, California, USA November 7-12, 1998 Society for  
Neuroscience  
. ISSN: 0190-5295.  
DT Conference  
LA English

L19 ANSWER 5 OF 12 MEDLINE DUPLICATE  
2  
AN 1998409316 MEDLINE

DN 98409316  
 TI Direct association of \*\*\*presenilin\*\*\* -1 with beta-catenin.  
 AU Murayama M; Tanaka S; Palacino J; Murayama O; Honda T; Sun X; Yasutake K;  
 Nihonmatsu N; \*\*\*Wolozin B\*\*\* ; Takashima A  
 CS Laboratory for Alzheimer's Disease, Brain Science Institute, RIKEN, Saitama, Japan.  
 SO FEBS LETTERS, (1998 Aug 14) 433 (1-2) 73-7.  
 Journal code: EUH. ISSN: 0014-5793.  
 CY Netherlands  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals; Cancer Journals  
 EM 199812  
 EW 19981201  
 AB Families bearing mutations in the \*\*\*presenilin\*\*\* -1 (PS1) gene develop Alzheimer's disease (AD). However, the mechanism through which PS1 causes AD is unclear. The co-immunoprecipitation with PS1 in transfected COS-7 cells indicates that PS1 directly interacts with endogenous beta-catenin, and the interaction requires residues 3224-50 of PS1 and 445-676 of beta-catenin. Both proteins are co-localized in the endoplasmic reticulum. Over-expression of PS1 reduces the level of cytoplasmic beta-catenin, and inhibits beta-catenin-F cell factor-regulated transcription. These results indicate that PS1 plays a role as inhibitor of the beta-catenin signal, which may be connected with the AD dysfunction.

L19 ANSWER 6 OF 12 MEDLINE DUPLICATE  
 3  
 AN 1998220955 MEDLINE  
 DN 98220955  
 TI Regulation of apoptosis by \*\*\*presenilin\*\*\* 1.  
 AU \*\*\*Wolozin B\*\*\* ; Alexander P; Palacino J  
 CS Department of Pharmacology, Loyola University Medical Center, Maywood, IL 60153, USA.; bwolozin@wvpo.it.luc.edu  
 SO NEUROBIOLOGY OF AGING, (1998 Jan-Feb) 19 (1 Suppl) S23-7.  
 Journal code: NX5. ISSN: 0197-4580.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199808  
 EW 19980803  
 AB Familial Alzheimer's disease is transmitted as an autosomal dominant disorder and, in 5-10% of the cases, is caused by mutations in the coding regions of two homologous genes, \*\*\*Presenilin\*\*\* 1 and 2

(PS1 and PS2). Previously, we have shown that PS2, a homolog of PS1, regulates apoptosis induced in neurons by trophic withdrawal or Abeta, and in T-cells by Fas ligand. We now report that PS1 also regulates apoptosis. Both wild-type and the H115Y mutant form of PS1 enhance Fas-mediated apoptosis in Jurkat cells. We also observed that wild-type and the H115Y mutant form of PS1 differentially regulate Jun Kinase, an important enzyme regulating apoptosis.

L19 ANSWER 7 OF 12 EMBASE COPYRIGHT 1999 ELSEVIER SCI. B.V.  
 AN 1998132311 EMBASE  
 TI Regulation of apoptosis by \*\*\*presenilin\*\*\* 1.  
 AU \*\*\*Wolozin B\*\*\* ; Alexander P; Palacino J.  
 CS B. Wolozin, Department of Pharmacology, Loyola University Medical Center, Building 102, 2160 South First Avenue, Maywood, IL 60153, United States.  
 bwolozin@wvpo.it.luc.edu  
 SO Neurobiology of Aging, (1998) 19(SUPPL. 1 (S23-S27)). Refs: 27  
 ISSN: 0197-4580 CODEN: NEAGDO  
 PUI S01974580980000414  
 CY United States  
 DT Journal; Conference Article  
 FS 008 Neurology and Neurosurgery  
 020 Gerontology and Geriatrics  
 021 Developmental Biology and Teratology  
 022 Human Genetics  
 032 Psychiatry  
 LA English  
 SL English  
 AB Familial Alzheimer's disease is transmitted as an autosomal dominant disorder and, in 5-10% of the cases, is caused by mutations in the coding regions of two homologous genes. \*\*\*Presenilin\*\*\* 1 and 2 (PS1 and PS2). Previously, we have shown that PS2, a homolog of PS1, regulates apoptosis induced in neurons by trophic withdrawal or A.beta., and in T-cells by Fas ligand. We now report that PS1 also regulates apoptosis. Both wild-type and the H115Y mutant form of PS1 enhance Fas-mediated apoptosis in Jurkat cells. We also observed that wild-type and the H115Y mutant form of PS1 differentially regulate Jun Kinase, an important enzyme regulating apoptosis.

L19 ANSWER 8 OF 12 BIOSIS COPYRIGHT 1999 BIOSIS AN 1997:527282 BIOSIS  
 DN PREV199799826485  
 TI Mutant PS1 stimulates the JNK signal transduction cascade.  
 AU Palacino, J. J.; Alexander, P.; St. George-Hyslop, P.; Takashima, A.; Schultz, R.; \*\*\*Wolozin, B.\*\*\*  
 CS Dep. Pharmacol., Loyola Univ. Chicago Med. Cent., Maywood, IL 60153 USA  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 1117.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295.  
 DT Conference; Abstract; Conference  
 LA English  
 L19 ANSWER 9 OF 12 BIOSIS COPYRIGHT 1999 BIOSIS AN 1997:527284 BIOSIS  
 DN PREV199799826487  
 TI \*\*\*Presenilin\*\*\* -2 couples with the signal transduction system of the RAGE receptor.  
 AU \*\*\*Wolozin, B.\*\*\* ; Alexander, P.; Stern, D.; Schmidt, A. M.; Yan, S. D.  
 CS Dep. Pharmacol., Loyola Univ. Chicago Med. Cent., Maywood, IL 60153 USA  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 1117.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295.  
 DT Conference; Abstract; Conference  
 LA English  
 L19 ANSWER 10 OF 12 MEDLINE DUPLICATE  
 4  
 AN 97094860 MEDLINE  
 DN 97094860  
 TI Requirement of the familial Alzheimer's disease gene PS2 for apoptosis.  
 Oposing effect of ALG-3.  
 AU Vito P; \*\*\*Wolozin B\*\*\* ; Ganjei J K; Iwasaki K; Lacana E; D'Adamo L  
 CS T-Cell Molecular Biology Unit, Laboratory of Cellular and Molecular Immunology, NIAID, National Institutes of Health, Maryland 20892, USA.; lldadamo@atlas.niaid.nih.gov  
 SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1996 Dec 6) 271 (49) 31025-8.  
 Journal code: HIV. ISSN: 0021-9258.  
 CY United States

DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
OS GENBANK-U49111; GENBANK-U57324;  
GENBANK-U57325  
EM 199703  
AB ALG-3, a truncated mouse homologue of the chromosome 1  
familial Alzheimer's disease gene PS2, rescues T hybridoma 3DO cells  
from T-cell  
receptor-induced apoptosis by inhibiting Fas ligand induction and  
Fas  
signaling. Here we show that ALG-3 transfected 3DO cells express  
a  
COOH-terminal PS2 polypeptide. Overexpression of PS2 in  
ALG-3 transfected  
3DO cells reconstitutes sensitivity to receptor-induced cell death,  
suggesting that the artificial PS2 polypeptide functions as a  
dominant  
negative mutant of PS2. ALG-3 and antisense PS2 protect PC12  
cells from  
glutamate-induced apoptosis but not from death induced by  
hydrogen  
peroxide or the free radical MPP+. Thus, the PS2 gene is required  
for some  
forms of cell death in diverse cell types, and its function is opposed  
by  
ALG-3.

L19 ANSWER 11 OF 12 MEDLINE DUPLICATE  
5  
AN 97094374 MEDLINE  
DN 97094374  
TI Participation of \*\*\*presenilin\*\*\* 2 in apoptosis: enhanced  
basal  
activity conferred by an Alzheimer mutation.  
AU \*\*\*Wolozin B\*\*\* ; Iwasaki K; Vito P; Ganjei J K; Lacan'a E;  
Sunderland  
T; Zhao B; Kuslak J W; Wasco W; D'Adamo L  
CS Unit on Alzheimer Biology, Laboratory of Clinical Science,  
National  
Institute of Mental Health, Building 10, Room 3D41, 9000  
Rockville Pike,  
Bethesda, MD 20892, USA.. ldamio@atlas.nimh.nih.gov  
SO SCIENCE, (1996 Dec 6) 274 (5293) 1710-3.  
Journal code: UJ7. ISSN: 0036-8075.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199703  
AB Overexpression of the familial Alzheimer's disease gene  
\*\*\*Presenilin\*\*\*  
2 (PS2) in nerve growth factor-differentiated PC12 cells increased  
apoptosis induced by trophic factor withdrawal or beta-amyloid.  
Transfection of antisense PS2 conferred protection against  
apoptosis

induced by trophic withdrawal in nerve growth factor-differentiated  
or  
amyloid precursor protein-expressing PC12 cells. The apoptotic  
cell death  
induced by PS2 protein was sensitive to pertussis toxin, suggesting  
that  
heterotrimeric GTP-binding proteins are involved. A PS2 mutation  
associated with familial Alzheimer's disease was found to generate  
a  
molecule with enhanced basal apoptotic activity. This gain of  
function  
might accelerate the process of neurodegeneration that occurs in  
Alzheimer's disease, leading to the earlier age of onset  
characteristic of  
familial Alzheimer's disease.

L19 ANSWER 12 OF 12 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1996:489438 BIOSIS  
DN PREV199699211794  
TI PS2 participates in cellular apoptosis.  
AU \*\*\*Wolozin, B. (1)\*\*\* ; Vito, P.; Ganjei, K.; Iwasaki, K.;  
Lacana, E.;  
D'Adamo, L.  
CS (1) Section Geriatric Psychiatry, NIMH, NIAID, Bethesda, MD  
20892 USA  
SO Society for Neuroscience Abstracts, (1996) Vol. 22, No. 1-3, pp.  
729.  
Meeting Info.: 26th Annual Meeting of the Society for  
Neuroscience  
Washington, D.C., USA November 16-21, 1996  
ISSN: 0190-5295.  
DT Conference  
LA English  
=> e stern d/au  
E1 1 STERN CURTIS/AU  
E2 11 STERN CURTIS H/AU  
E3 527 -> STERN D/AU  
E4 41 STERN D A/AU  
E5 100 STERN D B/AU  
E6 148 STERN D F/AU  
E7 32 STERN D H/AU  
E8 2 STERN D I/AU  
E9 42 STERN D J/AU  
E10 48 STERN D L/AU  
E11 246 STERN D M/AU  
E12 69 STERN D N/AU  
=> s e3  
L20 527 \*STERN D\*/AU  
=> s l20 and presenilin/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L21 1 L20 AND PRESENILIN/AB,BI  
=> d  
L21 ANSWER 1 OF 1 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1997:527284 BIOSIS  
DN PREV19979826487  
TI \*\*\*Presenilin\*\*\* -2 couples with the signal transduction system  
of the  
RAGE receptor.  
AU Wolozin, B.; Alexander, P.; \*\*\*Stern, D.\*\*\* ; Schmidt, A.  
M.; Yan, S.  
D.  
CS Dep. Pharmacol., Loyola Univ. Chicago Med. Cent., Maywood,  
IL 60153 USA  
SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp.  
1117.  
Meeting Info.: 27th Annual Meeting of the Society for  
Neuroscience New  
Orleans, Louisiana, USA October 25-30, 1997  
ISSN: 0190-5295.  
DT Conference; Abstract; Conference  
LA English  
=> s l9  
'AB' IS NOT A VALID FIELD CODE  
1 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
2 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L22 56 L9  
=> s l22 and neuron7/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L23 5 L22 AND NEURON7/AB,BI  
=> dup rem l23  
PROCESSING COMPLETED FOR L23  
L24 5 DUP REM L23 (0 DUPLICATES REMOVED)  
=> d l- bib ab  
YOU HAVE REQUESTED DATA FROM 5 ANSWERS -  
CONTINUE? Y(N)y

L24 ANSWER 1 OF 5 CAPLUS COPYRIGHT 1999 ACS  
AN 1998:643914 CAPLUS  
DN 130:50786  
TI RAGE-A beta. interactions in the pathophysiology of Alzheimer's disease  
AU Yan, Shi Du; Stern, David; Kane, Michael D.; Kuo, Yu-Min; Lampert, Heather  
C.; Rohrer, Alex E.  
CS Department of Pathology, Surgery, Medicine and Physiology, College of Physicians and Surgeons, Columbia University, New York, NY, 10032, USA  
SO Restor. Neurol. Neurosci. (1998), 12(2,3), 167-173  
CODEN: RNNEEL, ISSN: 0922-6028  
PB IOS Press  
DT Journal  
LA English  
AB RAGE is a cell surface mol. primarily identified for its capacity to bind advanced glycation end-products and amphoterin. Immunocytochem. studies demonstrated that in Alzheimer's disease (AD) the expression of RAGE is elevated in \*\*\*neurons\*\*\* close to neuritic plaque beta-amyloid (A beta) deposits and in the cells of A beta. cong. vessels. Crosslinking of surface bound A beta. 1-40 to endothelial cells, yielded a band of 50 kDa identified as RAGE. Using the sol. extracellular domain of recombinant human RAGE, we found that A beta. binds to RAGE with a Kd = 57 nM, a value close to those found for mouse brain endothelial cells and rat cortical \*\*\*neurons\*\*\*. The interaction of A beta. with RAGE in \*\*\*neuronal\*\*\*, endothelial, and RAGE-transfected COS-1 cells induced oxidative stress, as assessed by the TBARS and MTT assays. ELISA demonstrated a 2.5 times increase of RAGE in AD over control brains. Activated microglia also showed elevated expression of RAGE. In the BV-2 microglial cell line, RAGE bound A beta. in a dose dependent manner with a Kd of 25 nM. Sol. A beta. induced the migration of microglia along a concn. gradient, while immobilized A beta. arrested this migration. A beta.-RAGE interaction also activated NF-kappa B, resulting in \*\*\*neuronal\*\*\* up-regulation of macrophage-colony stimulating factor (M-CSF) which also induced microglial migration. Apparently, RAGE-A beta. interactions play an important role in the pathophysiol. of Alzheimer's disease.

L24 ANSWER 2 OF 5 CAPLUS COPYRIGHT 1999 ACS  
AN 1997:525836 CAPLUS  
DN 127:204001  
TI Binding of beta-amyloid protein by an \*\*\*advanced\*\*\*  
\*\*\*glycation\*\*\* - \*\*\*end\*\*\* - \*\*\*product\*\*\*  
possible treatment of Alzheimer's disease  
IN Stern, David; Schmidt, Ann Marie; Yan, Shi Du  
PA Trustees of Columbia University, USA  
SO PCT Int. Appl., 91 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN/CNT 1  
PATENT NO. KIND DATE APPLICATION NO.  
DATE  
PI WO 9726913 A1 19970731 WO 97-US857 19970121  
W: AU, CA, JP, MX  
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE  
AU 9718327 A1 19970820 AU 97-18327 19970121  
PRAIUS 96-592070 19960126  
WO 97-US857 19970121  
AB The beta-amyloid protein binds to a cell-surface RAGE (receptor for advanced glycation end products) in neural cells and induces neurotoxic damage typical of Alzheimer's disease. This interaction may be a useful target for treatment of Alzheimer's disease. Binding assays for the identification and characterization of beta-amyloid-binding proteins used to identify the interaction of beta-amyloid with RAGE are described. Peptides capable of inhibiting the interaction are reported.

L24 ANSWER 3 OF 5 CAPLUS COPYRIGHT 1999 ACS  
AN 1997:544866 CAPLUS  
DN 127:201264  
TI Beta amyloid toxicity does not require RAGE protein  
AU Liu, Y.; Dargusch, R.; Schubert, D.  
CS The Salk Institute for Biological Studies, La Jolla, CA, 92037, USA  
SO Biochem. Biophys. Res. Commun. (1997), 237(1), 37-40  
CODEN: BBRC9A; ISSN: 0006-291X  
PB Academic  
DT Journal  
LA English  
AB It has been suggested that a receptor for advanced glycation end products (RAGE) is the nerve cell receptor for amyloid beta protein (A beta). To det. if this is indeed the case, two neural cell lines as well as rat cortical \*\*\*neurons\*\*\* were examd. for the presence of the mRNA for RAGE by PCR and northern blot anal. Although lung was strongly

pos., in no case was RAGE mRNA detected in the cultured neural cells.  
Glycated albumin is a major ligand for RAGE and the cell surface RAGE protein is trypsin sensitive. In agreement with the mRNA data, trypsin treatment did not alter A beta. toxicity, nor did glycated albumin modify the A beta. response. It follows that RAGE is not the neural receptor for A beta..

L24 ANSWER 4 OF 5 CAPLUS COPYRIGHT 1999 ACS  
AN 1995:898715 CAPLUS  
DN 124:26311  
TI The receptor for advanced glycation end products (RAGE) is a cellular binding site for amphoterin. Mediation of neurite outgrowth and co-expression of RAGE and amphoterin in the developing nervous system  
AU Hori, Osamu; Brett, Jerold; Slaterry, Timothy; Cao, Rong; Zhang, Jinghua; Chen, Jing Xian; Nagashima, Mariko; Lundh, Erik R.; Vijay, Sharmila, et al.  
CS Coll. Physicians Surg., Columbia Univ., New York, NY, 10032, USA  
SO J. Biol. Chem. (1995), 270(43), 25752-61  
CODEN: JBCHA3; ISSN: 0021-9258  
DT Journal  
LA English  
AB The receptor of advanced glycation end products (RAGE), a newly-identified member of the Ig superfamily, mediates interactions of advanced glycation end product (AGE)-modified proteins with endothelium and other cell types. Survey of normal tissues demonstrated RAGE expression in situations in which accumulation of AGEs would be unexpected, leading to the hypothesis that under physiol. circumstances, RAGE might mediate interaction with ligands distinct from AGEs. Sequential chromatog. of bovine lung ext. identified polypeptides with Mr values of .apprxq. 12,000 (p12) and .apprxq. 23,000 (p23) which bound RAGE. NH2-terminal and internal protein sequence data for p23 matched that reported previously for amphoterin. Amphoterin purified from rat brain or recombinant rat amphoterin bound to purified sRAGE in a saturable and dose-dependent manner, blocked by anti-RAGE IgG or a sol. form of RAGE (sRAGE). Cultured embryonic rat

\*\*\*neurons\*\*\*, which express RAGE, displayed dose-dependent binding of 125I-amphoterin which was prevented by blockade of RAGE using antibody to the receptor or excess sol. receptor (sRAGE). A functional correlate of RAGE-amphoterin interaction was inhibition by anti-RAGE F(ab)2 of neurite formation by cortical \*\*\*neurons\*\*\* specifically on amphoterin-coated substrates. Consistent with a potential role for RAGE-amphoterin interaction in development, amphoterin and RAGE mRNA/antigen were co-localized in developing rat brain. These data indicate that RAGE has physiol. relevant ligands distinct from AGEs which are likely, via their interaction with the receptor, to participate the physiol processes outside of the context of diabetes and accumulation of AGEs.

L24 ANSWER 5 OF 5 CAPLUS COPYRIGHT 1999 ACS  
 AN 1994:240942 CAPLUS  
 DN 120:240942  
 TI Survey of the distribution of a newly characterized receptor for advanced glycation end products in tissues  
 AU Brett, Jerold; Schmidt, Ann Marie; Yan, Shi Du; Zou, Yu Shan; Weidman, Elliott; Pinsky, David; Nowygrod, Roman; Neeper, Michael; Przysiecki, Craig; et al.  
 CS Coll. Physicians Surg., Columbia Univ., New York, NY, 10032, USA  
 SO Am. J. Pathol. (1993), 143(6), 1699-712  
 LA English  
 DT Journal  
 AB Advanced glycation end products (AGEs), the final products of non-enzymic glycation and oxidn. of proteins, are found in the plasma and accumulate in the tissues during aging and at an accelerated rate in diabetes. A novel integral membrane protein, termed receptor for AGE (RAGE), forms a central part of the cell surface binding site for AGEs. Using monospecific, polyclonal antibody raised to human recombinant and bovine RAGE, immunostaining of bovine tissues showed RAGE in the vasculature, endothelium, and smooth muscle cells and in mononuclear cells in the tissues. Consistent with these data, RAGE antigen and mRNA were identified in cultured bovine endothelium, vascular smooth muscle, and monocyte-derived macrophages. RAGE antigen was also

visualized in bovine cardiac myocytes as well as in cultures of neonatal rat cardiac myocytes and in neural tissue where motor \*\*\*neurons\*\*\*, peripheral nerves, and a population of cortical \*\*\*neurons\*\*\* were pos. In situ hybridization confirmed the presence of RAGE mRNA in the tissues, and studies with rat PC12 pheochromocytes indicated that they provide a \*\*\*neuronal\*\*\* -related cell culture model for examg. RAGE expression. Pathol. studies of human atherosclerotic plaques showed infiltration of RAGE-expressing cells in the expanded intima. These results indicate that RAGE is present in multiple tissues and suggest the potential relevance of AGE-RAGE interactions for modulating properties of the vasculature as well as neural and cardiac function, prominent areas of involvement in diabetes and in the normal aging process.

=> s l9 and glial/ab,bi  
 'AB' IS NOT A VALID FIELD CODE  
 'AB' IS NOT A VALID FIELD CODE  
 2 FILES SEARCHED...  
 'AB' IS NOT A VALID FIELD CODE  
 'AB' IS NOT A VALID FIELD CODE  
 'AB' IS NOT A VALID FIELD CODE  
 L25 1 L9 AND GLIAL/AB,BI

=> d  
 L25 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS  
 AN 1997:525836 CAPLUS  
 DN 127:204001  
 TI Binding of beta-amyloid protein by an \*\*\*advanced\*\*\*  
 \*\*\*glycation\*\*\* \*\*\*end\*\*\* - \*\*\*product\*\*\*  
 \*\*\*receptor\*\*\* and possible treatment of Alzheimer's disease  
 IN Stern, David; Schmidt, Ann Marie; Yan, Shi Du  
 PA Trustees of Columbia University, USA  
 SO PCT Int. Appl., 91 pp.  
 DT Patent  
 LA English  
 FAN CNT 1  
 PATENT NO. KIND DATE APPLICATION NO.  
 DATE

PI WO 9726913 A1 19970731 WO 97-US857 19970121  
 W: AU, CA, JP, MX  
 RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

AU 9718327 A1 19970820 AU 97-18327 19970121  
 PRAI US 96-592070 19960126  
 WO 97-US857 19970121

=> d ab  
 L25 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS  
 AB The beta-amyloid protein binds to a cell-surface RAGE (receptor for advanced glycation end products) in neural cells and induces neurotoxic damage typical of Alzheimer's disease. This interaction may be a useful target for treatment of Alzheimer's disease. Binding assays for the identification and characterization of beta-amyloid-binding proteins used to identify the interaction of beta-amyloid with RAGE are described. Peptides capable of inhibiting the interaction are reported.

=> s l9 and microglial/ab,bi  
 'AB' IS NOT A VALID FIELD CODE  
 'AB' IS NOT A VALID FIELD CODE  
 2 FILES SEARCHED...  
 'AB' IS NOT A VALID FIELD CODE  
 'AB' IS NOT A VALID FIELD CODE  
 'AB' IS NOT A VALID FIELD CODE  
 L26 1 L9 AND MICROGLIAL/AB,BI

=> d  
 L26 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS  
 AN 1998:643914 CAPLUS  
 DN 130:50786  
 TI RAGE-A.beta. interactions in the pathophysiology of Alzheimer's disease  
 AU Yan, Shi Du; Stern, David; Kane, Michael D.; Kuo, Yu-Min; Lampert, Heather C.; Roher, Alex E.  
 CS Department of Pathology, Surgery, Medicine and Physiology, College of Physicians and Surgeons, Columbia University, New York, NY, 10032, USA  
 SO Restor. Neurol. Neurosci. (1998), 12(2,3), 167-173  
 CODEN: RNNEEL; ISSN: 0922-6028  
 PB IOS Press  
 DT Journal  
 LA English  
 => d ab

L26 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS  
 AB RAGE is a cell surface mol. primarily identified for its capacity to bind

advanced glycation end-products and amphoterin.  
Immunocytochem. studies  
demonstrated that in Alzheimer's disease (AD) the expression of RAGE is elevated in neurons close to neuritic plaque beta-amyloid (A. beta.) deposits and in the cells of A. beta. contig. vessels. Crosslinking of surface bound A. beta. 1-40 to endothelial cells, yielded a band of 50 kDa identified as RAGE. Using the sol. extracellular domain of recombinant human RAGE, we found that A. beta. binds to RAGE with a Kd = 57 nM, a value close to those found for mouse brain endothelial cells and rat cortical neurons. The interaction of A. beta. with RAGE in neuronal, endothelial, and RAGE-transfected COS-1 cells induced oxidative stress, as assessed by the TBARS and MTT assays. ELISA demonstrated a 2.3 times increase of RAGE in AD over control brains. Activated microglia also showed elevated expression of RAGE. In the BV-2 \*\*\*microglial\*\*\* cell line, RAGE bound A. beta. in a dose dependent manner with a Kd of 25 nM. Sol. A. beta. induced the migration of microglia along a concn. gradient, while immobilized A. beta. arrested this migration. A. beta.-RAGE interaction also activated NF- $\kappa$ B, resulting in neuronal up-regulation of macrophage-colony stimulating factor (M-CSF) which also induced \*\*\*microglial\*\*\* migration. Apparently, RAGE-A. beta. interactions play an important role in the pathophysiol. of Alzheimer's disease.

=> s 19 and astrocyte#ab,bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
2 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L27 1 L9 AND ASTROCYTE#AB,BI

=> d

L27 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1999 ACS  
AN 1998:760707 CAPLUS  
DN 130:107885  
TI Cell activation by glycated proteins. AGE receptors, receptor recognition factors and functional classification of AGEs  
AU Thormalley, Paul J.  
CS Department of Biological Sciences, University of Essex, Essex, CO4 3SQ, UK  
SO Cell. Mol. Biol. (Paris) (1998), 44(7), 1013-1023

CODEN: CMOBEF; ISSN: 0145-5680  
PB C.M.B. Association  
DT Journal; General Review  
LA English

=> s 19 and endothelial/ab,bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
2 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L28 18 L9 AND ENDOTHELIAL/AB,BI

=> dup rem 128

PROCESSING COMPLETED FOR L28  
L29 15 DUP REM L28 (3 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 15 ANSWERS -  
CONTINUE? Y(N)Y

L29 ANSWER 1 OF 15 CAPLUS COPYRIGHT 1999 ACS  
AN 1999:182259 CAPLUS  
TI Characterization of the \*\*\*Advanced\*\*\* \*\*\*Glycation\*\*\*  
\*\*\*End\*\*\*  
- \*\*\*Product\*\*\* \*\*\*Receptor\*\*\* Complex in Human Vascular  
\*\*\*Endothelial\*\*\* Cells  
AU Sutt, Alan W.; He, Cijiang; Vlassara, Helen  
CS Department of Ophthalmology, Royal Victoria Hospital, Queen's University of Belfast, Belfast, BT12 6BA, UK  
SO Biochem. Biophys. Res. Commun. (1999), 256(3), 549-556  
CODEN: BBRC99; ISSN: 0006-291X  
PB Academic Press  
DT Journal  
LA English  
AB Advanced glycation end products (AGEs) have been implicated as causal factors in the vascular complications of diabetes and it is known that these products interact with cells through specific receptors. The AGE-receptor complex, originally described as p60 and p90, has been characterised in hemopoietic cells and the component proteins identified and designated AGE-R1, -R2 and -R3. In the current study we have characterised this receptor in human umbilical vein \*\*\*endothelial\*\*\* cells (HUVECs) and elucidated several important biol. properties which may impact on AGE mediated vascular disease. 125I-AGE-BSA binding

to HUVEC monolayers was detd. with and without various cold competitors. The synthetic AGE, 2-(2-furyl)-4-(furyl)-1H-imidazole (FFI)-BSA, failed to compete with AGE-BSA binding unlike observations already reported in hemopoietic cells. The ability of 125I-AGE-BSA to bind to sepd. HUVEC plasma membrane (PM) proteins was also examd. and the binding at specific bands inhibited by antibodies to each component of the AGE-receptor complex. Western blotting of whole cell and PM fractions, before and after exposure to AGE-BSA, revealed that AGE-R1, -R2 and -R3 are subject to upregulation upon exposure to their ligand, a phenomenon which was also demonstrated by immunofluorescence of non-permeabilised cells. mRNA expression of each AGE-receptor component was apparent in HUVECs, with the AGE-R2 and -R3 gene expression being upregulated upon exposure to AGEs in a time-dependent manner. A phosphorylation assay in combination with AGE-R2 immunopptn. demonstrated that this component of the receptor complex is phosphorylated by acute exposure to AGE-BSA. These results indicate the presence of a conserved AGE-receptor complex in endothelium which demonstrates subtle differences to other cell-types. In response to AGE-modified mols., this complex is subject to upregulation, while the AGE-R2 component also displays increased phosphorylation possibly leading to enhanced signal transduction. (c) 1999 Academic Press.  
L29 ANSWER 2 OF 15 CAPLUS COPYRIGHT 1999 ACS  
AN 1998:760707 CAPLUS  
DN 130:107885  
TI Cell activation by glycated proteins. AGE receptors, receptor recognition factors and functional classification of AGEs  
AU Thormalley, Paul J.  
CS Department of Biological Sciences, University of Essex, Essex, CO4 3SQ, UK  
SO Cell. Mol. Biol. (Paris) (1998), 44(7), 1013-1023  
CODEN: CMOBEF; ISSN: 0145-5680  
PB C.M.B. Association  
DT Journal; General Review  
LA English



AB A review, with approx. 72 refs. Proteins modified by advanced glycation end products (AGE) bind to cell surface receptors and other AGE binding proteins. AGE-binding receptors are: scavenger receptors types I and II, the receptor for advanced glycation end products (RAGE), oligosaccharyl transferase-48 (OST-48, AGE-R1), 80K-H phosphoprotein (AGE-R2) and galectin-3 (AGE-R3). AGE receptors are found in monocytes, macrophages, \*\*\*endothelial\*\*\* cells, pericytes, podocytes and astrocytes and microglia. AGE-modified proteins also bind to lysozyme and lactoferrin. A review of the evidence for receptors binding AGE-modified protein binding in vivo is presented. Scavenger receptors have only been shown to bind proteins modified by AGE to a much higher extent than found in vivo. 80K-H phosphoprotein is involved in FGFR3 signal transduction to MAP kinase, and may be involved in AGE-receptor signal transduction. Whether all of these proteins bind AGE-modified proteins in vivo is not yet clear. Cell activation in response to AGE-modified proteins is associated with increased expression of extracellular matrix proteins, vascular adhesion molecules, cytokines and growth factors. Depending on the cell type and concurrent signaling, this is associated with chemotaxis, angiogenesis, oxidative stress, cell proliferation or programmed cell death (PCD). Receptor recognition factors for agonism at the AGE receptor have been little studied but to date hydroimidazolones appear to be the most likely candidates. Pharmacol. inhibition of AGE receptor-mediated cell activation with specific antagonists may provide the basis for therapeutic intervention in diseases where AGE accumulation is a suspected etiol. factor vascular complications of diabetes, macrovascular disease, renal insufficiency and Alzheimer's disease.

L29 ANSWER 3 OF 15 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1998:345185 BIOSIS  
 DN PREV199800345185  
 TI Differential expression of receptors for advanced glycation end products on monocytes in patients with IDDM.  
 AU Festa, A.; Schmoelzer, B.; Schernthaner, G.; Menzel, E. J. (1)  
 CS (1) Dep. Immunol., Univ. Vienna, Borschkeg. 8A, A-1090

Vienna Austria  
 SO Diabetologia, (June, 1998) Vol. 41, No. 6, pp. 674-680.  
 ISSN: 0012-186X.  
 DT Article  
 LA English  
 AB Accelerated modification of proteins by glucose terminating in the formation of advanced glycation endproducts (AGEs) is one of the main pathogenetic mechanisms of diabetes-associated complications. One pathway by which AGEs may exert their effects is by interaction with specific receptors initially identified on macrophages, monocytes and \*\*\*endothelial\*\*\* cells. As AGE-induced autoocrine upregulation of AGE receptors has been observed in vitro, we hypothesized that AGE-binding might be enhanced in diabetic patients to compensate for the elevated levels of circulating AGEs. We therefore examined the expression of AGE-binding sites on peripheral monocytes, serum levels of AGEs and AGE-induced cytokine production in patients with insulin-dependent diabetes mellitus (IDDM) compared to age-matched, healthy control subjects. In patients, AGE-binding capacity was significantly increased and there was only one class of binding sites, as revealed by Scatchard analysis (1.8 X 10<sup>5</sup> vs 1.4 X 10<sup>5</sup> binding sites per cell). Affinity of binding was, however, similar (K<sub>a</sub> 1.5 X 10<sup>6</sup> vs 1.4 X 10<sup>6</sup> mol<sup>-1</sup>). Saturation of binding was reached at 2.0-3.0 μmol/l with AGE-bovine serum albumin (BSA) as ligand. In contrast, cytometry using fluorescein isothiocyanate-labelled AGE-proteins showed no saturability and reversibility of AGE-binding up to 80 μmol/l, indicating non-specific binding in this concentration range. Again, this non-specific binding was significantly higher in IDDM patients. In addition, we found much higher levels of circulating AGEs in patients as compared to controls and studied possible functional consequences of increased AGE binding in vitro, monocyte stimulation by AGEs triggering cytokine release to a similar extent in patients and controls, i.e. independently of the AGE-binding capacity. Our finding of an enhanced overall AGE-binding capacity of peripheral monocytes in IDDM could be instrumental in limiting the plasma concentration of AGEs, the non-specific binding coming into play

after saturation of specific binding sites by higher plasma AGE-levels. Both binding strategies may act in concert as "damage limitation mechanisms" in the development of AGE-dependent diabetic complications.

L29 ANSWER 4 OF 15 EMBASE COPYRIGHT 1999 ELSEVIER  
 SCI. B.V.  
 AN 1998020373 EMBASE  
 TI American Diabetes Association Annual Meeting, 1997:  
 \*\*\*Endothelial\*\*\* dysfunction, neuropathy and the diabetic foot, diabetic mastopathy, and erectile dysfunction.  
 AU Bloomgarden Z.T.  
 SO Diabetes Care, (1998) 21/1 (183-189).  
 Refs: 5  
 ISSN: 0149-5992 CODEN: DICAD2  
 CY United States  
 DT Journal; Conference Article  
 FS 003 Endocrinology  
 006 Internal Medicine  
 037 Drug Literature Index  
 LA English

L29 ANSWER 5 OF 15 CAPLUS COPYRIGHT 1999 ACS  
 AN 1998:643914 CAPLUS  
 DN 130:50786  
 TI RAGE-A beta. interactions in the pathophysiology of Alzheimer's disease  
 AU Yan, Shi Du; Stern, David; Kane, Michael D.; Kuo, Yu-Min; Lampert, Heather C.; Rohrer, Alex E.  
 CS Department of Pathology, Surgery, Medicine and Physiology, College of Physicians and Surgeons, Columbia University, New York, NY, 10032, USA  
 SO Restor. Neurol. Neurosci. (1998), 12(2,3), 167-173  
 CODEN: RNNEEL; ISSN: 0922-6028  
 PB IOS Press  
 DT Journal  
 LA English  
 AB RAGE is a cell surface mol. primarily identified for its capacity to bind advanced glycation end-products and amphoterin. Immunocytochem. studies demonstrated that in Alzheimer's disease (AD) the expression of RAGE is elevated in neurons close to neuritic plaque beta-amyloid (A beta.) deposits and in the cells of A beta. cong. vessels. Crosslinking of surface bound A beta. 1-40 to \*\*\*endothelial\*\*\* cells, yielded a band of 50 kDa identified as RAGE. Using the sol. extracellular domain of recombinant human RAGE, we found that A beta. binds to RAGE with a K<sub>d</sub> = 57

nM, a value close to those found for mouse brain \*\*\*endothelial\*\*\* cells and rat cortical neurons. The interaction of A. beta. with RAGE in neuronal, \*\*\*endothelial\*\*\*, and RAGE-transfected COS-1 cells induced oxidative stress, as assessed by the TBARS and MTT assays. ELISA demonstrated a 2.5 times increase of RAGE in AD over control brains. Activated microglia also showed elevated expression of RAGE. In the BV-2 microglial cell line, RAGE bound A. beta. in a dose dependent manner with a Kd of 25 nM. Sol. A. beta. induced the migration of microglia along a concn. gradient, while immobilized A. beta. arrested this migration. A. beta.-RAGE interaction also activated NF-kappa.B, resulting in neuronal up-regulation of macrophage-colony stimulating factor (M-CSF) which also induced microglial migration. Apparently, RAGE-A. beta. interactions play an important role in the pathophysiol. of Alzheimer's disease.

L29 ANSWER 6 OF 15 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1997:340038 BIOSIS  
DN PREV199799639241  
TI Characterization and functional analysis of the promoter of RAGE, the receptor for advanced glycation end products.  
AU Li, Jianfeng; Schmidt, Ann Marie (1)  
CS (1) Columbia Univ. Coll. Phys. Surg., 630 W. 168 St., P S 11-518, New York, NY 10032 USA  
SO Journal of Biological Chemistry, (1997) Vol. 272, No. 26, pp. 16498-16506.  
ISSN: 0021-9258.  
DT Article  
LA English  
AB The receptor for advanced glycation end products, RAGE, is a member of the immunoglobulin superfamily of cell surface molecules expressed on a range of cell types. Ligation of RAGE perturbs homeostatic mechanisms and, potentially, provides a basis for cellular dysfunction in pathologic situations in which its ligands accumulate. To understand factors underlying RAGE expression, we cloned the 5'-flanking region of the RAGE gene and characterized putative regulatory motifs. Analysis of the putative promoter region revealed the presence of three potential NF-kappa-B-like and two SP1 binding sites. Transient transfection

of vascular \*\*\*endothelial\*\*\* and smooth muscle cells using chimeric 5'-deletion constructs linked to luciferase reporter revealed that the region -1543/-587 contributed importantly to both basal and stimulated expression of the RAGE gene. This region of the RAGE gene contained three putative NF-kappa-B-like binding sites and was responsible for increased luciferase activity observed when \*\*\*endothelial\*\*\* or smooth muscle cells were stimulated with lipopolysaccharide. DNase I footprinting assays and electrophoretic mobility shift assay revealed that two of the three NF-kappa-B-like binding sites (1 and 2) were likely functional and responsive to stimuli. Upon simultaneous mutation of NF-kappa-B-like sites 1 and 2, both basal promoter expression and response to stimulation with LPS, as measured by relative luciferase activity, were significantly diminished. These results point to NF-kappa-B-dependent mechanisms regulating cellular expression of RAGE and suggest a means of linking RAGE to the inflammatory response.

L29 ANSWER 7 OF 15 CAPLUS COPYRIGHT 1999 ACS  
AN 1997:686811 CAPLUS  
DN 127:344270  
TI Advanced glycation end product (AGE)-mediated induction of tissue factor in cultured \*\*\*endothelial\*\*\* cells is dependent on RAGE  
AU Bierhaus, Angelika; Illmer, Thomas; Kasper, Michael; Luther, Thomas; Quehenberger, Peter; Trischler, Hans; Wahl, Peter; Ziegler, Reinhard; Muller, Martin; Nawroth, Peter P.  
CS Department of Internal Medicine I, University of Heidelberg, Germany  
SO Circulation (1997), 96(7), 2262-2271  
CODEN: CIRCZ; ISSN: 0009-7322  
PB American Heart Association  
DT Journal  
LA English  
AB Binding of advanced glycation end products (AGEs) to the cellular surface receptor (RAGE) induces translocation of the transcription factor NF-kappa.B into the nucleus and NF-kappa.B-mediated gene expression. This study examines the role of RAGE in the AGE albumin-mediated induction of \*\*\*endothelial\*\*\* tissue factor, known to be partly controlled by NF-kappa.B. \*\*\*Endothelial\*\*\* cells (ECs) were incubated in the

presence of an 18-mer phosphorothioate oligodeoxynucleotide antisense to the 5'-coding sequence of the RAGE gene (antisense RAGE; 0.1 .mu.mol/L). Sense oligonucleotides (sense RAGE; 0.1 .mu.mol/L) of the same region served as controls. The cellular uptake of oligonucleotides was controlled by immunofluorescence microscopy. RAGE transcription was suppressed by antisense RAGE, as demonstrated by RT-PCR reactions. AGE albumin-mediated activation of cultured ECs was studied after 48 h preincubation of ECs with antisense or sense RAGE. Electrophoretic mobility shift assays and Western blot anal. demonstrated that the albumin-induced translocation of NF-kappa.B from the cytoplasm into the nucleus was suppressed in the presence of antisense RAGE but not by sense RAGE. In parallel, AGE albumin-mediated tissue factor transcription, activity, and antigen were significantly reduced in ECs exposed to antisense RAGE, whereas sense RAGE (and nonspecific oligonucleotides) did not influence tissue factor expression. In conclusion, activation of ECs and induction of tissue factor by AGE albumin in ECs is dependent on RAGE.

L29 ANSWER 8 OF 15 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1997:362973 BIOSIS  
DN PREV199799654906  
TI Advanced glycation end products and their receptors co-localise in rat organs susceptible to diabetic microvascular injury.  
AU Soulis, T.; Thallas, V.; Youssef, S.; Gilbert, R. E.; McWilliam, R. G.; Murray-Mcintosh, R. P.; Cooper, M. E. (1)  
CS (1) Dep. Med., Univ. Melbourne, Austin Australia  
SO Diabetologia, (1997) Vol. 40, No. 6, pp. 619-628.  
ISSN: 0012-186X.  
DT Article  
LA English  
AB Advanced glycation end products (AGEs) are believed to play an important role in the development of diabetic complications. AGEs are increased in experimental diabetes and treatment with the inhibitor of advanced glycation end products, aminoguanidine, has been shown to attenuate the level of these products in tissues undergoing complications. Recently, an AGE-binding protein has been isolated from bovine lung \*\*\*endothelial\*\*\* cells and termed the receptor for advanced glycated end products

(RAGE).

The present study sought to determine the distribution of AGE and RAGE in tissues susceptible to the long-term complications of diabetes including the kidney, eye, nerve, arteries as well as in a tissue resistant to such complications, the lung. Using polyclonal antisera both AGE and RAGE were found to co-localize in the renal glomerulus. AGE staining was clearly increased with age and was further increased by diabetes. Aminoguanidine treatment reduced AGE accumulation in the kidney.

Co-localization of AGE

and RAGE was demonstrated in the inner plexiform layer and the inner limiting membrane of the retina and in nerve bundles from mesenteric

arteries. In the aorta, both AGE and RAGE were found in the intima, media

and adventitia. Medial staining was increased in diabetes and was reduced

by aminoguanidine treatment. A similar pattern was observed for RAGE in

the aorta. In the lung, RAGE was found widely distributed throughout the

lung whereas the distribution of AGE staining was more limited, primarily

localizing to macrophages. The co-localization of AGEs and RAGE in sites

of diabetic microvascular injury suggests that this ligand-receptor interaction may represent an important mechanism in the genesis of diabetic complications.

L29 ANSWER 9 OF 15 BIOSIS COPYRIGHT 1999 BIOSIS

AN 1997:130388 BIOSIS

DN PREV199799422201

TI Advanced glycation end products (AGEs) co-localize with AGE receptors in

the retinal vasculature of diabetic and of AGE-infused rats.

AU Stitt, Alan W.; Li, Yong M.; Gardiner, Thomas A.; Buccala, Richard; Archer,

Desmond B.; Vlassara, Helen (1)

CS (1) Picower Inst. Med. Res., 350 Community Drive, Manhasset, NY 11030 USA

SO American Journal of Pathology, (1997) Vol. 150, No. 2, pp.

523-531.

ISSN: 0002-9440.

DT Article

LA English

AB Advanced glycation end products (AGEs), formed from the nonenzymatic

glycation of proteins and lipids with reducing sugars, have been implicated in many diabetic complications; however, their role in

diabetic retinopathy remains largely unknown. Recent studies suggest that

the

cellular actions of AGEs may be mediated by AGE-specific receptors (AGE-R). We have examined the immunolocalization of AGEs and

AGE-R components R1 and R2 in the retinal vasculature at 2, 4, and 8

months after STZ-induced diabetes as well as in nondiabetic rats infused with AGE

bovine serum albumin for 2 weeks. Using polyclonal or monoclonal anti-AGE

antibodies and polyclonal antibodies to recombinant AGE-R1 and AGE-R2,

immunoreactivity (IR) was examined in the complete retinal vascular tree

after isolation by trypsin digestion. After 2, 4, and 8 months of diabetes, there was a gradual increase in AGE IR in basement

membrane. At 8 months, pericytes, smooth muscle cells, and \*\*\*endothelial\*\*\*

cells of the retinal vessels showed dense intracellular AGE IR. AGE epitopes

stained most intensely within pericytes and smooth muscle cells but less

in basement membrane of AGE-infused rats compared with the diabetic group.

Retinas from normal or bovine-serum-albumin-infused rats were largely

negative for AGE IR. AGE-R1 and -R2 colocalized strongly with AGEs of

vascular \*\*\*endothelial\*\*\* cells, pericytes, and smooth muscle cells

of either normal, diabetic, or AGE-infused rat retinas, and this distribution did not vary with each condition. The data indicate that

AGEs accumulate as a function of diabetes duration first within the basement

membrane and then intracellularly, co-localizing with cellular AGE-Rs.

Significant AGE deposits appear within the pericytes after long-term

diabetes or acute challenge with AGE infusion conditions associated with

pericyte damage. Co-localization of AGEs and AGE-Rs in retinal cells

points to possible interactions of pathogenic significance.

L29 ANSWER 10 OF 15 MEDLINE DUPLICATE

I AN 97368045 MEDLINE

DN 97368045

TI Recombinant \*\*\*advanced\*\*\* \*\*\*glycation\*\*\*

\*\*\*product\*\*\* \*\*\*receptor\*\*\* pharmacokinetics in normal and diabetic

rats.

AU Renard C; Chappey O; Wautier M P; Nagashima M; Lundh E;

Morser J; Zhao L;

Schmidt A M; Schermmann J M; Wautier J L

CS Laboratoire de Recherche en Biologie Vasculaire et Cellulaire, Universite

Paris 7, Hôpital Lariboisière, France.

SO MOLECULAR PHARMACOLOGY, (1997 Jul) 52 (1) 54-62.

Journal code: NGR. ISSN: 0026-895X.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals; Cancer Journals

EM 199710

EW 19971003

AB Vascular dysfunction in patients with diabetes mellitus is related to advanced glycation end product (AGE) formation. We previously showed that

AGEs produce an increase in vascular permeability and generated an oxidant

stress after binding to the receptor (RAGE) present on endothelium. RAGE,

a 35-kDa protein that belongs to the immunoglobulin superfamily, has been

cloned from a rat lung cDNA library, and recombinant rat soluble RAGE

(rR-RAGE) has been produced in insect cells. The sequence of RAGE is

highly conserved between human and rat. We studied the biological effect

of rR-RAGE and pharmacokinetics of 125I-rR-RAGE after intraperitoneal administration in normal and streptozotocin-induced

diabetic rats. rR-RAGE prevented albumin or inulin transfer through a

bovine aortic \*\*\*endothelial\*\*\* cell monolayer, restored the hyperpermeability observed in diabetic rats or induced in normal

diabetic rat red blood cells, and corrected the reactive oxygen intermediate production after intravenous or intraperitoneal

administration. After intravenous injection of 125I-rR-RAGE, the distribution half-life was longer (p < or = 0.01) in diabetic (0.15

and 4.01 hr) than in normal (0.02 and 0.21 hr) rats, as was the case for the

elimination half-lives (diabetic, 57.17 hr; normal, 26.02 hr; p < or = 0.01). Distribution volume was higher in diabetic than in normal

rats (6.94 and 3.24 liter/kg, respectively; p = 0.049). Our study showed that

rR-RAGE was biologically active in vivo and slowly cleared, which suggests

it could be considered as a potential therapy.

L29 ANSWER 11 OF 15 BIOSIS COPYRIGHT 1999 BIOSIS

AN 1997:98644 BIOSIS

DN PREV19979937847

TI A novel mechanism for the pathogenesis of diabetic retinopathy

- involving glucose-modified proteins.  
AU Lambourne, B. J.; Molinatti, P. A.; Chibber, R.; Kohner, E. M.; Sonksen, P. H.  
CS Diabetic Retinopathy Unit, Div. Med., UMDS, St. Thomas' Hosp., London UK  
SO International Journal of Microcirculation Clinical and Experimental, (1996) Vol. 16, No. 4, pp. 214.  
Meeting Info.: 1st Joint National Vascular Meeting of the British Microcirculation Society, the British Society for Cardiovascular Research and the Royal Society of Medicine Forum on Angiology Exeter, England, UK  
April 17-19, 1996  
ISSN: 0167-6865.  
DT Conference; Abstract  
LA English
- L29 ANSWER 12 OF 15 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1996:375628 BIOSIS  
DN PREV199699097984  
TI A novel cellular receptor for advanced glycation end products.  
AU Schmidt, Ann Marie (1); Hori, Osamu; Cao, Rong; Yan, Shi Du; Brett, Jerold; Wautier, Jean-Luc; Ogawa, Satoshi; Kuwabara, Keisuke; Masumoto, Masayasu; Stern, David  
CS (1) Dep. Physiol., P and S 11-518, Columbia Univ., Coll. Phys. Surg., 630 W. 168th, New York, NY 10032 USA  
SO Diabetes, (1996) Vol. 45, No. SUPPL. 3, pp. S77-S80.  
ISSN: 0012-1797.  
DT Article  
LA English  
AB Exposure of proteins to reducing sugars results in nonenzymatic glycation with the ultimate formation of advanced glycation end products (AGEs). One means through which AGEs modulate cellular functions is through binding to specific cell surface acceptor molecules. The receptor for AGEs (RAGE) is such a receptor and is a newly identified member of the immunoglobulin superfamily expressed on \*\*\*endothelial\*\*\* cells (ECs), mononuclear phagocytes (MPs), and vascular smooth muscle cells (SMCs) in both *vivo* and *in vitro*. Binding of AGEs to RAGE results in induction of cellular oxidant stress, as exemplified by the generation of thiobarbituric acid-reactive substances, expression of heme oxygenase type 1, and activation of the transcription factor NF-kappa-B, with consequences for a range of cellular functions. AGEs on the surface of diabetic red cells enhance binding to \*\*\*endothelial\*\*\* RAGE and result in enhanced oxidant stress in the vessel wall. By using reagents to selectively block access to RAGE, the role of this receptor in AGE-mediated perturbation of cellular properties can be dissected in detail.  
L29 ANSWER 13 OF 15 CAPLUS COPYRIGHT 1999 ACS  
AN 1994:266466 CAPLUS  
DN 120:266466  
TI The \*\*\*endothelial\*\*\* cell binding site for advanced glycation products consists of a complex: an integral membrane protein and a lactoferrin-like polypeptide  
AU Schmidt, Ann Marie; Mora, Rozalia; Cao, Rong; Yan, Shi Du; Brett, Jerold; Ramakrishnan, Rajasekhar; Tsang, T. Christopher; Simionescu, Maya; Stern, David  
CS Coll. Physicians Surg., Columbia Univ., New York, NY, 10032, USA  
SO J. Biol. Chem. (1994), 269(13), 9882-8  
CODEN: JBCHA3; ISSN: 0021-9258  
DT Journal  
LA English  
AB Advanced glycation end products (AGEs), formed as the result of extended interaction of proteins with ketoses, modulate central properties of \*\*\*endothelial\*\*\* cells and mononuclear phagocytes by interacting with a cell surface binding site comprised of a novel integral membrane protein (receptor for AGE = RAGE) and a lactoferrin-like polypeptide (LF-L), the latter having sequence identity to lactoferrin (LF). To further understand this cellular binding site, the interaction of RAGE with LF-L and LF was characterized. By ligand blotting and a solid state competitive binding assay, 125I-LF-L and 125I-LF bound to RAGE immobilized on nitrocellulose membranes or polypropylene tubes in a time-dependent and reversible manner, demonstrating a high affinity component with Kd approx 100 pM. The interaction of 125I-LF-L and 125I-LF with RAGE was independent of iron in LF and was competed by addn. of an excess of unlabeled carboxyl-terminal portion of LF. Crosslinking studies with purified 125I-LF-L and RAGE, in the presence of disuccinimidyl
- substrate, showed a new, slowly migrating band, corresponding to a complex of RAGE and LF-L, and crosslinking on mouse aortic \*\*\*endothelial\*\*\* cells showed two new slowly migrating bands on immunoblotting visualized with both anti-RAGE IgG and anti-LF-L IgG. These data lead the authors to propose that the \*\*\*endothelial\*\*\* cell surface binding site for AGEs consists of LF-L bound noncovalently to RAGE anchored in the cell membrane.  
L29 ANSWER 14 OF 15 CAPLUS COPYRIGHT 1999 ACS  
AN 1994:531448 CAPLUS  
DN 121:131448  
TI Advanced glycation end products (AGEs) on the surface of diabetic erythrocytes bind to the vessel wall via a specific receptor inducing oxidant stress in the vasculature: a link between surface-associated AGEs and diabetic complications  
AU Wautier, J.-L.; Wautier, M.-P.; Schmidt, A.-M.; Anderson, G. M.; Hori, O.; Zoukourian, C.; Capron, L.; Chappey, O.; Yan, S.-D.; et al.  
CS Coll. Physicians and Surgeons, Columbia Univ., New York, NY, 10032, USA  
SO Proc. Natl. Acad. Sci. U. S. A. (1994), 91(16), 7742-6  
CODEN: PNASA6; ISSN: 0027-8424  
DT Journal  
LA English  
AB Vascular complications are an important cause of morbidity and mortality in patients with diabetes. The extent of vascular complications has been linked statistically to enhanced adherence of diabetic erythrocytes to \*\*\*endothelial\*\*\* cells (ECs) and to the accumulation of a class of glycated proteins termed advanced glycation end products (AGEs). The authors hypothesized that formation of AGEs on the surface of diabetic erythrocytes could mediate their interaction with ECs leading to binding and induction of vascular dysfunction. Enhanced binding of diabetic erythrocytes to ECs was blocked by preincubation of erythrocytes with anti-AGE IgG or preincubation of ECs with antibodies to the AGE (RAGE). Immunoblotting of cultured human ECs and immunostaining of normal/diabetic human tissue confirmed the presence of RAGE in the vessel

wall. Binding of diabetic erythrocytes to endothelium generated an oxidant stress, as measured by prodn. of thiobarbituric acid-reactive substances (TBARS) and activation of the transcription factor NF-kappa B, both of which were blocked by probucol or anti-RAGE IgG. Erythrocytes from diabetic rats infused into normal rats had an accelerated, early phase of clearance that was prevented, in part, by antibody to RAGE. Liver tissue from rats infused with diabetic erythrocytes showed elevated levels of TBARS, which was prevented by pretreatment with anti-RAGE IgG or probucol. Thus, erythrocyte surface AGEs can function as ligands that interact with RAGE on endothelium. The extensive contact of diabetic erythrocytes bearing surface-assocd. AGEs with vessel wall RAGE could be important in the development of vascular complications.

L29 ANSWER 15 OF 15 CAPLUS COPYRIGHT 1999 ACS  
AN 1995:278135 CAPLUS  
DN 122:233401  
TI AGE-receptors and in vivo biological effects of AGEs  
AU Viassara, Helen  
CS The Picower Institute for Medical Research, Manhasset/New York, 11030, USA  
SO Spec. Publ. - R. Soc. Chem. (1994), 151(Maillard Reactions in Chemistry, Food, and Health), 254-61  
CODEN: SROCDQ; ISSN: 0260-6291  
DT Journal; General Review  
LA English  
AB A review with 32 refs. on \*\*\*advanced\*\*\* \*\*\*glycation\*\*\* \*\*\*end\*\*\* \*\*\*product\*\*\* (AGE) \*\*\*receptors\*\*\* and the biol. effects of AGEs is presented. Surface receptors for AGEs are found on macrophages, T-lymphocytes, \*\*\*endothelial\*\*\* cells (EC), mesangial cells (MS), fibroblasts, and smooth muscle cells. Binding of AGEs to these receptors tends to a range of cellular responses including monocyte chemotaxis, activation, growth factor release, increased matrix prodn., increased EC permeability, and procoagulant activity. A no. of these responses can be inhibited by anti-AGE-receptor antibodies, supporting the role of AGE-ligand/receptor interactions in these events. Evidence for similar AGE-mediated biol. effects in vivo was obtained recently: short-term (4-8 wk) exogenous AGE administration to normal rats and rabbits led to multiple vascular defects including vascular

permeability, mononuclear activation, and vasodilatory impairment. Longer treatment with AGEs (3 mo) led to arterial basement membrane thickening, mesangial expansion, and glomerulosclerotic changes. These alterations were largely prevented by simultaneous treatment with aminoguanidine. These studies suggest that the interaction of de novo implanted, reactive AGEs with cellular AGE-receptors of otherwise healthy tissues can generate renal, and vascular pathol. similar to that seen in diabetes, in the absence of either the genetic or the metabolic abnormalities linked to diabetes. Progressive loss of kidney function correlates with increasing circulating AGE levels, presumably reflecting tissue AGE-degrdn. products which are not cleared by the failing kidneys. The pronounced (apprx. 8-fold) increase in serum AGEs obsd. in diabetic anephric patients, a group particularly susceptible to accelerated atherosclerosis, indicates that uncleared "reactive" AGEs may be available for enhanced interaction with cellular AGE-receptors, accelerating existing pathol.

=> s l9 and mononuclear/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
2 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
4 FILES SEARCHED...  
L30 6 L9 AND MONONUCLEAR/AB,BI  
=> dup rem l30  
PROCESSING COMPLETED FOR L30  
L31 6 DUP REM L30 (0 DUPLICATES REMOVED)  
=> d l- bib ab  
YOU HAVE REQUESTED DATA FROM 6 ANSWERS -  
CONTINUE? Y(N)y

L31 ANSWER 1 OF 6 CAPLUS COPYRIGHT 1999 ACS  
AN 1997:525836 CAPLUS  
DN 127:204001  
TI Binding of beta-amyloid protein by an \*\*\*advanced\*\*\* \*\*\*glycation\*\*\* \*\*\*end\*\*\* - \*\*\*product\*\*\* \*\*\*receptor\*\*\* and possible treatment of Alzheimer's disease  
IN Stern, David; Schmidt, Ann Marie; Yan, Shi Du  
PA Trustees of Columbia University, USA

SO PCT Int. Appl., 91 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN,CNT 1  
PATENT NO. KIND DATE APPLICATION NO.  
DATE

P1 WO 9726913 AI 19970731 WO 97-US857 19970121  
W: AU, CA, JP, MX  
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE  
AU 9718327 AI 19970820 AU 97-18327 19970121  
PRAI US 96-592070 19960126  
WO 97-US857 19970121  
AB The beta-amyloid protein binds to a cell-surface RAGE (receptor for advanced glycation end products) in neural cells and induces neurotoxic damage typical of Alzheimer's disease. This interaction may be a useful target for treatment of Alzheimer's disease. Binding assays for the identification and characterization of beta-amyloid-binding proteins used to identify the interaction of beta-amyloid with RAGE are described. Peptides capable of inhibiting the interaction are reported.

L31 ANSWER 2 OF 6 CAPLUS COPYRIGHT 1999 ACS  
AN 1996:552734 CAPLUS  
DN 125:230717  
TI The receptor for advanced glycation end products (RAGE) is a central mediator of the interaction of AGE- beta.2microglobulin with human \*\*\*mononuclear\*\*\* phagocytes via an oxidant-sensitive pathway: implications for the pathogenesis of dialysis-related amyloidosis  
AU Miyata, Toshio; Hori, Osamu; Zhang, JingHua; Yan, Shi Du; Ferran, Luis; Iida, Yoshiyasu; Schmidt, Ann Marie  
CS Dep. Int. Med., Nagoya Univ. Sch. Med., Nagoya, 461, Japan  
SO J. Clin. Invest. (1996), 98(5), 1088-1094  
CODEN: JCINAO; ISSN: 0021-9738  
DT Journal  
LA English  
AB An important component of amyloid fibrils in dialysis-related amyloidosis is a form of beta.2-microglobulin modified with advanced glycation end products (AGEs) of the Maillard reaction, known as AGE-beta.2M. The authors demonstrate here that the interaction of AGE-beta.2M with \*\*\*mononuclear\*\*\* phagocytes (MPs), cells important in the pathogenesis of the inflammatory arthropathy of dialysis-related amyloidosis, is mediated by the receptor for AGEs, or RAGE. 125I-AGE-beta.2M

bound to immobilized RAGE or to MPs in a specific, dose-dependent manner, a process inhibited in the presence of RAGE blockade. AGE-beta.2M-mediated monocyte chemotaxis was prevented by excess sRAGE or anti-RAGE IgG. Induction of tumor necrosis factor- $\alpha$ . (TNF) expression by MPs exposed to AGE-beta.2M resulted from engagement of RAGE, as appearances of TNF transcripts and TNF antigen release into culture supernatants were prevented by addn. of sRAGE; a process mediated, at least in part, by oxidant stress. AGE-beta.2M reduced cytochrome c and the elaboration of TNF by MPs was inhibited by N-acetylcysteine. Consistent with these data, immunohistochem. studies of AGE-laden amyloid deposits of a long-term hemodialysis patient reveals pos. staining for RAGE in the MPs infiltrating these lesions. These data indicate that RAGE is a central binding site for AGEs formed in vivo and suggest that AGE-beta.2M-MP-RAGE interaction likely contributes to the initiation of an inflammatory response in amyloid deposits of long-term hemodialysis patients, a process which may ultimately lead to bone and joint destruction.

L31 ANSWER 3 OF 6 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1996:375628 BIOSIS  
 DN PREV19969097984  
 AU Schmidt, Ann Marie (1); Horn, Osamu; Cao, Rong; Yan, Shi Du; Brett, Jerold; Wautier, Jean-Luc; Ogawa, Satoshi; Kuwabara, Keisuke; Masayasu, Stern, David  
 CS (1) Dep. Physiol., P and S 11-518, Columbia Univ., Coll. Phys. Surg., 630 W. 168th, New York, NY 10032 USA  
 SO Diabetes, (1996) Vol. 45, No. SUPPL. 3, pp. S77-S80. ISSN: 0012-1797.  
 DT Article  
 LA English  
 AB Exposure of proteins to reducing sugars results in nonenzymatic glycation with the ultimate formation of advanced glycation end products (AGEs). One means through which AGEs modulate cellular functions is through binding to specific cell surface acceptor molecules. The receptor for AGEs (RAGE) is such a receptor and is a newly identified member of the immunoglobulin superfamily expressed on endothelial cells (ECs),

\*\*\*mononuclear\*\*\* phagocytes (MPs), and vascular smooth muscle cells (SMCs) in both *in vivo* and *in vitro*. Binding of AGEs to RAGE results in induction of cellular oxidant stress, as exemplified by the generation of thiobarbituric acid-reactive substances, expression of heme oxygenase type I, and activation of the transcription factor NF-kappa-B, with consequences for a range of cellular functions. AGEs on the surface of diabetic red cells enhance binding to endothelial RAGE and result in enhanced oxidant stress in the vessel wall. By using reagents to selectively block access to RAGE, the role of this receptor in AGE-mediated perturbation of cellular properties can be dissected in detail.

L31 ANSWER 4 OF 6 CAPLUS COPYRIGHT 1999 ACS  
 AN 1994:266466 CAPLUS  
 DN 120:266466  
 TT The endothelial cell binding site for advanced glycation end products consists of a complex: an integral membrane protein and a lactoferrin-like polypeptide  
 AU Schmidt, Ann Marie; Mora, Rozalia; Cao, Rong; Yan, Shi Du; Brett, Jerold; Ramakrishnan, Rajasekhar; Tsang, T. Christopher; Simionescu, Maya; Stern, David  
 CS Coll. Physicians Surg., Columbia Univ., New York, NY, 10032, USA  
 SO J. Biol. Chem. (1994), 269(13), 9882-8  
 CODEN: JBCHA3; ISSN: 0021-9258  
 DT Journal  
 LA English  
 AB Advanced glycation end products (AGEs), formed as the result of the extended interaction of proteins with ketoses, modulate central properties of endothelial cells and \*\*\*mononuclear\*\*\* phagocytes by interacting with a cell surface binding site comprised of a novel integral membrane protein (receptor for AGE = RAGE) and a lactoferrin-like polypeptide (LF-L), the latter having sequence identity to lactoferrin (LF). To further understand this cellular binding site, the interaction of RAGE with LF-L and LF was characterized. By ligand blotting and a solid state competitive binding assay, 125I-LF-L and 125I-LF bound to RAGE immobilized on nitrocellulose membranes or polypropylene tubes in a

time-dependent and reversible manner, demonstrating a high affinity component with Kd approx. 100 pM. The interaction of 125I-LF-L and 125I-LF with RAGE was independent of iron in LF and was competed by addn. of an excess of unlabeled carboxyl-terminal portion of LF. Crosslinking studies with purified 125I-LF-L and RAGE, in the presence of disuccinimidyl suberate, showed a new, slowly migrating band, corresponding to a complex of RAGE and LF-L, and crosslinking on mouse aortic endothelial cells showed two new slowly migrating bands on immunoblotting visualized with both anti-RAGE IgG and anti-LF-L IgG. These data lead the authors to propose that the endothelial cell surface binding site for AGEs consists of LF-L bound noncovalently to RAGE anchored in the cell membrane.

L31 ANSWER 5 OF 6 CAPLUS COPYRIGHT 1999 ACS  
 AN 1995:278135 CAPLUS  
 DN 122:233401  
 TT AGE-receptors and *in vivo* biological effects of AGEs  
 AU Vlassara, Helen  
 CS The Picower Institute for Medical Research, Manhasset/New York, 11030, USA  
 SO Spec. Publ. - R. Soc. Chem. (1994), 151 (Maillard Reactions in Chemistry, Food, and Health), 234-61  
 CODEN: SROCDQ; ISSN: 0260-6291  
 DT Journal, General Review  
 LA English  
 AB A review with 32 refs. on \*\*\*advanced\*\*\* \*\*\*glycation\*\*\* and \*\*\*end\*\*\* \*\*\*product\*\*\* (AGE) \*\*\*receptors\*\*\* and the biol. effects of AGEs is presented. Surface receptors for AGEs are found on macrophages, T-lymphocytes, endothelial cells (EC), mesangial cells (MS), fibroblasts, and smooth muscle cells. Binding of AGEs to these receptors leads to a range of cellular responses including monocyte chemotaxis, activation, growth factor release, increased matrix prodn., increased EC permeability, and procoagulant activity. A no. of these responses can be inhibited by anti-AGE-receptor antibodies, supporting the role of AGE-ligand/receptor interactions in these events. Evidence for similar AGE-mediated biol. effects *in vivo* was obtained recently: short-term (4-8 wk) exogenous AGE administration to normal rats and rabbits led

to multiple vascular defects including vascular permeability, \*\*\*mononuclear\*\*\* activation, and vasodilatory impairment.

Longer treatment with AGEs (3 mo) led to arterial basement membrane thickening, mesangial expansion, and glomerulosclerotic changes. These alterations were largely prevented by simultaneous treatment with aminoguanidine.

These studies suggest that the interaction of de novo implanted, reactive AGEs with cellular AGE-receptors of otherwise healthy tissues can generate renal, and vascular pathol. similar to that seen in diabetes, in the absence of either the genetic or the metabolic abnormalities linked to diabetes. Progressive loss of kidney function correlates with increasing circulating AGE levels, presumably reflecting tissue AGE-degradn. products which are not cleared by the failing kidneys. The pronounced (apprx. 8-fold) increase in serum AGEs obsd. in diabetic anephric patients, a group particularly susceptible to accelerated atherosclerosis, indicates that uncaptured "reactive" AGEs may be available for enhanced interaction with cellular AGE-receptors, accelerating existing pathol.

L31 ANSWER 6 OF 6 CAPLUS COPYRIGHT 1999 ACS  
AN 1994:240942 CAPLUS  
DN 120:240942

T1 Survey of the distribution of a newly characterized receptor for glycation end products in tissues

AU Brett, Jerold; Schmidt, Ann Marie; Yan, Shi Du; Zou, Yu Shan; Weidman, Elliott; Pinsky, David; Nowygrod, Roman; Neepet, Michael; Przysiecki, Craig; et al.

CS Coll. Physicians Surg., Columbia Univ., New York, NY, 10032, USA

SO Am. J. Pathol. (1993), 143(6), 1699-712  
CODEN: AJPA44; ISSN: 0002-9440

DT Journal  
LA English

AB Advanced glycation end products (AGEs), the final products of non-enzymic glycation and oxidn. of proteins, are found in the plasma and accumulate in the tissues during aging and at an accelerated rate in diabetes. A novel integral membrane protein, termed receptor for AGE (RAGE), forms a central part of the cell surface binding site for AGEs. Using monospecific, polyclonal antibody raised to human recombinant and bovine

RAGE, immunostaining of bovine tissues showed RAGE in the vasculature, endothelium, and smooth muscle cells and in \*\*\*mononuclear\*\*\* cells in the tissues. Consistent with these data, RAGE antigen and mRNA were identified in cultured bovine endothelium, vascular smooth muscle, and monocyte-derived macrophages. RAGE antigen was also visualized in bovine cardiac myocytes as well as in cultures of neonatal rat cardiac myocytes and in neural tissue where motor neurons, peripheral nerves, and a population of cortical neurons were pos. In situ hybridization confirmed the presence of RAGE mRNA in the tissues, and studies with rat PC12 pheochromocytes indicated that they provide a neuronal-related cell culture model for examg. RAGE expression. Pathol. studies of human atherosclerotic plaques showed infiltration of RAGE-expressing cells in the expanded intima. These results indicate that RAGE is present in multiple tissues and suggest the potential relevance of AGE-RAGE interactions for modulating properties of the vasculature as well as neural and cardiac function, prominent areas of involvement in diabetes and in the normal aging process.

=> s 19 and tumor#/#ab,bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
2 FILES SEARCHED...

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
4 FILES SEARCHED...

L32 4 L9 AND TUMOR#/#AB,BI

=> dup rem l32

PROCESSING COMPLETED FOR L32  
L33 4 DUP REM L32 (0 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 4 ANSWERS -  
CONTINUE? Y(N)?y

L33 ANSWER 1 OF 4 MEDLINE  
AN 1999009111 MEDLINE  
DN 99009111

T1 A redox-triggered ras-effector interaction. Recruitment of phosphatidylinositol 3'-kinase to Ras by redox stress.

AU Deora A A; Win T; Vanhaesebroeck B; Lander H M

CS Department of Biochemistry, Cornell University Medical College, New York, New York 10021, USA.  
NC GM55509 (NIGMS)  
AI37637 (NIAID)  
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1998 Nov 6) 273 (45) 29923-8.  
Journal code: HIV. ISSN: 0021-9258.

CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199902  
EW 19990204  
AB Reactive free radical species are known to trigger biochemical events culminating in transcription factor activation and modulation of gene expression. The cytosolic signaling events triggered by free radicals that result in nuclear responses are largely unknown. Here we identify a signaling cascade triggered immediately upon redox activation of Ras. We examined two physiologically relevant models of redox signaling: 1) nitric oxide in human T cells, and 2) advanced glycation end product in pheochromocytoma cells. Reactive free radical species generated by nitric oxide donors and the interaction of \*\*\*advanced\*\*\* glycation\*\*\* with its \*\*\*receptor\*\*\* led to the recruitment of p85/p110 phosphatidylinositol 3'-kinase (PI3K) to the plasma membrane, where it associated directly with the effector domain of Ras and became activated. Only the p110beta and p110delta (but not p110alpha) catalytic subunits were recruited by redox-activated Ras.

Activation of downstream targets of PI3K such as protein kinase B/Akt and mitogen-activated protein kinase was found to be PI3K dependent. Our study demonstrates that nitrosative and oxidative stressors trigger Ras-dependent and PI3K-regulated events in cells and define a biochemical pathway that is triggered by redox signaling.

L33 ANSWER 2 OF 4 CAPLUS COPYRIGHT 1999 ACS  
AN 1997:525836 CAPLUS  
DN 127:204001

T1 Binding of beta -amyloid protein by an \*\*\*advanced\*\*\* glycation\*\*\* - \*\*\*end\*\*\* - \*\*\*product\*\*\* receptor\*\*\* and possible treatment of Alzheimer's disease

IN Stern, David, Schmidt, Ann Marie; Yan, Shi Du  
PA Trustees of Columbia University, USA  
SO PCT Int. Appl., 91 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN/CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.
WO 97-05857	A1	19970731	WO 97-05857

PI WO 9726913 A1 19970731 WO 97-05857 19970121

W: AU, CA, JP, MX

RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,

MC, NL, PT, SE

AU 9718327 A1 19970820 AU 97-18327 19970121

PRAI US 96-592070 19960126

WO 97-05857 19970121

AB The beta-amyloid protein binds to a cell-surface RAGE

(receptor for

advanced glycation end products) in neural cells and induces

neurotoxic

damage typical of Alzheimer's disease. This interaction may be a

useful

target for treatment of Alzheimer's disease. Binding assays for the

identification and characterization of beta-amyloid-binding

proteins

used to identify the interaction of beta-amyloid with RAGE are

described. Peptides capable of inhibiting the interaction are

reported.

L33 ANSWER 3 OF 4 CAPLUS COPYRIGHT 1999 ACS

AN 1996:552734 CAPLUS

DN 125/230717

TI The receptor for advanced glycation end products (RAGE) is a

central

mediator of the interaction of AGE-beta.2microglobulin with

human

mononuclear phagocytes via an oxidant-sensitive pathway:

implications for

the pathogenesis of dialysis-related amyloidosis

AU Miyata, Toshio; Hori, Osamu; Zhang, JingHua; Yan, Shi Du;

Ferran, Luis;

Ida, Yoshiyasu; Schmidt, Ann Marie

CS Dep. Int. Med., Nagoya Univ. Sch. Med., Nagoya, 461, Japan

SO J. Clin. Invest. (1996), 98(5), 1088-1094

CODEN: JCINAO; ISSN: 0021-9738

DT Journal

LA English

AB An important component of amyloid fibrils in dialysis-related

amyloidosis

is a form of beta.2-microglobulin modified with advanced

glycation end

products (AGEs) of the Maillard reaction, known as

AGE-beta.2M. The

authors demonstrate here that the interaction of AGE-beta.2M with

mononuclear phagocytes (MPs), cells important in the pathogenesis

of the

inflammatory arthropathy of dialysis-related amyloidosis, is  
mediated by

the receptor for AGEs, or RAGE. 125I-AGE-beta.2M bound to

immobilized

RAGE or to MPs in a specific, dose-dependent manner, a process

inhibited

in the presence of RAGE blockade. AGE-beta.2M-mediated

monocyte

chemotaxis was prevented by excess sRAGE or anti-RAGE IgG.

Induction of

\*\*\*tumor\*\*\* necrosis factor-alpha. (TNF) expression by MPs

exposed to

AGE-beta.2M resulted from engagement of RAGE, as appearances

of TNF

transcripts and TNF antigen release into culture supernatants were

prevented by addn. of sRAGE, a process mediated, at least in part,

by

oxidant stress. AGE-beta.2M reduced cytochrome c and the

elaboration of

TNF by MPs was inhibited by N-acetylcysteine. Consistent with

these data,

immunohistochem. studies of AGE-laden amyloid deposits of a

long-term

hemodialysis patient reveals pos. staining for RAGE in the MPs

infiltrating these lesions. These data indicate that RAGE is a

central

binding site for AGEs formed in vivo and suggest that

AGE-beta.2M-MP-RAGE

interaction likely contributes to the initiation of an inflammatory

response in amyloid deposits of long-term hemodialysis patients, a

process

which may ultimately lead to bone and joint destruction.

L33 ANSWER 4 OF 4 MEDLINE

AN 95096076 MEDLINE

DN 95096076

TI Receptor-mediated endocytic uptake of methylglyoxal-modified

serum

albumin. Competition with advanced glycation end

product-modified serum

albumin at the \*\*\*advanced\*\*\* \*\*\*glycation\*\*\*

\*\*\*end\*\*\*

\*\*\*product\*\*\* \*\*\*receptor\*\*\*

AU Westwood M E; McLellan A C; Thornalley P J

CS Department of Chemistry and Biological Chemistry, University

of Essex,

Colchester, United Kingdom.

SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1994 Dec 23)

269 (51) 32293-8.

Journal code: HIV. ISSN: 0021-9258.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals; Cancer Journals

EM 199503

AB Methylglyoxal binds and irreversibly modifies arginine and lysine

residues

in bovine serum albumin (BSA) under physiological conditions,  
producing a

protein with an increased net negative charge at physiological pH.

At 4

degrees C, methylglyoxal-modified BSA (MG-BSA) was bound by

cell surface

receptors on murine P388D1 macrophages. The apparent

dissociation constant

KD value was 435 +/- 2 nM, and there were 8.89 +/- 0.02 x 10(5)

receptors/cell (n = 6), compare with an apparent KD value of 263

+/- 52 nM

and 10.17 +/- 0.93 x 10(5) receptors/cell (n = 11) for advanced

glycation

end product-modified BSA (AGE-BSA). AGE-BSA competed with

MG-BSA for

binding to a common receptor; however, a component of

AGE-BSA receptor

binding could not be displaced by MG-BSA, and a component of

MG-BSA

receptor binding could not be displaced by AGE-BSA, suggesting

that there

are binding sites for both AGE-BSA and MG-BSA, competitive

and

noncompetitive, to MG-BSA and AGE-BSA on P388D1 cells at 4

degrees C. At

37 degrees C, receptor binding of AGE-BSA and MG-BSA was

followed by

endocytosis and lysosomal degradation of the modified protein.

Methylglyoxal-modified proteins are ligands for the AGE receptor,

and

their formation and metabolism may be linked to the development

of

diabetic complications.

=> s i9 and pc12/ab,bi

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

2 FILES SEARCHED...

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

L34 2 L9 AND PC12/AB,BI

=> dup rem l34

PROCESSING COMPLETED FOR L34

L35 2 DUP REM L34 (0 DUPLICATES REMOVED)

=> d l - bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS -

CONTINUE? Y(N)/y

L35 ANSWER 1 OF 2 CAPLUS COPYRIGHT 1999 ACS

AN 1997:544866 CAPLUS

DN 127:201264



TI Beta amyloid toxicity does not require RAGE protein  
AU Liu, Y.; Dargusch, R.; Schubert, D.  
CS The Salk Institute for Biological Studies, La Jolla, CA, 92037, USA  
SO Biochem. Biophys. Res. Commun. (1997), 237(1), 37-40  
CODEN: BBRC99; ISSN: 0006-291X  
PB Academic  
DT Journal  
LA English  
AB It has been suggested that a receptor for advanced glycation end products (RAGE) is the nerve cell receptor for amyloid .beta. protein (A.beta.).  
To det. if this is indeed the case, two neural cell lines as well as rat cortical neurons were examd. for the presence of the mRNA for RAGE by PCR  
and northern blot anal. Although lung was strongly pos., in no case was RAGE mRNA detected in the cultured neural cells. Glycated albumin is a major ligand for RAGE and the cell surface RAGE protein is trypsin sensitive. In agreement with the mRNA data, trypsin treatment did not alter A.beta. toxicity, nor did glycated albumin modify the A.beta. response. It follows that RAGE is not the neural receptor for A.beta..

L35 ANSWER 2 OF 2 CAPLUS COPYRIGHT 1999 ACS  
AN 1994:240942 CAPLUS  
DN 120:240942  
TI Survey of the distribution of a newly characterized receptor for advanced glycation end products in tissues  
AU Brett, Jerold; Schmidt, Ann Marie; Yan, Shi Du; Zou, Yu Shan; Weidman, Elliott; Pinsky, David; Nowygrod, Roman; Neeper, Michael; Przysiecki, Craig; et al  
CS Coll. Physicians Surg., Columbia Univ., New York, NY, 10032, USA  
SO Am. J. Pathol. (1993), 143(6), 1699-712  
CODEN: AJPA44; ISSN: 0002-9440  
DT Journal  
LA English  
AB Advanced glycation end products (AGEs), the final products of glycation and oxidn. of proteins, are found in the plasma and accumulate in the tissues during aging and at an accelerated rate in diabetes. A novel integral membrane protein, termed receptor for AGE (RAGE), forms a central part of the cell surface binding site for AGEs. Using monospecific, polyclonal antibody raised to human recombinant and bovine RAGE, immunostaining of bovine tissues showed RAGE in the vasculature,

endothelium, and smooth muscle cells and in mononuclear cells in the tissues. Consistent with these data, RAGE antigen and mRNA were identified in cultured bovine endothelium, vascular smooth muscle, and monocyte-derived macrophages. RAGE antigen was also visualized in bovine cardiac myocytes as well as in cultures of neonatal rat cardiac myocytes and in neural tissue where motor neurons, peripheral nerves, and a population of cortical neurons were pos. In situ hybridization confirmed the presence of RAGE mRNA in the tissues, and studies with rat \*\*\*PC12\*\*\* pheochromocytes indicated that they provide a neuronal-related cell culture model for examg. RAGE expression. Pathol. studies of human atherosclerotic plaques showed infiltration of RAGE-expressing cells in the expanded intima. These results indicate that RAGE is present in multiple tissues and suggest the potential relevance of AGE-RAGE interactions for modulating properties of the vasculature as well as neural and cardiac function, prominent areas of involvement in diabetes and in the normal aging process.

=> s 11 and pc12/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L36 88 L1 AND PC12/AB,BI  
=> s 136 and 19  
'AB' IS NOT A VALID FIELD CODE  
2 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L37 0 L36 AND L9  
=> s 11 and tumor#/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
3 FILES SEARCHED...  
'AB' IS NOT A VALID FIELD CODE  
L38 37 L1 AND TUMOR#/AB,BI  
=> dup rem 138  
PROCESSING COMPLETED FOR L38  
L39 31 DUP REM L38 (6 DUPLICATES REMOVED)

=> d 1 - bib ab  
YOU HAVE REQUESTED DATA FROM 31 ANSWERS - CONTINUE? Y(N/Y)

L39 ANSWER 1 OF 31 CAPLUS COPYRIGHT 1999 ACS  
AN 1999:96387 CAPLUS  
DN 130:164015  
TI Characterization of transcription factor Sel-10 and its use in drug screening  
IN Greenwald, Iva; Hubbard, E. Jane  
PA The Trustees of Columbia University in the City of New York, USA  
SO PCT Int. Appl., 86 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN/CNT 1  
PATENT NO. KIND DATE APPLICATION NO.  
DATE  
PI WO 9905307 AI 19990204 WO 98-US15335 19980723  
W: AU, CA, JP, MX, US  
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE  
PRAI US 97-899578 19970724  
AB This invention provides a cDNA sequence of the Sel-10 gene and the transcription factor thus encoded, which appears to be a member of the Cdc4 family of proteins. Genetic evidence indicates that Sel-10 is a neg. regulator of lin-12 mediated signaling in C. elegans, whereby lin-12 activity is controlled by controlling lin-12/Notch protein levels. Since Notch can induce mammalian \*\*\*tumors\*\*\* and since sel-10 downregulates Notch activity, it is suggested that sel-10 behaves as a \*\*\*tumor\*\*\* suppressor. This invention further provides methods for identifying compds. that are capable of treating cancer and Alzheimer's disease.

L39 ANSWER 2 OF 31 MEDLINE  
AN 1999101168 MEDLINE  
DN 99101168  
TI Dual roles of proteasome in the metabolism of \*\*\*presenilin\*\*\*  
1.  
AU Honda T; Yasutake K; Nihonmatsu N; Mercken M; Takahashi H; Murayama O; Murayama M; Sato K; Omori A; Tsubuki S; Saido T C; Takashima A  
CS Laboratory for Alzheimer's Disease, Brain Science Institute, RIKEN, Saitama, Japan.

SO JOURNAL OF NEUROCHEMISTRY, (1999 Jan) 72 (1) 255-61.  
Journal code: JAV. ISSN: 0022-3042.

CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199903  
EW 19990304

AB \*\*\*Presenilin\*\*\* 1 (PS1) has been identified as a causative gene for most early-onset familial Alzheimer's disease. Biochemical studies revealed that PS1 exists predominantly as two processed fragments in cells and brain tissues. We prepared stably transfected cells expressing the wild-type and familial Alzheimer's disease-associated mutants of PS1 and investigated the enzyme that participates in the metabolism of PS1. After treatment of the cells with proteasome inhibitors, the full-length PS1 was significantly accumulated. The levels of N- and C-terminal fragments were also increased. The accumulation of PS1 with a deletion of exon 10, which is unable to be processed, on treatment of the transfected cells with lactacystin indicated that proteasome can degrade full-length PS1.

A synthetic peptide that includes the processing region of PS1 was cleaved by 20S proteasome at the putative processing sites after Met288 and Glu299. Metabolic labeling experiments showed that the appearance of the N-terminal fragment was attenuated by the inhibitor. Finally, 28-kDa N- and 20-kDa C-terminal fragments were generated by purified PS1 in vitro. These data indicated that the proteasome pathway is involved in PS1 processing. These results demonstrate that the proteasome pathway plays dual roles in processing and degradation of PS1.

L39 ANSWER 3 OF 31 CAPLUS COPYRIGHT 1999 ACS  
AN 1998:682417 CAPLUS  
DN 129:286713  
TI Diagnosis of genetic disease arising from frameshift mutation by RT-PCR and hybridization or antibody assay, and treatment with hammerhead ribozyme cleavage of defective mRNA  
IN Van Leeuwen, Frederik W.; Grosveld, Franklin G.; Burbach, Johannes Peter  
Henri  
PA Royal Netherlands Academy of Arts and Sciences, Neth.;

Erasmus University  
Rotterdam; University of Utrecht  
SO PCT Int. Appl., 258 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN CNT 1  
PATENT NO. KIND DATE APPLICATION NO.  
DATE

P1 WO 9845322 A2 19981015 WO 98-IB705 19980402  
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM, RW, GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG  
AU 9870715 A1 19981030 AU 98-70715 19980402  
PRAI US 97-43163 19970410  
WO 98-IB705 19980402

AB The invention relates to methods and reagents for the diagnosis of a disease, specifically neurodegenerative diseases, caused by or associated with frameshift mutations, by RT-PCR and nucleic acid probe hybridization or by antibody detection of disease protein epitopes, and for treating these diseases via ribozyme cleavage of the mutant mRNA. The diagnostic methods include the steps of providing a body fluid or tissue sample from a patient, and analyzing the sample for the presence of an RNA mol. having a frameshift mutation or a protein encoded thereby, wherein the presence of the mutated RNA mol. or encoded protein is indicative of the disease. The therapeutic treatments include administering substances which selectively eliminate mutated RNA mol. from the cell, such as a genetic construct which provides for the synthesis of a ribozyme capable of selective cleavage of the mutant RNA. Neurodegenerative diseases such as Alzheimer's disease Down's syndrome are preferred examples to which these diagnostic and therapeutic method may be applied.

L39 ANSWER 4 OF 31 MEDLINE  
AN 1999069372 MEDLINE

DN 99069372  
TI Abrogation of the \*\*\*presenilin\*\*\* 1/beta-catenin interaction and preservation of the heterodimeric \*\*\*presenilin\*\*\* 1 complex following caspase activation.  
AU Tesco G; Kim T W; Diehlmann A; Beyreuther K; Tanzi R E  
CS Genetics and Aging Unit, Massachusetts General Hospital and Harvard Medical School, Charlestown, Massachusetts 02129, USA.  
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1998 Dec 18) 273 (51) 33909-14.  
Journal code: HIV. ISSN: 0021-9258.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199903  
EW 19990305

AB beta-Catenin has previously been shown to interact with \*\*\*presenilin\*\*\* 1 (PS1) in transfected cells. Here we report that beta-catenin co-immunoprecipitates with the endogenous C-terminal fragment of \*\*\*presenilin\*\*\* 1 (PS1-CTF) but not with the endogenous CTF of \*\*\*presenilin\*\*\* 2 (PS2-CTF) in H4 human neuroglioma cells. During staurosporine (STS)-induced cell death, beta-catenin and PS1-CTF undergo a caspase-mediated cleavage. After 12 h of STS treatment, the beta-catenin PS1-CTF interaction is abrogated. While PS1-CTF immunoprecipitated with all caspase-cleaved species of beta-catenin, beta-catenin holoprotein did not co-immunoprecipitate with the "alternative" caspase-derived PS1-CTF (PS1-aCTF). Thus, the abrogation of the beta-catenin PS1-CTF complex was due to caspase cleavage of PS1-CTF. beta-Catenin co-immunoprecipitated with PS1-NTF, but only when PS1-NTF was associated with PS1-CTF. Even though PS1-NTF.CTF complex stability was not altered by caspase cleavage, its ability to bind beta-catenin was abolished. Thus, while the PS1-NTF.CTF complex is preserved after caspase cleavage, it may no longer be fully functional.

L39 ANSWER 5 OF 31 MEDLINE  
AN 1999047359 MEDLINE  
DN 99047359  
TI Prominent expression of \*\*\*presenilin\*\*\* -1 in senile plaques and reactive astrocytes in Alzheimer's disease brain.  
AU Weggen S; Diehlmann A; Buslei R; Beyreuther K; Bayer T A  
CS Department of Psychiatry, University of Bonn Medical Center, Germany.  
SO NEUROREPORT, (1998 Oct 5) 9 (14) 3279-83.

Journal code: A6M. ISSN: 0959-4965.

CY ENGLAND; United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199903  
EW 19990304  
AB Mutations in the \*\*\*presenilin\*\*\* 1 (PS-1) gene account for most cases of autosomal dominant early-onset familial Alzheimer's disease (AD). In order to elucidate the cellular expression profile of PS-1 we used a novel N-terminal monoclonal antibody against human PS-1. Immunohistochemical staining was observed strongly in senile plaques, and reactive astrocytes of gray and white matter. Neuronal immunoreactivity, however, was found to be only moderate. RT-PCR analysis of PS-1 mRNA revealed expression throughout human development as well as in human glioma cell lines.  
Altered PS-1 function may contribute to plaque formation in AD.

L39 ANSWER 6 OF 31 MEDLINE  
AN 1998421807 MEDLINE  
DN 98421807  
TI \*\*\*Presenilin\*\*\* 1 mutations linked to familial Alzheimer's disease  
increase the intracellular levels of amyloid beta-protein 1-42 and its N-terminally truncated variant(s) which are generated at distinct sites.

AU Sudoh S; Kawamura Y; Sato S; Wang R; Saïdo T C; Oyama F; Sakaki Y; Komano H; Yanagisawa K  
CS Department of Dementia Research, National Institute for Longevity Sciences, Obu, Aichi, Japan.  
SO JOURNAL OF NEUROCHEMISTRY, (1998 Oct) 71 (4) 1535-43.  
Journal code: JAV. ISSN: 0022-3042.

CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199812  
AB Mutations in the \*\*\*presenilin\*\*\* genes PS1 and PS2 cause the most common form of early-onset familial Alzheimer's disease. The influence of PS1 mutations on the generation of endogenous intracellular amyloid beta-protein (A beta) species was assessed using a highly sensitive immunoblotting technique with inducible mouse neuroblastoma (Neuro 2a) cell lines expressing the human wild-type (wt) or mutated PS1

(M146L or delta exon 10). The induction of mutated PS1 increased the intracellular levels of two distinct A beta species ending at residue 42 that were likely to be A beta1-42 and its N-terminally truncated variant(s) A beta x-42. The induction of mutated PS1 resulted in a higher level of intracellular A beta1-42 than of intracellular A beta x-42, whereas extracellular levels of A beta1-42 and A beta x-42 were increased proportionally. In addition, the intracellular generation of these A beta42 species in wt and mutated PS1-induced cells was completely blocked by brefeldin A, whereas it exhibited differential sensitivities to monensin: the increased accumulation of intracellular A beta x-42 versus inhibition of intracellular A beta1-42 generation. These data strongly suggest that A beta x-42 is generated in a proximal Golgi, whereas beta1-42 is generated in a distal Golgi and/or a post-Golgi compartment. Thus, it appears that PS1 mutations enhance the degree of 42-specific gamma-secretase cleavage that occurs in the normal beta-amyloid protein processing pathway (a) in the endoplasmic reticulum or the early Golgi apparatus prior to beta-secretase cleavage or (b) in the distinct sites where A beta x-42 and A beta1-42 are generated.

L39 ANSWER 7 OF 31 MEDLINE  
AN 1998294913 MEDLINE  
DN 98294913  
TI Caspase-mediated cleavage is not required for the activity of \*\*\*presenilins\*\*\* in amyloidogenesis and NOTCH signaling.  
AU Brockhaus M; Grunberg J; Rohrig S; Loetscher H; Wittenburg N; Baumeister R; Jacobsen H; Haass C  
CS F. Hoffmann-La Roche Ltd, Pharma Division, Preclinical CNS Research-Gene Technology, Basel, Switzerland.  
SO NEUROREPORT, (1998 May 11) 9 (7) 1481-6.  
Journal code: A6M. ISSN: 0959-4965.

CY ENGLAND; United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199810  
EW 19981005  
AB The Alzheimer's disease (AD) associated \*\*\*presenilin\*\*\* (PS) proteins are proteolytically processed. One of the processing pathways involves cleavage by caspases. Pharmacological inhibition of caspases is currently being discussed as a treatment for a variety of neurodegenerative

diseases, including AD. We therefore inhibited caspase mediated processing of PS-1 and PS-2 in cells transfected with wt and mutant PS by mutating the substrate recognition site or by using specific peptide aldehydes known to block caspases. We found that the inhibition of caspase mediated processing of PS proteins does not decrease its amyloidogenic activity. PS cDNA constructs with mutations in the caspase cleavage site are biologically active in *Caenorhabditis elegans* such as the wt human PS proteins, demonstrating that caspase-mediated cleavage is not required for the physiological PS function in NOTCH signaling.

L39 ANSWER 8 OF 31 EMBASE COPYRIGHT 1999 ELSEVIER  
SCI. B.V.DUPLICATE 1  
AN 1998240134 EMBASE  
TI Inhibition of \*\*\*presenilin\*\*\* 1 expression is promoted by p53 and p21(WAF-1) and results in apoptosis and \*\*\*tumor\*\*\* suppression.  
AU Roperch J.-P.; Alvaro V.; Prieur S.; Tuynder M.; Nemani M.; Lethrosne F.; Plouffe L.; Gendron M.-C.; Israeli D.; Dausset J.; Oren M.; Telerman A.  
CS A. Telerman, Fondation Jean Dausset-CEPH, 27 rue Juliette Dodu, 75010 Paris, France. Telerman@cephb.fr  
SO Nature Medicine, (1998) 4/7 (835-838).  
Refs: 20

ISSN: 1078-8956 CODEN: NAMEFI  
CY United States  
DT Journal; Article  
FS 005 General Pathology and Pathological Anatomy  
022 Human Genetics  
LA English  
SL English  
AB Previously, we cloned a cDNA fragment, TSIP 2 ( \*\*\*tumor\*\*\* suppressor inhibited pathway clone 2), that detects by northern blot analysis of M1-LTR6 cells a 3-kb mRNA downregulated during p53-induced apoptosis. Cloning the full-length TSIP 2 cDNA showed that it corresponds to the \*\*\*presenilin\*\*\* 1 (PS1) gene, in which mutations have been reported in early-onset familial Alzheimer's disease. Here we demonstrate that PS1 is downregulated in a series of model systems for p53-dependent and p53-independent apoptosis and \*\*\*tumor\*\*\* suppression. To investigate the biological relevance of this downregulation, we stably transfected

U937 cells with antisense PS1 cDNA. The downregulation of PS1 in these U937 transfectants results in reduced growth with an increased fraction of the cells in apoptosis. When injected into mice homozygous for severe combined immunodeficiency disease (scid/scid mice), these cells show a suppression of their malignant phenotype. Our results indicate that PS1, initially identified in a neurodegenerative disease, may also be involved in the regulation of cancer-related pathways.

L39 ANSWER 9 OF 31 MEDLINE  
AN 1998330911 MEDLINE  
DN 98330911  
TI Stable association of \*\*\*presenilin\*\*\* derivatives and absence of \*\*\*presenilin\*\*\* interactions with APP.

AU Thinakaran G; Regard J B; Bouton C M; Harris C L; Price D L; Borchelt D R; Sisodia S S  
CS Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205-2196.  
NC 1P01 AG 14248 (NIA)  
5 P50 AG 05146 (NIA)  
SO NEUROBIOLOGY OF DISEASE, (1998 Apr) 4 (6) 438-53.  
Journal code: CJUN. ISSN: 0969-9961.

CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199811  
EW 19981104  
AB Mutations in two related genes, \*\*\*presenilin\*\*\* 1 and 2 \*\*\*presenilin\*\*\* 2 (PS1 and PS2), cosegregate with Alzheimer's disease.  
PS1 and PS2 are highly homologous polytopic membrane proteins that are subject to endoproteolytic cleavage in vivo. The resulting N- and C-terminal derivatives are the predominant PS-related species that accumulate in cultured cells and tissue. In earlier studies, we demonstrated that PS1 N- and C-terminal derivatives accumulate to 1:1 stoichiometry and that the absolute levels of fragments are established by a tightly regulated and saturable mechanism. These findings led to the suggestion that the levels of PS1 derivatives might be determined by their association with limiting cellular components. In this study, we use situ chemical cross-linking and coimmunoprecipitation analyses to document that the N- and C-terminal derivatives of either PS1 or PS2 can be

coisolated. Moreover, and in contrast to published reports which documented that PS1 and PS2 form stable heteromeric assemblies with the beta-amyloid precursor protein (APP), we have failed to provide evidence for physiological complexes between PS1 and PS2 holoproteins or their derivatives with APP.

L39 ANSWER 10 OF 31 MEDLINE  
AN 1998363099 MEDLINE  
DN 98363099  
TI Effect of steroid receptors, pS2 and cathepsin D on the outcome of elderly breast cancer patients: an exploratory investigation.

AU Coradini D; Biganzoli E; Boracchi P; Bombardieri E; Seregni E; De Palo G; Martelli G; Di Fronzo G  
CS Division of Experimental Oncology C, Istituto Nazionale per lo Studio e la Cura dei Tumori, Milan, Italy. coradini@istitutotumori.mi.it  
SO INTERNATIONAL JOURNAL OF CANCER, (1998 Aug 21) 79 (4) 305-11.  
Journal code: GQU. ISSN: 0020-7136.

CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199810  
EW 19981004  
AB In 83 elderly breast cancer patients, oestrogen and progesterone receptors (ER, PgR), pS2 and cathepsin D (CathD) were evaluated for their possible prognostic role on disease-free survival (DFS). The biomarkers were determined on the same cytosol by using immunoradiometric assays, and the variables were considered on a continuous scale. Univariate analysis indicated a linear relationship between logarithmic hazard ratio (log(HR)) and the log(ER) and log(PgR) concentration, but a non-linear relationship between log(HR) and CathD. As regards pS2, there was no evidence of a relationship with log(HR). In multivariate analysis, log(ER) content did not have a significant prognostic role, whereas log(PgR) retained a significant prognostic role. As regards the predictive ability, log(PgR) was the best discriminator of outcome followed by CathD, whereas the contribution of log(ER) was negligible. In multivariate analysis, 2 models were considered: one with log(ER), pS2, CathD and the interaction between

pS2 and CathD, and another with log(PgR), pS2, CathD and the interaction between pS2 and CathD. In the first model, log(ER) content did not have a significant prognostic role, whereas in the second model log(PgR) retained a significant prognostic role. Our findings indicate that the quantitative determination of pS2 and CathD, in addition to steroid receptors, on the same cytosolic fraction could be a complementary tool to describe all breast cancer patients rather than just the elderly and that the use of a continuous scale, instead of a simple dichotomous "status", may improve the biological information supplied by the variables.

L39 ANSWER 11 OF 31 CAPLUS COPYRIGHT 1999 ACS  
AN 1998707646 CAPLUS  
DN 130:151570  
TI Relationship between immunoinflammatory reactions and Alzheimer's disease

AU Du, Zeyang; Li, Xiaoyu  
CS Shanghai Institute of Medicine, Chinese Academy of Sciences, Shanghai, 200031, Peop. Rep. China  
SO Shengli Xueue Jinzhan (1998), 29(3), 253-256  
CODEN: SLKHA8; ISSN: 0559-7765

PB Zhongguo Shengli Xuehui  
DT Journal; General Review  
LA Chinese  
AB A review with 10 refs. was reported on the relationship between immunoinflammatory reactions and Alzheimer's disease (AD) with the subsections as follows: (1) the pathol. characteristics of AD; (2) the proofs of immunoinflammatory reactions; (3) the natural inhibitors in vivo; (4) conclusions. The medicine inhibiting or blocking the immunoinflammatory reactions in CNS may play an important role in the prevention of AD.

L39 ANSWER 12 OF 31 MEDLINE  
AN 1999090638 MEDLINE  
DN 99090638  
TI Measurement of pS2 protein in pancreatic cyst fluids. Evidence for a potential role of pS2 protein in the pathogenesis of mucinous cystic \*\*\*tumors\*\*\*

AU Yang J M; Lee J; Southern J F; Warshaw A L; Dhanak E; Lewandrowski K B  
CS Department of Pathology, Massachusetts General Hospital, Harvard Medical School, Boston, USA.  
SO INTERNATIONAL JOURNAL OF PANCREATOLOGY, (1998 Dec) 24 (3) 181-6.

Journal code: IJP. ISSN: 0169-4197.

CY United States  
DT Journal: Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199905  
EW 19990504  
AB CONCLUSION: Elevated levels of the growth factor pS2 protein in the cyst  
fluids of mucinous cystic \*\*\*tumors\*\*\* correlate with earlier observations using immunohistochemical techniques showing that pS2 protein is expressed by these \*\*\*tumors\*\*\*. The markedly elevated levels of pS2 protein compared to normal plasma values suggest that this growth factor may be important in the pathogenesis of pancreatic mucinous cystic \*\*\*tumors\*\*\*. BACKGROUND: Cystic lesions of the pancreas include inflammatory pseudocysts, serous cystadenomas, and mucinous cystic \*\*\*tumors\*\*\*, some of which are malignant. Previous studies using immunohistochemical techniques have shown that virtually all pancreatic mucinous \*\*\*tumors\*\*\* express pS2 protein. pS2 protein is a growth factor that is believed to be important in the normal process of inflammation and repair. We measured pS2 protein and other growth factors in pancreatic cyst fluids to assess their potential pathophysiologic and diagnostic significance. METHODS: Levels of pS2 protein were measured in 54 pancreatic cyst fluids by radioimmunoassay. The growth factors, epidermal growth factor (EGF), transforming growth factor-alpha (TGF-alpha), and insulin-like growth factors I and II (IGF-I, IGF-II) were measured in 22 cyst fluids using commercial immunoassays. RESULTS: Mucinous cysts exhibited significantly higher levels of cyst fluid pS2 protein than nonmucinous lesions, including pseudocysts and serous cystadenomas (median: 78,303 pg/mL; range: 218-361,176 pg/mL vs median: 886 pg/mL; range: 0-14,206 pg/mL; p = 0.0001). The level of pS2 in mucinous \*\*\*tumors\*\*\* was markedly higher than plasma values (median: 392 pg/mL). Levels of pS2 protein in malignant mucinous lesions tended to be higher than those in benign mucinous cysts, but this difference was not statistically significant (median: 88,817 vs 64,350 pg/mL; p = 0.159).

Levels of other growth factors, including EGF, TGF-alpha, IGF-I, and IGF-II, did not discriminate among the different cyst types, and the values were within normal plasma ranges.

L39 ANSWER 13 OF 31 MEDLINE  
AN 1998249534 MEDLINE  
DN 98249534  
TI The pS2 protein in colorectal carcinomas and metastases.  
AU Hackel C; Falkenberg B; Gunther T; Lippert H; Roessner A  
CS Institute of Pathology, Otto-von-Guericke University, Magdeburg, Germany.  
carsten.hackel@medizin.uni-magdeburg.de  
SO PATHOLOGY, RESEARCH AND PRACTICE, (1998) 194 (3) 171-6.

Journal code: PBZ. ISSN: 0344-0338.  
CY GERMANY: Germany, Federal Republic of  
DT Journal: Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199809  
EW 19980901  
AB Expression of pS2 protein in 50 primary \*\*\*tumors\*\*\*, metastases and recurrent \*\*\*tumors\*\*\* of colorectal carcinomas has been analyzed by immunohistochemistry. Sixty percent of the primary \*\*\*tumors\*\*\* were at least focally positive for the antigen. There was no correlation between pS2 expression and histologic grade of the lesions. In contrast, pS2 expression in T4 and T3 \*\*\*tumors\*\*\* was significantly higher than in T2 carcinomas. Immunoreactions in carcinomas with distant metastases (M1) were stronger than in M0 cases. However, this difference did not reach statistical significance. The presence of lymph node metastases did not correlate with pS2 expression. High expression of pS2 in T4 carcinomas together with the finding of pronounced expression of the antigen at invasion fronts in single cases could be interpreted as a function in \*\*\*tumor\*\*\* cell invasion and motility. However, in metastases and recurrent \*\*\*tumors\*\*\*, pS2 expression did not differ from primary lesions (53% positive lesions). All in all, under consideration of the latter finding in particular and together with the randomly distributed immunopositive \*\*\*tumor\*\*\* cells and clusters in the majority of cases, it is more likely that the expression pattern of pS2 in colorectal carcinomas is a result of overall \*\*\*tumor\*\*\* cell heterogeneity.

L39 ANSWER 14 OF 31 MEDLINE  
AN 1999038014 MEDLINE

DN 99038014  
TI Tamoxifen aziridine binding to cytosolic proteins from human breast specimens is negatively associated with estrogen receptors, progesterone receptors, pS2, and cathepsin-D.  
AU Navarro D; Doreste H; Cabrera J J; Morales M; Diaz-Chico J C; Diaz-Chico B  
N  
CS Dept. Endocrinologia Celular y Molecular, Centro de Ciencias de la Salud, Universidad de Las Palmas de Gran Canaria, Las Palmas, Spain.  
SO BREAST CANCER RESEARCH AND TREATMENT, (1998 Jul) 50 (2) 155-66.  
Journal code: A8X. ISSN: 0167-6806.

CY Netherlands  
DT Journal: Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199903  
EW 19990304  
AB [3H]Tamoxifen Aziridine ([3H]TAZ) is a derivative of the antiestrogen tamoxifen that covalently labels the Estrogen Receptor (ER), and perhaps other uncharacterized proteins. In a previous article we described that [3H]TAZ binds to a cytosolic protein from human uterine tissues that shares some, but not all, the ER properties. Here we have extended these studies to [3H]TAZ binding to cytosol proteins from human breast cancer specimens, and studied its quantitative association with other molecular markers and clinico-pathological variables. Cytosols were obtained in hypotonic buffer containing 20 mM molybdate and protease inhibitors, incubated with [3H]TAZ, and subjected to Sucrose Gradient Analysis (SGA). A [3H]TAZ labeled peak that consistently migrated with the 4S fractions was found in most of the assayed cytosols (range of 0 to 1278 fmol/mg p.). The 4S peak of [3H]TAZ was partially inhibited by both estrogens and antiestrogens. When [3H]E2 was used instead of [3H]TAZ, only an 8S peak was detected. [3H]TAZ was covalently bound to a protein with an apparent MW of 65 kDa, as determined by SDS-PAGE and fluorography. The mean of [3H]TAZ binding was significantly higher in the subgroups of samples classified as ER-, PR-, pS2- or cathepsin D-, than in the respective positive subgroups (P < 0.01 in all the cases). [3H]TAZ binding

was not associated with clinico-pathological variables, except that its mean was significantly larger in \*\*\*tumors\*\*\* larger than 5 cm than in smaller \*\*\*tumors\*\*\*. These results, and those previously reported, suggest that: 1) [3H]TAZ labels a cytosolic protein present in human breast cancers and uterine tissues that does not share all the ER properties, and 2) the [3H]TAZ binding by breast cancer cytosols is negatively associated with markers of estrogen dependency, and its quantification may provide valuable information on antiestrogen responsiveness of a given \*\*\*tumor\*\*\*.

L39 ANSWER 15 OF 31 MEDLINE  
 AN 1998409316 MEDLINE  
 DN 98409316  
 TI Direct association of \*\*\*presenilin\*\*\* -1 with beta-catenin.  
 AU Murayama M; Tanaka S; Palacino J; Murayama O; Honda T; Sun X; Yasutake K;  
 Nihonmatsu N; Wolozin B; Takashima A  
 CS Laboratory for Alzheimer's Disease, Brain Science Institute, RIKEN, Saitama, Japan.  
 SO FEBS LETTERS, (1998 Aug 14) 433 (1-2) 73-7.  
 Journal code: EUH. ISSN: 0014-5793.  
 CY Netherlands  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199812  
 EW 19981201  
 AB Families bearing mutations in the \*\*\*presenilin\*\*\* -1 (PS1) gene develop Alzheimer's disease (AD). However, the mechanism through which PS1 causes AD is unclear. The co-immunoprecipitation with PS1 in transfected COS-7 cells indicates that PS1 directly interacts with endogenous beta-catenin, and the interaction requires residues 322450 of PS1 and 445-676 of beta-catenin. Both proteins are co-localized in the endoplasmic reticulum. Over-expression of PS1 reduces the level of cytoplasmic beta-catenin, and inhibits beta-catenin-T cell factor-regulated transcription. These results indicate that PS1 plays a role as inhibitor of the beta-catenin signal, which may be connected with the AD dysfunction.

L39 ANSWER 16 OF 31 MEDLINE  
 AN 1998330346 MEDLINE  
 DN 98330346  
 TI Lack of specific association of \*\*\*presenilin\*\*\* -1 (PS-1)

protein with plaques and tangles in Alzheimer's disease.  
 AU Xia M Q; Berezovska O; Kim T W; Xia W M; Liao A; Tanzi R E; Selkoe D;  
 Hyman B T  
 CS Alzheimer's Research Unit, Department of Neurology, Massachusetts General Hospital-East, Charlestown 02129, USA.  
 NC AG05134 (NIA)  
 AG08487 (NIA)  
 AG14744 (NIA)  
 SO JOURNAL OF THE NEUROLOGICAL SCIENCES, (1998 Jun 1) 158 (1) 15-23.  
 Journal code: JBI. ISSN: 0022-510X.  
 CY Netherlands  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199812  
 EW 19981203  
 AB Missense mutations in the \*\*\*presenilin\*\*\* -1 (PS-1) gene are causally related to the majority of familial early-onset Alzheimer's disease (FAD).  
 PS-1 immunohistochemical expression in normal human brain and in brains with Alzheimer's disease (AD) has so far been controversial. Here, we report a study of PS-1 expression in brains, cell lines and peripheral blood mononuclear cells using a panel of well characterized PS-1-specific antibodies. These antibodies were characterized by immunofluorescent staining of PS-1 transfectants followed by flow cytometric analysis. In human brain, widespread neuronal staining was observed. PS-1 immunoreactivity was primarily confined to neuronal cell bodies and proximal dendrites. Weaker staining of microglia was also detected, in accord with the finding of PS-1 immunoreactivity in monocytes.  
 PS-1 expression is not particularly associated with neurons either containing or spared from neurofibrillary tangles, nor with senile plaques. The level of PS-1 expression does not differ between normal and AD brains. Immunoprecipitation from AD, FAD and control brains revealed only a 32 kDa N-terminal fragment and an 18-20 kDa C-terminal fragment. Little or no full length PS-1 was detected. The enriched presence of PS-1 in neurons implies an important role in neuronal function, however, the lack of apparent association of its expression with AD pathology signifies the need for a better understanding of its pathophysiological role.

L39 ANSWER 17 OF 31 CAPLUS COPYRIGHT 1999 ACS  
 AN 19981559 CAPLUS  
 DN 12873898  
 TI Transgenic animals expressing perlecan and amyloid genes at high levels and methods of identifying compounds for the treatment of amyloidoses  
 IN Snow, Alan; Fukuchi, Ken-ichiro; Hassell, John  
 PA University of Washington, USA  
 SO PCT Int. Appl., 146 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN/CNT 1  
 PATENT NO. KIND DATE APPLICATION NO.  
 DATE

PI WO 9746664 AI 19971211 WO 97-US9875  
 19970606  
 W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE,  
 ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR,  
 LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU,  
 SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, AM, AZ,  
 BY, KG, KZ, MD, RU, TJ, TM  
 RW: GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB,  
 GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN,  
 ML, MR, NE, SN, TD, TG  
 AU 9736402 AI 19980105 AU 97-36402 19970606  
 PRAI US 96-17830 19960606  
 WO 97-US9875 19970606  
 AB Transgenic animals expressing a foreign gene for a perlecan, or genes for perlecan and an amyloid are constructed for use in the testing of compds. that can alter the rate or extent of amyloid deposition.  
 Over-expression of perlecan and amyloid proteins results in animals showing symptoms closer to amyloidoses than found in animals only over-expressing an amyloid gene, esp. Alzheimer's disease. Over-expression of a gene encoding domains 1-V of mouse perlecan and the 695-amino acid .beta.-amyloid in P19 cells led to an up-regulation of .beta.-amyloid synthesis and secretion. P19 cells induced to form neurons degenerated when the perlecan gene was overexpressed.

L39 ANSWER 18 OF 31 MEDLINE  
 DUPLICATE  
 2

AN 1998019211	MEDLINE	DN 98019198	TI Generation of anti-apoptotic ***presenilin*** -2 polypeptides by alternative transcription, proteolysis, and caspase-3 cleavage.	TI Endoproteolytic processing and stabilization of wild-type and mutant ***presenilin***	
DN 98019211		AU Vito P; Ghayur T; D'Adamo L	CS T Cell Molecular Biology Unit, Laboratory of Cellular and Molecular Immunology, NIAID, National Institutes of Health, Bethesda, Maryland 20892, USA.	AU Ratovitski T; Slunt H H; Thinakaran G; Price D L; Sisodia S S; Borchelt D R	
TI Evidence that levels of ***presenilins*** (PS1 and PS2) are coordinately regulated by competition for limiting cellular factors.		SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1997 Nov 7) 272 (45) 28315-20	Journal code: HIV. ISSN: 0021-9258.	CS Division of Neuropathology, Johns Hopkins School of Medicine, Baltimore, Maryland 21205, USA.	
AU Thinakaran G; Harris C L; Ratovitski T; Davenport F; Slunt H H; Price D L;		CY United States	DT Journal; Article; (JOURNAL ARTICLE)	NC NS10580 (NINDS)	
Borchelt D R; Sisodia S S		LA English	FS Priority Journals; Cancer Journals	AG07914 (NIA)	
CS Department of Pathology, The Johns Hopkins University School of Medicine,		OS GENBANK-U57325	EM 199802	AG05146 (NIA)	
Baltimore, Maryland 21205-2196, USA.		EW 19980204	AB PS2, the chromosome 1 familial Alzheimer's disease gene, has been shown to be involved in programmed cell death by three complementary experimental approaches. Reduction of PS2 protein levels by antisense RNA protects from apoptosis, whereas overexpression of an Alzheimer's PS2 mutant cell death induced by several stimuli. In addition, ALG-3, a truncated PS2 cDNA, encodes an artificial COOH-terminal PS2 segment that dominantly inhibits apoptosis. Here we describe a physiological COOH-terminal PS2 polypeptide (PS2s, Met298-Ile448) generated by both an alternative PS2 transcript and proteolytic cleavage. We find that PS2s protects transfected cells from Fas- and ***tumor*** necrosis factor alpha (TNFalpha)-induced apoptosis. Furthermore, a similar anti-apoptotic COOH-terminal PS2 polypeptide (PS2Ccas) is generated by caspase-3 cleavage at Asp329. These results suggest that caspase-3 not only activates pro-apoptotic substrates but also generates a negative feedback signal in which PS2Ccas antagonizes the progression of cell death. Thus, whereas PS2 is required for apoptosis, PS2s and PS2Ccas oppose this process, and the balance between PS2 and these COOH-terminal fragments may dictate the cell fate.	SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1997 Sep 26) 272 (39) 24536-41.	
gopal@welchlink.welch.jhu.edu		DN 97450985	MEDLINE	Journal code: HIV. ISSN: 0021-9258.	
NC IPO1 AG14248 (NIA)		DN 97450985	MEDLINE	CY United States	
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1997 Nov 7) 272 (45) 28415-22		L39 ANSWER 20 OF 31	MEDLINE	DT Journal; Article; (JOURNAL ARTICLE)	
Journal code: HIV. ISSN: 0021-9258.		AN 97450985	MEDLINE	LA English	
CY United States		DN 97450985	MEDLINE	FS Priority Journals; Cancer Journals	
DT Journal; Article; (JOURNAL ARTICLE)				EM 199802	
LA English				EW 19980204	
FS Priority Journals; Cancer Journals				AB Mutations in two related genes, PS1 and PS2, account for the majority of early onset cases of familial Alzheimer's disease. PS1 and PS2 are homologous polytopic membrane proteins that are processed endoproteolytically into two fragments in vivo. In the present report we examine the fate of endogenous PS1 and PS2 after overexpression of human PS1 or PS2 in mouse N2a neuroblastoma cell lines and human PS1 transgenic mice. Remarkably, in N2a cell lines and in brains of transgenic mice expressing human PS1, accumulation of human PS1 derivatives is accompanied by a compensatory, and highly selective, decrease in the steady-state levels of murine PS1 and PS2 derivatives. Similarly, the levels of murine PS1 derivatives are diminished in cultured cells overexpressing human PS2. To define the minimal sequence requirements for "replacement", we expressed familial Alzheimer's disease-linked experimental deletion variants of PS1. These studies revealed that compromised accumulation of murine PS1 and PS2 derivatives resulting from overexpression of human PS1 occurs in a manner independent of endoproteolytic cleavage. Our results are consistent with a model in which the abundance of PS1 and PS2 fragments is regulated coordinately by competition for limiting cellular factor(s).	SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1997 Sep 26) 272 (39) 24536-41.
		L39 ANSWER 19 OF 31	MEDLINE	Journal code: HIV. ISSN: 0021-9258.	
		AN 1998019198	MEDLINE	CY United States	
		DUPLICATE		DT Journal; Article; (JOURNAL ARTICLE)	

- L39 ANSWER 21 OF 31 MEDLINE  
AN 97442406 MEDLINE  
DN 97442406  
TI Transcriptional regulation of the mouse **\*\*\*presenilin\*\*\*** -1 gene.
- AU Mitsuda N; Roses A D; Vitek M P  
CS Division of Neurology, Duke University Medical Center, Durham, North Carolina 27710, USA.  
NC RO1 AG-13839 (NIA)  
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1997 Sep 19) 272 (38) 23489-97.  
Journal code: HIV. ISSN: 0021-9258.
- CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
OS GENBANK-AF007560  
EM 199712  
AB The **\*\*\*presenilin\*\*\*** -1 (PS-1) gene encodes at least three separate mRNA transcripts from its 12 exons, which are spread over 50 kilobase pairs of mouse DNA. The first transcript begins with exon 1A, whereas the other transcripts begin with exon 1B. Different portions of exon 1B are spliced to give long and short mRNAs. The expression of all of these transcripts depends on a single promoter located just upstream of exon 1A. Although this region lacks a TATA box and a number of common initiator sequences, it does contain a CAAT box, a heat-shock responsive element, a polyomavirus enhancer activator-3 site, an Ets 1-3 site, and multiple-Spl and multiple-Ap2 binding sites, which are typically found in eukaryotic promoters. We have combined a reporter gene with various portions of this putative PS-1 promoter and measured firefly luciferase activity relative to an internal renilla luciferase standard. We identified a 25-base pair fragment spanning the 5'-transcription start site of exon 1A as containing the core of the promoter activity. The sequences downstream of this region had undetectable promoter activity, suggesting that this core element is the gene's only promoter, and it controls expression of all three transcripts. Although human PS-1 mRNA expression is clearly different from the mouse PS-1 mRNA pattern, the human and mouse core promoters do share limited homology.
- L39 ANSWER 22 OF 31 MEDLINE  
AN 97268991 MEDLINE  
DN 97268991  
TI Endoproteolytic cleavage and proteasomal degradation of **\*\*\*presenilin\*\*\*** 2 in transfected cells.
- AU Kim T W; Pettingell W H; Hallmark O G; Moir R D; Wasco W; Tanzi R E  
CS Genetics and Aging Unit, Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Charlestown, Massachusetts 02129, USA.  
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1997 Apr 25) 272 (17) 11006-10.  
Journal code: HIV. ISSN: 0021-9258.
- CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199707  
AB Mutations in the **\*\*\*presenilin\*\*\*** genes, PS1 and PS2, cause a major portion of early onset familial Alzheimer's disease (FAD). The biological roles of the **\*\*\*presenilins\*\*\*** and how their pathological mutations confer FAD are unknown. In this study, we set out to examine the processing and degradation pathways of PS2. For regulated expression of PS2, we have established inducible cell lines expressing PS2 under the tight control of the tetracycline-responsive transactivator. Western blot analysis revealed that PS2 was detected as an approximately 53-55-kDa polypeptide (54-kDa PS2) as well as a high molecular mass form (HMW-PS2). Using a stably transfected, inducible cell system, we have found that PS2 is proteolytically cleaved into two stable cellular polypeptides including an approximately 20-kDa C-terminal fragment and an approximately 34-kDa N-terminal fragment. PS2 is polyubiquitinated in vivo, and the degradation of PS2 is inhibited by proteasome inhibitors, N-acetyl-L-leucinal-L-norleucinal and lactacystin. Our studies suggest that PS2 normally undergoes endoproteolytic cleavage and is degraded via the proteasome pathway.
- L39 ANSWER 23 OF 31 MEDLINE  
AN 97289724 MEDLINE  
DN 97289724  
TI Evidence for phosphorylation and oligomeric assembly of **\*\*\*presenilin\*\*\***
1.  
AU Seeger M; Nordstedt C; Petanceska S; Kovacs D M; Gouras G K; Hahne S; Fraser P; Levesque L; Czernik A J; George-Hyslop P S; Sisodia S S; Thinakaran G; Tanzi R E; Greengard P; Gandy S  
CS Laboratory of Alzheimer Research, Department of Neurology and Neuroscience, Cornell University Medical College, New York, NY 10021, USA.  
NC AG09464 (NIA)  
AG11508 (NIA)  
AG13780 (NIA)  
+  
SO PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1997 May 13) 94 (10) 5090-4.  
Journal code: PV3. ISSN: 0027-8424.
- CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199708  
EW 19970801  
AB Pathogenic mutations in **\*\*\*presenilin\*\*\*** 1 (PS1) are associated with approximately 50% of early-onset familial Alzheimer disease. PS1 is endoproteolytically cleaved to yield a 30-kDa N-terminal fragment (NTF) and an 18-kDa C-terminal fragment (CTF). Using COS7 cells transfected with human PS1, we have found that phorbol 12, 13-dibutyrate and forskolin increase the state of phosphorylation of serine residues of the human CTF. Phosphorylation of the human CTF resulted in a shift in electrophoretic mobility from a single major species of 18 kDa to a doublet of 20-23 kDa. This mobility shift was also observed with human PS1 that had been transfected into mouse neuroblastoma (N2a) cells. Treatment of the phosphorylated CTF doublet with phage lambda protein phosphatase eliminated the 20- to 23-kDa doublet while enhancing the 18-kDa species, consistent with the interpretation that the electrophoretic mobility shift was due to the addition of phosphate to the 18-kDa species. The NTF and CTF eluted from a gel filtration column at an estimated mass of over 100 kDa, suggesting that these fragments exist as an oligomerized species. Upon phosphorylation of the PS1 CTF, the apparent mass of the NTF- or CTF-containing oligomers was unchanged. Thus, the association of



- PS1 fragments may be maintained during cycles of phosphorylation/dephosphorylation of the PS1 CTF.
- L39 ANSWER 24 OF 31 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1997-533425 BIOSIS  
DN PREV199799832628  
TI The role of \*\*\*presenilin\*\*\* -1 in the response of PC12 cells to nerve growth factor.  
AU Lah, J. J.; Bennett-Desmelik, J. A.; Heilman, C. J.; Nash, N. R.; Greenamyre, J. T.; Levey, A. I.  
CS Emory Univ., Dep. Neurol., Atlanta, GA USA  
SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 2167.  
Meeting Info.: 27th Annual Meeting of the Society for Neuroscience New Orleans, Louisiana, USA October 25-30, 1997  
ISSN: 0190-5295.  
DT Conference; Abstract; Conference  
LA English
- L39 ANSWER 25 OF 31 MEDLINE  
AN 97437409 MEDLINE  
DN 97437409  
TI Neuronal expression and intracellular localization of \*\*\*presenilins\*\*\* in normal and Alzheimer disease brains.  
AU Huynh D P; Vinters H V; Ho D H; Ho V V; Puls S M  
CS Neurogenetics Laboratory, Burns and Allen Research Institute, Cedars-Sinai Medical Center, Los Angeles, CA 90048, USA.  
NC P30 AG 10123 (NIA)  
SO JOURNAL OF NEUROPATHOLOGY AND EXPERIMENTAL NEUROLOGY, (1997 Sep) 56 (9) 1009-17.  
Journal code: JBR. ISSN: 0022-3069.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199712  
AB The expression patterns of \*\*\*presenilin\*\*\* 1 (PS1) and \*\*\*presenilin\*\*\* 2 (PS2) in human normal and Alzheimer disease (AD) brains were investigated using antibodies to specific N-terminal peptides of PS1 (Alzh14A and Alzh14B) and PS2 (Alzh1A-AB). The antibodies to peptides Alzh14A (Alzh14A-AB) and Alzh14B (Alzh14B-AB) detected the full-length protein (approximately 63 kDa) and the N-terminal-processed fragment (36 kDa) of PS1, while the Alzh1A-AB detected mainly the N-terminal-processed fragment (36 kDa) of PS2.
- Immunofluorescent staining detected by confocal microscopy suggested that both native PS1 and PS2 are localized mainly in the Golgi/ER apparatus. Immunohistochemical studies of human temporal lobes from 2 normal and 5 sporadic Alzheimer brains revealed high levels of PS1 and PS2 expression in the granule cell layer and pyramidal neurons of the hippocampus. Strong immunoreactivity was found in reactive astrocytes and neurofibrillary tangles of all 5 Alzheimer brains. In contrast, only 2 sporadic Alzheimer brains showed \*\*\*presenilin\*\*\* -positive neuritic plaques. These observations suggest that \*\*\*presenilins\*\*\* may be involved in the pathology of some cases of sporadic AD.
- L39 ANSWER 26 OF 31 MEDLINE  
AN 1998063306 MEDLINE  
DN 98063306  
TI Determination of a cleavage site of \*\*\*presenilin\*\*\* 2 protein in stably transfected SH-SY5Y human neuroblastoma cell lines.  
AU Shirotani K; Takahashi K; Ozawa K; Kunishita T; Tabira T  
CS Division of Demyelinating Disease and Aging, National Institute of Neuroscience, Tokyo, Japan.  
SO BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1997 Nov 26) 240 (3) 728-31.  
Journal code: 9Y8. ISSN: 0006-291X.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199803  
EW 19980303  
AB Mutations in the \*\*\*presenilin\*\*\* 1 (PS1) and \*\*\*presenilin\*\*\* 2 (PS2) genes are associated with early-onset autosomal dominant Alzheimer's disease, and the gene products are endoproteolytically processed to yield N-terminal fragments (NTF) and C-terminal fragments (CTF). We have studied the cleavage site of the PS2 protein in stably transfected human neuroblastoma cells. The 23 kD PS2-CTF was isolated by a combination of anion exchange chromatography and affinity chromatography and directly sequenced. The N-terminus of the PS2-CTF started at residue 307, which indicated that the cleavage occurs between Lys306 and Leu307 in
- the proximal portion of the large hydrophilic loop. This site is close to the cleavage positions observed in the PS1 protein.
- L39 ANSWER 27 OF 31 MEDLINE  
AN 97474235 MEDLINE  
DN 97474235  
TI Alzheimer's disease-associated \*\*\*presenilin\*\*\* 1 in neuronal cells: evidence for localization to the endoplasmic reticulum-Golgi intermediate compartment  
AU Culvenor J G; Maher F; Evin G; Maltchiodi-Albedi F; Cappai R; Underwood J R; Davis J B; Karran E H; Roberts G W; Beyreuther K; Masters C L  
CS Department of Pathology, The University of Melbourne, Parkville, Victoria, Australia.. j.culvenor@pathology.unimelb.edu.au  
SO JOURNAL OF NEUROSCIENCE RESEARCH, (1997 Sep 15) 49 (6) 719-31.  
Journal code: KAC. ISSN: 0360-4012.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199801  
AB The recently identified Alzheimer's disease-associated \*\*\*presenilin\*\*\* 1 and 2 (PS1 and PS2) genes encode two homologous multi membrane-spanning proteins. Rabbit antibodies to the N-terminal domain of PS1 detected PS1 in human neuroblastoma SH-SY5Y wild type and PS1 transfectants (SY5Y-PS1) as well as in mouse P19, in CHO-K1 and CHO-APP770 transfect cells, in rat cerebellar granule and hippocampal neurons, and astrocytes. Immunoblotting detected full-length protein of 50 kDa, and a major presumptive cleavage product of 30 kDa. The immunofluorescence pattern resembled labeling of the endoplasmic reticulum-Golgi intermediate compartment (ERGIC) marker protein ERGIC-53. PS1 distribution showed slight condensation after brefeldin A and more marked incubation of cells at 16 degrees C, characteristic of the ERGIC compartment. Double labeling showed colocalization of ERGIC-53 with PS1 in the SY5Y-PS1 cells. PS1 labeling of SY5Y-PS1 and P19 cells showed overlap of the cis-Golgi marker p210 and colocalization with p210 after brefeldin A which causes redistribution of p210 to the ERGIC. Expression of PS1 did not change in level or cellular distribution during development of

- neurons in culture. Double labeling for the amyloid precursor protein (APP) and PSI on SY5Y-PS1 cells and CHO-APP770 cells showed some overlap under control conditions. These results indicate that PS1 is a resident protein of the ERGIC and could be involved in trafficking of proteins, including APP, between the ER and Golgi compartments.
- L39 ANSWER 28 OF 31 MEDLINE  
AN 97364828 MEDLINE  
DN 97364828  
TI Alternative cleavage of Alzheimer-associated \*\*\*presenilins\*\*\* during apoptosis by a caspase-3 family protease.  
AU Kim T W, Pettingell W H, Jung Y K, Kovacs D M, Tanzi R E  
CS Genetics and Aging Unit, Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA 02129, USA.  
SO SCIENCE, (1997 Jul 18) 277 (5324) 373-6.  
Journal code: UJ7. ISSN: 0036-8075.
- CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199805  
EW 19980504  
AB Most cases of early-onset familial Alzheimer's disease (FAD) are caused by mutations in the genes encoding the \*\*\*presenilin\*\*\* 1 (PS1) and PS2 proteins, both of which undergo regulated endoproteolytic processing. During apoptosis, PS1 and PS2 were shown to be cleaved at sites distal to their normal cleavage sites by a caspase-3 family protease. In cells expressing PS2 containing the asparagine-141 FAD mutant, the ratio of alternative to normal PS2 cleavage fragments was increased relative to wild-type PS2-expressing cells, suggesting a potential role for apoptosis-associated cleavage of \*\*\*presenilins\*\*\* in the pathogenesis of Alzheimer's disease.
- L39 ANSWER 29 OF 31 MEDLINE  
AN 97092711 MEDLINE  
DN 97092711  
TI Familial Alzheimer's disease-linked \*\*\*presenilin\*\*\* 1 variants elevate Abeta1-42/1-40 ratio in vitro and in vivo.  
AU Borcheit D R, Thinakaran G, Eckman C B, Lee M K, Davenport F, Ratovitsky T, Prada C M, Kim G, Seekins S, Yager D, Slunt H H, Wang R, Seeger M.
- Levey A I; Gandy S E; Copeland N G; Jenkins N A; Price D L; Younkin S G; Sisodia S S  
CS Department of Pathology, The Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA.  
NC AG05146 (NIA)  
NS 20471 (NINDS)  
AG05689 (NIA)  
+  
SO NEURON, (1996 Nov) 17 (5) 1005-13.  
Journal code: AN8. ISSN: 0896-6273.
- CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199703  
AB Mutations in the \*\*\*presenilin\*\*\* 1 (PS1) and \*\*\*presenilin\*\*\* 2 genes cosegregate with the majority of early-onset familial Alzheimer's disease (FAD) pedigrees. We now document that the Abeta1-42(43)/Abeta1-40 ratio in the conditioned media of independent N2a cell lines expressing three FAD-linked PS1 variants is uniformly elevated relative to cells expressing similar levels of wild-type PS1. Similarly, the Abeta1-42(43)/Abeta1-40 ratio is elevated in the brains of young transgenic animals coexpressing a chimeric amyloid precursor protein (APP) and an FAD-linked PS1 variant compared with brains of transgenic mice expressing APP alone or transgenic mice coexpressing wild-type human PS1 and APP. These studies provide compelling support for the view that one mechanism by which these mutant PS1 cause AD is by increasing the extracellular concentration of Abeta peptides terminating at 42(43), species that foster Abeta deposition.
- L39 ANSWER 30 OF 31 MEDLINE  
AN 96216717 MEDLINE  
DN 96216717  
TI Regional and cellular \*\*\*presenilin\*\*\* 1 gene expression in human and rat tissues.  
AU Suzuki T, Nishiyama K, Murayama S, Yamamoto A, Sato S, Kanazawa I, Sakaki Y  
CS Human Genome Center, Institute of Medical Science, University of Tokyo, Japan.  
SO BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1996 Feb 27) 219 (3) 708-13.
- Journal code: 9Y8. ISSN: 0006-291X.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199609  
AB \*\*\*Presenilin\*\*\* 1 (PSNLI) is a novel causative gene for early-onset familial Alzheimer's disease (EOFAD). We have examined the regional and cellular distribution of PSLN1 gene expression in normal human and rat tissues. In situ hybridization and Northern blot analysis showed that PSNLI mRNA was ubiquitously expressed in many different organs. We also demonstrated that PSNLI mRNA was expressed predominantly in the neuronal cells of the central nervous system, but only at low-level in glial cells. Furthermore, the distribution of PSNLI mRNA in human and rodent brains was similar.
- L39 ANSWER 31 OF 31 MEDLINE  
AN 97179560 MEDLINE  
DN 97179560  
TI \*\*\*Presenilin\*\*\* -1 is processed into two major cleavage products in neuronal cell lines.  
AU Ward R V; Davis J B; Gray C W; Barton A J; Bresciani L G; Caivano M; Murphy V F; Duff K; Hutton M; Hardy J; Roberts G W; Karran E  
H  
CS Department of Molecular Neuropathology, Smithkline Beecham Pharmaceuticals, Harlow, Essex, UK.  
SO NEURODEGENERATION, (1996 Dec) 5 (4) 293-8.  
Journal code: B99. ISSN: 1055-8330.
- CY ENGLAND: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199706  
EW 19970604  
AB \*\*\*Presenilin\*\*\* -1 (PS-1) has been identified as the protein encoded by the chromosome 14 locus that, when mutated, leads to familial Alzheimer's disease (FAD). Using PS-1 transfected SHSY5Y neuroblastoma cells, we have demonstrated by immunodetection, using polyclonal antibodies, that PS-1 is processed to give two fragments: an N-terminal 28 kDa fragment, and a C-terminal 18 kDa fragment. In a number of non-transfected cell types, most PS-1 is detected as the cleaved products. The molecular weights of the PS-1 cleavage products suggest that the cleavage point will most probably be within a region of the

hydrophilic  
loop domain coded for by either exon 8 or 9 of the PS-1 gene. The clustering of FAD mutations within exon 8 strongly suggests that it encodes a key functional domain. It seems likely that the cleavage of PS-1 is crucial to some aspect of its functionality. An understanding of this process will give insights into the pathology of AD, and may offer new opportunities for therapeutic intervention.

=> s11 and mononuclear/ab,bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L40 8 L1 AND MONONUCLEAR/AB,BI

=> dup rem l40

PROCESSING COMPLETED FOR L40  
L41 2 DUP REM L40 (6 DUPLICATES REMOVED)

L41 ANSWER 1 OF 2 MEDLINE DUPLICATE 1  
AB Sporadic inclusion-body myositis (s-IBM) is the most common progressive muscle disease of older persons. The muscle biopsy demonstrates \*\*\*mononuclear\*\*\* cell inflammation and vacuolated muscle fibers containing paired helical filaments and 6- to 10-nm fibrils, both resembling those of Alzheimer disease brain and Congo red term hereditary inclusion-body myopathies (h-IBMs) designates autosomal-recessive or autosomal-dominant disorders with muscle biopsies cytopathologically similar to s-IBM but without inflammation. Vacuolated muscle fibers of both s-IBM and the h-IBMs contain accumulations of several "Alzheimer-characteristic proteins" including beta-amyloid protein and beta-amyloid precursor protein, and their paired helical filaments are composed of phosphorylated tau. We used six well characterized antibodies against several residues of \*\*\*presenilin\*\*\* 1 (PS1) to immunostain muscle biopsies of 12 patients with s-IBM, 5 patients with autosomal-recessive inclusion-body myopathy, and 16 normal and disease controls. Seventy to eighty percent of the vacuolated muscle fibers of both s-IBM and autosomal-recessive inclusion-body myopathy had inclusions

that were strongly PS1-immunoreactive, which by immunoelectron microscopy localized mainly to paired helical filaments and 6- to 10-nm filaments. None of the control biopsies had PS1-positive inclusions characteristic of the s- and h-IBM abnormal muscle fibers. Mutations of the newly discovered PS1 gene are responsible for early-onset familial Alzheimer disease (AD), and PS1 is abnormally accumulated in sporadic and familial AD brain. Our study provides the first demonstration of PS1 abnormality in non-neural tissue and in diseases other than AD and suggests that the cytopathogenesis in AD brain and IBM muscle may share similarities.

L41 ANSWER 2 OF 2 MEDLINE DUPLICATE 2  
AB Missense mutations in the \*\*\*presenilin\*\*\* -1 (PS-1) gene are causally related to the majority of familial early-onset Alzheimer's disease (FAD). PS-1 immunohistochemical expression in normal human brain and in brains with Alzheimer's disease (AD) has so far been controversial. Here, we report a study of PS-1 expression in brains, cell lines and peripheral blood \*\*\*mononuclear\*\*\* cells using a panel of well characterized PS-1-specific antibodies. These antibodies were characterized by immunofluorescent staining of PS-1 transfectants followed by flow cytometric analysis. In human brain, widespread neuronal staining was observed. PS-1 immunoreactivity was primarily confined to neuronal cell bodies and proximal dendrites. Weaker staining of microglia was also detected, in accord with the finding of PS-1 immunoreactivity in monocytes. PS-1 expression is not particularly associated with neurons either containing or spared from neurofibrillary tangles, nor with senile plaques. The level of PS-1 expression does not differ between normal and AD brains. Immunoprecipitation from AD, FAD and control brains revealed only a 32 kDa N-terminal fragment and an 18-20 kDa C-terminal fragment. Little or no full length PS-1 was detected. The enriched presence of PS-1 in neurons implies an important role in neuronal function, however, the lack of apparent association of its expression with AD pathology signifies the need for a better understanding of its pathophysiological role.

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 2 ANSWERS - CONTINUE? Y(N)y

L41 ANSWER 1 OF 2 MEDLINE DUPLICATE 1  
AN 199806425 MEDLINE  
DN 98206425  
TI Light and electron microscopic immunolocalization of \*\*\*presenilin\*\*\* 1 in abnormal muscle fibers of patients with sporadic inclusion-body myositis and autosomal-recessive inclusion-body myopathy. AU Askamas V; Engel W K; Yang C C; Alvarez R B; Lee V M; Wisniewski T CS USC Neuromuscular Center, Los Angeles, California 90017-1912, USA. SO AMERICAN JOURNAL OF PATHOLOGY, (1998 Apr) 152 (4) 889-95. Journal code: JRS. ISSN: 0002-9440. CY United States DT Journal: Article; (JOURNAL ARTICLE) LA English FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals EM 199807 EW 19980701 AB Sporadic inclusion-body myositis (s-IBM) is the most common progressive muscle disease of older persons. The muscle biopsy demonstrates \*\*\*mononuclear\*\*\* cell inflammation and vacuolated muscle fibers containing paired helical filaments and 6- to 10-nm fibrils, both resembling those of Alzheimer disease brain and Congo red positivity. The term hereditary inclusion-body myopathies (h-IBMs) designates autosomal-recessive or autosomal-dominant disorders with muscle biopsies cytopathologically similar to s-IBM but without inflammation. Vacuolated muscle fibers of both s-IBM and the h-IBMs contain accumulations of several "Alzheimer-characteristic proteins" including beta-amyloid protein and beta-amyloid precursor protein, and their paired helical filaments are composed of phosphorylated tau. We used six well characterized antibodies against several residues of \*\*\*presenilin\*\*\* 1 (PS1) to immunostain muscle biopsies of 12 patients with s-IBM, 5 patients with autosomal-recessive inclusion-body myopathy, and 16 normal and disease controls. Seventy to eighty percent of the vacuolated muscle fibers of both s-IBM and autosomal-recessive inclusion-body myopathy had

inclusions that were strongly PS1-immunoreactive, which by immunoelectron microscopy localized mainly to paired helical filaments and 6- to 10-nm filaments. None of the control biopsies had PS1-positive inclusions characteristic of the s- and h-IBM abnormal muscle fibers. Mutations of the newly discovered PS1 gene are responsible for early-onset familial Alzheimer disease (AD), and PS1 is abnormally accumulated in sporadic and familial AD brain. Our study provides the first demonstration of PS1 abnormality in non-neural tissue and in diseases other than AD and suggests that the cytopathogenesis in AD brain and IBM muscle may share similarities.

L41 ANSWER 2 OF 2 MEDLINE DUPLICATE 2  
AN 1998330346 MEDLINE  
DN 98330346  
TI Lack of specific association of \*\*\*presenilin\*\*\* 1 (PS-1)

protein with plaques and tangles in Alzheimer's disease.  
AU Xia M Q; Bereznovska O; Kim T W; Xia W M; Liao A; Tanzi R E; Selkoe D; Hyman B T  
CS Alzheimer's Research Unit, Department of Neurology, Massachusetts General Hospital-East, Charlestown 02129, USA.  
NC AG05134 (NIA)  
AG08487 (NIA)  
AG14744 (NIA)

SO JOURNAL OF THE NEUROLOGICAL SCIENCES, (1998 Jun 11) 158 (1) 15-23.  
Journal code: JBI. ISSN: 0022-510X.

CY Netherlands  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199812  
EW 19981203  
AB Missense mutations in the \*\*\*presenilin\*\*\* -1 (PS-1) gene are causally related to the majority of familial early-onset Alzheimer's disease (FAD). PS-1 immunohistochemical expression in normal human brain and in brains with Alzheimer's disease (AD) has so far been controversial. Here, we report a study of PS-1 expression in brains, cell lines and peripheral blood \*\*\*mononuclear\*\*\* cells using a panel of well characterized PS-1-specific antibodies. These antibodies were characterized by immunofluorescent staining of PS-1 transfectants followed by flow cytometric analysis. In human brain, widespread neuronal staining

was observed. PS-1 immunoreactivity was primarily confined to neuronal cell bodies and proximal dendrites. Weaker staining of microglia was also detected, in accord with the finding of PS-1 immunoreactivity in monocytes. PS-1 expression is not particularly associated with either containing or spared from neurofibrillary tangles, nor with senile plaques. The level of PS-1 expression does not differ between normal and AD brains. Immunoprecipitation from AD, FAD and control brains revealed only a 32 kDa N-terminal fragment and an 18-20 kDa C-terminal fragment.

Little or no full length PS-1 was detected. The enriched presence of PS-1 in neurons implies an important role in neuronal function, however, the lack of apparent association of its expression with AD pathology signifies the need for a better understanding of its pathophysiological role.

=> s 11 and endothelial/ab,bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L42 10 L1 AND ENDOTHELIAL/AB,BI

=> dup rem l42

PROCESSING COMPLETED FOR L42  
L43 4 DUP REM L42 (6 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 4 ANSWERS -  
CONTINUE? Y/(N):y

L43 ANSWER 1 OF 4 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1997471338 BIOSIS  
DN PREV1997970561  
TI \*\*\*Presenilin\*\*\* 1 is involved in cerebral amyloid angiopathy

of Alzheimer's disease affected brains.  
AU Hayashi Y (1); Fukatsu R.; Tsuzuki K.; Yoshida T.; Takamaru Y.; Sasaki N.; Yamaguchi H.; Fujii N.; Takahata N.  
CS (1) Dep. Neuropsychiatry, Sapporo Med. Univ., South 1, West 16, Chuo-ku, Sapporo 060 Japan  
SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 825.

Meeting Info.: 27th Annual Meeting of the Society for Neuroscience, Part 1  
New Orleans, Louisiana, USA October 25-30, 1997  
ISSN: 0190-5295.  
DT Conference; Abstract; Conference  
LA English

L43 ANSWER 2 OF 4 MEDLINE DUPLICATE 1  
AN 97404082 MEDLINE  
DN 97404082  
TI Superoxide free radical and intracellular calcium mediate A beta(1-42)

induced \*\*\*endothelial\*\*\* toxicity.  
AU Suo Z; Fang C; Crawford F; Mullan M  
CS Department of Psychiatry, University of South Florida, Tampa 33613, USA.  
SO BRAIN RESEARCH, (1997 Jul 11) 762 (1-2) 144-52.  
Journal code: B5L. ISSN: 0006-8993.

CY Netherlands  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199712  
EW 19971201  
AB The 39-42 amino acid residue amyloid beta peptide (A beta), the major protein component in senile plaques and cerebrovascular amyloidosis in the brain in Alzheimer's disease (AD), has been shown to be neurotoxic in vitro. Accumulating data from several areas suggest that cerebrovascular dysfunction and damage may also play a significant role in the AD process.

For instance, we have recently demonstrated enhanced vasoconstriction and resistance to relaxation in intact rat aorta treated with A beta [Thomas et al., beta-Amyloid-mediated vasoactivity and vascular \*\*\*endothelial\*\*\* damage, Nature, 380 (1996) 168-171]. Significant vessel damage occurred after thirty minutes of exposure, but could be prevented with superoxide dismutase. To further investigate the role of A beta toxicity on \*\*\*endothelial\*\*\* cells, we have applied A beta peptides to cultures of human aortic \*\*\*endothelial\*\*\* cells (HAEC). Our results show that both A beta(1-42) and A beta(25-35) are toxic to HAEC in a time- and dose-dependent manner, and that this toxicity can be partially prevented by the calcium channel blocker, verapamil, and the antioxidant, superoxide dismutase. The common form of A beta, A beta(1-40), which has been shown to be neurotoxic, is much less toxic to

HAEC. A beta toxicity to HAEC occurs within 30 min of treatment with relatively lower doses than those usually observed in primary cultured neurons and vascular smooth muscle cells. It was recently reported that a variety of mutations in the beta-amyloid protein precursor gene and the \*\*\*Presenilin\*\*\* -1 and -2 genes linked to early-onset familial AD cause an increase in the plasma concentration of A beta(1-42) in mutation carriers (Scheuner et al., Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vitro by the \*\*\*presenilin\*\*\* 1 and 2 and APP mutations linked to familial Alzheimer's disease, *Nature Med.*, 2 (1996) 864-870). Human aortic \*\*\*endothelial\*\*\* cells are more sensitive to A beta(1-42) than A beta(1-40), via a pathway involving an excess of superoxide free radicals and influx of extracellular calcium. Finally, we have evidence that both apoptotic and necrotic processes are activated by the A beta peptides in these \*\*\*endothelial\*\*\* cells.

L43 ANSWER 3 OF 4 CAPLUS COPYRIGHT 1999 ACS  
AN 1998:109324 CAPLUS  
DN 128:163691  
TI Central role of oxyradicals in the mechanism of amyloid b-peptide cytotoxicity  
AU Mattson, Mark P  
CS Sanders-Brown Res. Cent. on Aging and Dep. Anatomy & Neurobiol., Univ. Kentucky, Lexington, KY, 40636-0230, USA  
SO *Alzheimer's Dis. Rev.* (1997), 2(1/2), 1-14  
CODEN: ADREFN  
URL: <http://www.coa.uky.edu/ADReview/Mattson.htm>  
PB Sanders-Brown Center on Aging, University of Kentucky  
DT Journal: General Review; (online computer file)  
LA English  
AB A review and discussion with many refs. Overwhelming evidence indicates that cells in Alzheimer's disease brain are subjected to abnormally high levels of oxidative stress, and that amyloids are a focus of cellular and mol. oxidn. Recent studies suggest that amyloid b-peptide (Ab) plays a major role in promoting oxidative stress in neurons and glial cells, and that such oxidative stress can account for many of the metabolic and neurodegenerative alterations obsd. in AD brain. Ab induces membrane lipid peroxidn. in neurons which leads to impairment of ion-motive

ATPases, and glutamate and glucose transporters. These actions of Ab lead to membrane depolarization and energy failure which, in turn, promote excitotoxic and apoptotic degenerative depolarization and energy failure which, in turn, promote excitotoxic and apoptotic degenerative cascades involving calcium overload. Membrane oxidn., as induced by Ab, also disrupts coupling of metabotropic receptors to their GTP-binding proteins, which may account for the well-known cholinergic signaling deficits and assocd. cognitive impairment in AD. 4-Hydroxynonenal, an aldehydic prodn. of membrane lipid peroxidn., is implicated as a mediator of Ab-induced disruption of cellular ion and energy homeostasis, and neuronal apoptosis. Oxidative stress induced by Ab in microglia and astrocytes likely contributes to the inflammatory process in AD brain. Moreover, Ab-mediated oxidative damage to vascular \*\*\*endothelial\*\*\* cells may contribute to the impaired glucose transport and compromised barrier function of the cerebral vessels in AD. Finally, the possible links between mutations in \*\*\*presenilin\*\*\* genes, oxidative stress, and neuronal degeneration in AD are considered.

L43 ANSWER 4 OF 4 MEDLINE DUPLICATE 2  
AN 96234265 MEDLINE  
DN 96234265  
TI Widespread neuronal expression of the \*\*\*presenilin\*\*\* -1 early-onset Alzheimer's disease gene in the murine brain.  
AU Gribbs D H; Chen L S; Bende S M; LaFerla F M  
CS Department of Neurology, University of California, Irvine, USA.  
NC P30-AG01542 (NIA)  
SO AMERICAN JOURNAL OF PATHOLOGY, (1996 Jun) 148 (6) 1797-806.  
Journal code: 3RS. ISSN: 0002-9440.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Abridged Index Medicus Journals; Priority Journals; Cancer Journals  
EM 199610  
AB Mutations in the \*\*\*presenilin\*\*\* -1 (S182) gene have been genetically linked to early-onset Alzheimer's disease. To clarify the underlying molecular mechanism through which \*\*\*presenilin\*\*\* -1 is involved in the pathogenesis of this neurodegenerative disorder, the regional and

cellular transcription profile of this gene was characterized in primary cells isolated from the murine brain by Northern blot hybridization using digoxigenin-labeled riboprobes. Our results indicate that \*\*\*presenilin\*\*\* -1 mRNA transcripts are widely distributed throughout the adult mouse brain. Furthermore, immunohistochemical labeling of hybridized sections indicates that expression was predominantly localized to neuronal cells. Neurons in the hippocampus and cerebral cortex, which are severely compromised in Alzheimer's disease, showed prominent expression of \*\*\*presenilin\*\*\* -1. In contrast, white matter areas and \*\*\*endothelial\*\*\* cells do not appear to express \*\*\*presenilin\*\*\* -1 to detectable levels. \*\*\*presenilin\*\*\* -1 transcripts, however, are also present less frequently in certain nonneuronal cell populations such as ependymal cells in the choroid plexus. Analysis of primary cells isolated from murine brain supported the results obtained by in situ hybridization and showed that cultured primary neurons and astrocytes express \*\*\*presenilin\*\*\* -1. Overall, it appears that the pattern of \*\*\*presenilin\*\*\* -1 gene expression parallels that previously described for the amyloid precursor protein.  
=> s 11 and astrocyte#/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L44 61 LI AND ASTROCYTE#/AB,BI  
=> dup rem 144  
PROCESSING COMPLETED FOR L44  
L45 26 DUP REM L44 (35 DUPLICATES REMOVED)  
=> s 144 and presenilin-2/ab,bi  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L46 27 L44 AND PRESENILIN-2/AB,BI  
=> dup rem 146  
PROCESSING COMPLETED FOR L46

L47 13 DUP REM L46 (14 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 13 ANSWERS -  
CONTINUE? Y(N):y

L47 ANSWER 1 OF 13 MEDLINE DUPLICATE

AN 1998099802 MEDLINE

DN 98099802

TI Interaction of \*\*\*presenilins\*\*\* with the filamin family of actin-binding proteins.

AU Zhang W; Han S W; McKeel D W; Goate A; Wu J Y

CS Department of Pediatrics and Molecular Biology and Pharmacology, Washington University School of Medicine, St. Louis, Missouri 63110, USA.

NC AG-05861 (NIA)

AG00634 (NIA)

AG05681 (NIA)

SO JOURNAL OF NEUROSCIENCE, (1998 Feb 1) 18 (3) 914-22.

Journal code: JDF. ISSN: 0270-6474.

CY United States

DT Journal Article; (JOURNAL ARTICLE)

LA English

EM 199804

EW 19980402

AB Mutations in \*\*\*presenilin\*\*\* genes PS1 and PS2 account for approximately 50% of early-onset familial Alzheimer's disease (FAD). The PS1 and PS2 genes encode highly homologous transmembrane proteins related to the Caenorhabditis elegans sel-12 and spe-4 gene products. A hydrophilic loop region facing the cytoplasmic compartment is likely to be functionally important because at least 14 mutations in FAD patients have been identified in this region. We report here that the loop regions of PS1 and PS2 interact with nonmuscle filamin (actin-binding protein 280, ABP280) and a structurally related protein (filamin homolog 1, Fh1). Overexpression of PS1 appears to modify the distribution of ABP280 and Fh1 proteins in cultured cells. A monoclonal antibody recognizing ABP280 and Fh1 binds to blood vessels, \*\*\*astrocytes\*\*\*, neurofibrillary tangles, neuropil threads, and dystrophic neurites in the AD brain. Detection of ABP280/Fh1 proteins in these structures suggests that these \*\*\*presenilin\*\*\*-interacting proteins may be involved in the development

of AD and that interactions between \*\*\*presenilins\*\*\* and ABP280/Fh1 may be functionally significant. The ABP280 gene is located on the human X chromosome, whereas the newly identified Fh1 gene maps to human chromosome 3. These results provide a new basis for understanding the function of \*\*\*presenilin\*\*\* proteins and further implicate cytoskeletal elements in AD pathogenesis.

L47 ANSWER 2 OF 13 BIOSIS COPYRIGHT 1999 BIOSIS

AN 1999:52671 BIOSIS

DN PREV199900052671

TI Double transgenic mice carrying mutant amyloid beta precursor protein and \*\*\*presenilin\*\*\* 1 genes express accelerated Alzheimer-like phenotype.

AU Sugaya, K. (I); Jerome, S. (I); Bryan, D.; McKinney, M.; Duff, K.; Kumar, V. (I)

CS (I) Westside VA Med. Cent., Chicago, IL 60612 USA

SO Society for Neuroscience Abstracts, (1998) Vol. 24, No. 1-2, pp. 728.

Meeting Info.: 28th Annual Meeting of the Society for Neuroscience, Part 1 Los Angeles, California, USA November 7-12, 1998 Society for Neuroscience

DT Conference

LA English

L47 ANSWER 3 OF 13 MEDLINE DUPLICATE

AN 1998267265 MEDLINE

DN 98267265

TI Cloning and characterization of the \*\*\*presenilin\*\*\* - \*\*\*2\*\*\* gene promoter.

AU Pennypacker K R; Fuldner R; Xu R; Hernandez H; Dawbarn D; Mehta N; Perez-Tur J; Baker M; Hutton M

CS Department of Pharmacology and Therapeutics, University of South Florida, Tampa, FL 33612, USA.

NC AG14633 (NIA)

SO BRAIN RESEARCH. MOLECULAR BRAIN RESEARCH, (1998 May) 56 (1-2) 57-65.

Journal code: MBR. ISSN: 0169-328X.

CY Netherlands

DT Journal Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199903

EW 19990303

AB Mutations in the \*\*\*presenilin\*\*\* - \*\*\*2\*\*\* (PS-2) have been shown to cause early onset Alzheimer's disease (AD) in a series of families known as the Volga Germans and in an unrelated Italian kindred. Expression of the PS-2 gene is regulated during AD, aging, development and brain injury. Although expressed primarily in neurons, enhanced levels of PS-2 have been reported in \*\*\*astrocytes\*\*\* activated by neuronal damage. Understanding the regulation of the PS-2 gene may thus provide an insight into its role in AD. We have isolated a 3635 bp DNA fragment that contains 2934 bp of DNA sequence upstream from the PS-2 gene. Primer extension analysis was used to map three major transcriptional start sites within the PS-2 gene. The promoter sequence, upstream of each transcriptional start site, does not contain TATA or CAAT boxes but does contain several GC rich sites (Sp-1 and AP-2). A reporter gene construct containing the PS-2 promoter (PS2P, -2934 to +702) transfected into M17 cells drives basal transcription to 20% of the levels of the SV-40 viral promoter. Addition of NGF to PC-12 cells was found to upregulate the PS2P promoter and an NGF-responsive element was localized by deletion analysis between -403 and +13 within the promoter. Since the PS-2 gene has multiple start sites and the upstream sequence is GC rich with no TATA box, the PS-2 promoter is consistent with the GC class of 'housekeeping' genes. Copyright 1998 Elsevier Science B.V.

L47 ANSWER 4 OF 13 BIOSIS COPYRIGHT 1999 BIOSIS

AN 1999:32125 BIOSIS

DN PREV199900032125

TI Endogenous \*\*\*presenilin\*\*\* : 1. \*\*\*Presenilin\*\*\* and soluble APP in primary cultures of fetal rat \*\*\*astrocytes\*\*\* and neurons: Effects of 5-HT2A/C, adenylyl cyclase, or PGE2 stimulation on normal, alternative and novel proteolytic fragments.

AU Paradis, M. D.; Lee, R. K.; Wurtman, R. J.

CS Dep. Brain Cognitive Sci., MIT, Cambridge, MA 02139 USA

SO Society for Neuroscience Abstracts, (1998) Vol. 24, No. 1-2, pp. 6.

Meeting Info.: 28th Annual Meeting of the Society for Neuroscience, Part 1 Los Angeles, California, USA November 7-12, 1998 Society for

Neuroscience  
 ISSN: 0190-5295.  
 DT Conference  
 LA English

L47 ANSWER 5 OF 13 MEDLINE DUPLICATE  
 3

AN 1998041524 MEDLINE  
 DN 98041524  
 TI Cellular expression and proteolytic processing of  
 \*\*\*presenilin\*\*\*  
 proteins is developmentally regulated during neuronal  
 differentiation.  
 AU Capell A; Saffrich R; Olivo J C; Meyn L; Walter J; Grunberg J;  
 Mathews P;  
 Nixon R; Dotti C; Haass C  
 CS Central Institute of Mental Health, Department of Molecular  
 Biology,  
 Mannheim, Germany.  
 SO JOURNAL OF NEUROCHEMISTRY, (1997 Dec) 69 (6)  
 2432-40.  
 Journal code: JAV. ISSN: 0022-3042.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199802  
 EW 19980204  
 AB We have determined the expression of the Alzheimer's  
 disease-associated  
 proteins \*\*\*presenilin\*\*\* -I and \*\*\*presenilin\*\*\* - \*\*2\*\*\*  
 in  
 primary cultures of rat hippocampal neurons. Neurons highly  
 express  
 \*\*\*presenilin\*\*\* -I and \*\*\*presenilin\*\*\* - \*\*2\*\*\* ,  
 whereas both  
 proteins were not detected in \*\*\*astrocytes\*\*\* . Further, we  
 have  
 analyzed the subcellular localization and expression in rat  
 hippocampal  
 neurons during development. Although \*\*\*presenilin\*\*\*  
 proteins were  
 localized predominantly to the endoplasmic reticulum in  
 nonneuronal cells  
 transfected with \*\*\*presenilin\*\*\* cDNAs, in neurons,  
 \*\*\*presenilin\*\*\* proteins were also found in compartments not  
 staining  
 with antibodies to grp78(BiP). \*\*\*Presenilin\*\*\* -I and  
 \*\*\*presenilin\*\*\* - \*\*2\*\*\* were predominantly detected in  
 vesicular  
 structures within the somatodendritic compartment with much less  
 expression in axons. Polarized distribution of \*\*\*presenilin\*\*\*  
 -I and  
 \*\*\*presenilin\*\*\* - \*\*2\*\*\* differs slightly, with more  
 \*\*\*presenilin\*\*\* - \*\*2\*\*\* expressed in axons compared  
 with  
 \*\*\*presenilin\*\*\* -I. \*\*\*Presenilin\*\*\* expression was found

to be  
 developmentally regulated. \*\*\*Presenilin\*\*\* expression  
 strongly  
 increased during neuronal differentiation until full morphological  
 polarization and then declined. No full-length \*\*\*presenilin\*\*\*  
 -I or  
 \*\*\*presenilin\*\*\* - \*\*2\*\*\* could be detected within cell  
 lysates. At  
 early developmental stages the expected approximately 34-kDa  
 N-terminal  
 proteolytic fragment of \*\*\*presenilin\*\*\* -I and the  
 approximately  
 38-kDa fragment of \*\*\*presenilin\*\*\* - \*\*2\*\*\* were  
 detected. Later  
 during differentiation we predominantly detected a approximately  
 38-kDa  
 fragment for \*\*\*presenilin\*\*\* -I and a approximately 42-kDa  
 fragment  
 for \*\*\*presenilin\*\*\* - \*\*2\*\*\* . By epitope mapping, we show  
 that  
 these slower migrating peptides represent N-terminal proteolytic  
 fragments, cleaved C-terminal to the conventional site of  
 processing. It  
 is noteworthy that both \*\*\*presenilin\*\*\* -I and  
 \*\*\*presenilin\*\*\* -  
 \*\*2\*\*\* undergo alternative proteolytic cleavage at the same  
 stage of  
 neuronal differentiation. Regulation of \*\*\*presenilin\*\*\*  
 expression  
 and proteolytic processing might have implications for the  
 pathological as  
 well as the biological function of \*\*\*presenilins\*\*\* during aging  
 in  
 the human brain.

L47 ANSWER 6 OF 13 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997425474 BIOSIS  
 DN PREV199799724677  
 TI \*\*\*Presenilin\*\*\* 1 (S182) and 2 (STM2) mRNA and protein  
 in sporadic  
 Alzheimer's disease (AD).  
 AU Siopa, E. G. (1); Taylor, W. (1); Rubin, B. S.; Roses, A. D.;  
 Schmechel,  
 D.; Kuo-Leblanc, V. (1); Wei, Y.; Song, P. C. (1); King, J. C.;  
 Boteva,  
 K.; Mitsuda, N.; Gilbert, J. R.; Vitek, M. P.  
 CS (1) Brown Univ., Providence, RI USA  
 SO Brain Pathology, (1997) Vol. 7, No. 4, pp. 1204.  
 Meeting Info.: Xlifth International Congress of Neuropathology  
 Perth,  
 Western Australia, Australia September 7, 1997  
 ISSN: 1015-6305.  
 DT Conference; Abstract; Conference  
 LA English

L47 ANSWER 7 OF 13 MEDLINE DUPLICATE  
 4

AN 97437409 MEDLINE  
 DN 97437409  
 TI Neuronal expression and intracellular localization of  
 \*\*\*presenilins\*\*\*  
 in normal and Alzheimer disease brains.  
 AU Hyyth D P; Vinters H V; Ho D H; Ho V V; Puls S M  
 CS Neurogenetics Laboratory, Burns and Allen Research Institute,  
 Cedars-Sinai  
 Medical Center, Los Angeles, CA 90048, USA.  
 NC P30 AG 10123 (NIA)  
 SO JOURNAL OF NEUROPATHOLOGY AND  
 EXPERIMENTAL NEUROLOGY, (1997 Sep) 56 (9)  
 1009-17.  
 Journal code: JBR. ISSN: 0022-3069.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199712  
 AB The expression patterns of \*\*\*presenilin\*\*\* 1 (PS1) and  
 \*\*\*presenilin\*\*\* \*\*2\*\*\* (PS2) in human normal and  
 Alzheimer disease  
 (AD) brains were investigated using antibodies to specific  
 N-terminal  
 peptides of PS1 (Alzh14A and Alzh14B) and PS2 (Alzh1A-AB).  
 The antibodies  
 to peptides Alzh14A (Alzh14A-AB) and Alzh14B (Alzh14B-AB)  
 detected the  
 full-length protein (approximately 63 kDa) and the  
 N-terminal-processed  
 fragment (36 kDa) of PS1, while the Alzh1A-AB detected mainly  
 the  
 N-terminal-processed fragment (36 kDa) of PS2.  
 Immunofluorescent staining  
 detected by confocal microscopy suggested that both native PS1  
 and PS2 are  
 localized mainly in the Golgi/ER apparatus. Immunohistochemical  
 studies of  
 human temporal lobes from 2 normal and 5 sporadic Alzheimer  
 brains  
 revealed high levels of PS1 and PS2 expression in the granule cell  
 layer  
 and pyramidal neurons of the hippocampus. Strong  
 immunoreactivity was  
 found in reactive \*\*\*astrocytes\*\*\* and neurofibrillary tangles of  
 all  
 5 Alzheimer brains. In contrast, only 2 sporadic Alzheimer brains  
 showed  
 \*\*\*presenilin\*\*\* -positive neuritic plaques. These observations  
 suggest  
 that \*\*\*presenilins\*\*\* may be involved in the pathology of  
 some cases  
 of sporadic AD.

L47 ANSWER 8 OF 13 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997471357 BIOSIS  
 DN PREV199799770560

TI \*\*\*Presenilin\*\*\* -1 is closely related with neurofibrillary tangles in the Alzheimer's brain.  
 AU Tomidokoro, Y.; Ishiguro, K.; Igeta, Y.; Shizuka, M.; Kawabayashi, T.; Matsubara, E.; Kanai, M.; Harigaya, Y.; Okamoto, K.; Shoji, M.  
 CS Dep. Neurol., Gunma Univ. Sch. Med., 3-39-15 Showa-machi, Maebashi, Gunma 371 Japan  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 825.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience, Part 1  
 New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295  
 DT Conference; Abstract; Conference  
 LA English

L47 ANSWER 9 OF 13 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997:517865 BIOSIS  
 DN PREV199799817068  
 TI Amyotrophic lateral sclerosis and Alzheimer's disease. Lessons from model systems.  
 AU Price, D. L. (1); Wong, P. C.; Borchelt, D. R.; Pardo, C. A.; Thinakaran, G.; Doan, A. P.; Lee, M. K.; Martin, L. J.; Sisodia, S. S.  
 CS (1) Neuropathol. Lab., Johns Hopkins Univ. Sch. Med., 558 Ross Research Bldg., 720 Rutland Ave., Baltimore, MD 21205-2196 USA  
 SO Revue Neurologique (Paris), (1997) Vol. 153, No. 8-9, pp. 484-495.  
 ISSN: 0035-3787.  
 DT General Review  
 LA English  
 SL English; French  
 AB The human neurodegenerative diseases, including motor neuron disease and Alzheimer's disease (AD), are characterized by a selective involvement of certain regions of the brain/spinal cord and of selected populations of neurons. Sporadic amyotrophic lateral sclerosis (ALS) is an age-associated disease with cytoskeletal abnormalities and death of motor neurons; familial ALS (FALS), an autosomal dominant disease linked to mutations in superoxide dismutase 1 (SOD1), is manifested by inclusions and degeneration of motor neurons. Autosomal dominant familial AD (FAD), linked to mutations in \*\*\*presenilin\*\*\* (PS1 and PS2) genes or the amyloid precursor protein (APP) gene, shows brain abnormalities (e.g., neurofibrillary tangles, deposits of cndot -amyloid A cndot , and death of subsets of neurons) similar to those that occur in sporadic AD,

the risk of which is enhanced by the presence of one or two copies of apolipoprotein E4 (apoE4) alleles. To examine the mechanisms of these diseases, investigators have used a variety of animal models, including experimentally produced, spontaneously occurring, or genetically engineered models of disease. Studies of models of degeneration of motor neurons (axotomy) and cytoskeletal abnormalities seen in motor neuron disease (i.e., axonopathy induced by cndot , cndot , iminodipropionitrile (IDPN), hereditary canine spinal muscular atrophy (HCSMA), and neurofilament NF transgenic Tg mice) have demonstrated that NF-filled swellings of axons are related to alterations in the biology of NF transport. Tg mice with SOD1 mutations, which develop the clinical features of FALS, show selective degeneration of motor neurons, which is attributed to the acquisition of toxic properties by mutant SOD1. Models of AD include: aged monkeys that show both cognitive/memory deficits and cellular abnormalities (amyloid deposition/cytoskeletal abnormalities of neurons) in cortex and hippocampus; and Tg mice that express FAD-linked genes (i.e., APP and PS1) and show increased levels of A-42, amyloid deposits, dystrophic neurites, and local responses of \*\*\*astrocytes\*\*\* and microglia. This review discusses the behavioral/neuropathological features of AD, the results of investigations of the mechanisms of disease in model systems, and the potential utility of some of these models for testing new therapies.

L47 ANSWER 10 OF 13 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997:468096 BIOSIS  
 DN PREV199799767299  
 TI Expression of Alzheimer's disease related genes in three human astrocytic cell lines.  
 AU Shepherd, C. E.; Calvert, E. L.; Cambray-Deakin, M. A.; Pearson, R. C. A.  
 CS Dep. Biomedical Sci., Univ. Sheffield, S10 2TN UK  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 266.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience, Part 1  
 New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295.  
 DT Conference; Abstract; Conference  
 LA English

L47 ANSWER 11 OF 13 MEDLINE DUPLICATE  
 5  
 AN 97220077 MEDLINE  
 DN 97220077  
 TI Expression of \*\*\*presenilin\*\*\* -1 and -2 mRNAs in rat and Alzheimer's disease brains.  
 AU Takami K., Terai K., Matsuo A.; Walker D G; McGeer P L  
 CS Pharmaceutical Development Division, Takeda Chemical Industries, Ltd., Osaka, Japan.  
 SO BRAIN RESEARCH, (1997 Feb 14) 748 (1-2) 122-30.  
 Journal code: BSL. ISSN: 0006-8993.  
 CY Netherlands  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199707  
 AB Recently, new genetic linkages have been identified for early-onset familial Alzheimer's disease (AD). Mutations have been found in the \*\*\*presenilin\*\*\* (PS)-1 (S182) gene on chromosome 14 and the PS-2 (STM2/E5-a) gene on chromosome 1. We have investigated the distribution of gene expression of both \*\*\*presenilins\*\*\* in normal rat brain, and in human control and AD cases using in situ hybridization histochemistry. In normal rat brain, intense PS-1 mRNA expression was observed predominantly in neurons, particularly hippocampal pyramidal and dentate granular neurons and cerebellar Purkinje and granular neurons. The distribution of intensely expressing PS-2 mRNA cells was similar to that of PS-1, but additional groups in the brain stem and cortex were identified. Faint but significant mRNA expression of both PS genes was detected in white matter. In control human cases, the same neuronal cell types as seen in rat brain expressed both PS mRNAs in the hippocampus and cerebellum. In AD cases, the expression of both mRNAs was markedly decreased in the hippocampus but not in the cerebellum. In addition, PS-2 hybridization showed increased mRNA expression in \*\*\*astrocyte\*\*\* -like cells in affected areas of AD cases. The present data indicate that the PS genes may play important roles in specific neurons in normal brain, and that the decreased expression in neurons in sporadic AD brain may bear some relationship to



the pathogenesis.

L47 ANSWER 12 OF 13 CAPLUS COPYRIGHT 1999 ACS  
AN 1997:227178 CAPLUS  
DN 126:275585  
TI Familial Alzheimer disease gene: \*\*\*presenilin\*\*\* 1 and 2  
AU Tabira, Takeshi  
CS Natl. Inst. Neuroscience, Tokyo, 187, Japan  
SO Shinkai Kenkyu no Shinpo (1997), 41(1), 8-17  
CODEN: SKNSAF; ISSN: 0001-8724  
PB Igaku Shoin  
DT Journal; General Review  
LA Japanese  
AB A review, with 50 refs. Since the discovery of  
\*\*\*presenilin\*\*\* 1  
(PS1) and \*\*\*presenilin\*\*\* 2 (PS2) genes, about 2 yr  
has  
passed. Over 30 point mutations in PS1 gene were found in early  
onset  
familial Alzheimer's disease families in the world, and a mutation  
of PS2  
was found in the Volga-German family. However, in our studies,  
PS1  
mutations were found in less than 20% of Japanese families of  
Alzheimer's  
disease, and none was shown to have mutations in PS2. Therefore,  
important causative or risk factor genes are still missing. PS1 and  
PS2  
genes are very homologous, and numerous isoforms are produced  
by  
alternative splicing. The full length protein of PS1 is 47 kDa  
protein  
and it is supposed to function after cleavage into two fragments.  
Both  
PS1 and PS2 are expressed mainly in Golgi and ER of neurons, but  
a part of  
\*\*\*astrocytes\*\*\* that surround vessels and senile plaques also  
contain  
this substance. It is still immature to know the function of PS1 and  
PS2,  
and the pathomechanism of Alzheimer's disease due to mutations  
of these  
genes is still unknown.  
L47 ANSWER 13 OF 13 MEDLINE DUPLICATE  
6  
AN 97081125 MEDLINE  
DN 97081125  
TI Expression of \*\*\*presenilin\*\*\* 1 and 2 (PS1 and PS2) in  
human and  
murine tissues.  
AU Lee M K; Slunt H H; Martin L J; Thinakaran G; Kim G; Gandy  
S E; Seeger M;  
Koo E; Price D L; Sisodia S S  
CS Department of Pathology, The Johns Hopkins University School  
of Medicine,  
Baltimore, Maryland 21205, USA.

NC AG05146 (NIA)  
NS20471 (NINDS)  
SO JOURNAL OF NEUROSCIENCE, (1996 Dec 1) 16 (23)  
7513-25.  
Journal code: JDF. ISSN: 0270-6474.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199703  
EW 19970302  
AB Mutations in genes encoding related proteins, termed  
\*\*\*presenilin\*\*\*  
1 (PS1) and \*\*\*presenilin\*\*\* 2 (PS2), are linked to  
the  
majority of cases with early-onset familial Alzheimer's disease  
(FAD). To  
clarify potential function(s) of \*\*\*presenilins\*\*\* and  
relationships  
of \*\*\*presenilin\*\*\* expression to pathogenesis of AD, we  
examined the  
expression of PS1 and PS2 mRNA and PS1 protein in human and  
mouse.  
Semi-quantitative PCR of reverse-transcribed RNA (RT-PCR)  
analysis  
revealed that PS1 and PS2 mRNA are expressed ubiquitously and  
at  
comparable levels in most human and mouse tissues, including  
adult brain.  
However, PS1 mRNA is expressed at significantly higher levels in  
developing brain. In situ hybridization studies of mouse embryos  
revealed  
widespread expression of PS1 mRNA with a neural expression  
pattern that,  
in part, overlaps that reported for mRNA encoding specific Notch  
homologs.  
In situ hybridization analysis in adult mouse brain revealed that  
PS1 and  
PS2 mRNAs are enriched in neurons of the hippocampal formation  
and  
entorhinal cortex. Although PS1 and PS2 mRNA are expressed  
most  
prominently in neurons, lower but significant levels of PS1 and PS2  
transcripts are also detected in white matter glial cells. Moreover,  
cultured neurons and \*\*\*astrocytes\*\*\* express PS1 and PS2  
mRNAs. Using  
PS1-specific antibodies in immunoblot analysis, we demonstrate  
that PS1  
accumulates as approximately 28 kDa N-terminal and  
approximately 18 kDa  
C-terminal fragments in brain. Immunocytochemical studies of  
mouse brain  
reveal that PS1 protein accumulates in a variety of neuronal  
populations  
with enrichment in somatodendritic and neuropil compartments.  
=> s 11 and microglia#ab,bi

'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
'AB' IS NOT A VALID FIELD CODE  
L48 19 L1 AND MICROGLIA#AB,BI  
=> dup rem 148  
PROCESSING COMPLETED FOR L48  
L49 7 DUP REM L48 (12 DUPLICATES REMOVED)  
=> d 1 - bib ab  
YOU HAVE REQUESTED DATA FROM 7 ANSWERS -  
CONTINUE? Y(N)>  
L49 ANSWER 1 OF 7 MEDLINE DUPLICATE 1  
AN 1999047653 MEDLINE  
DN 99047653  
TI Insulin-degrading enzyme regulates extracellular levels of amyloid  
beta-protein by degradation.  
AU Qiu W Q; Walsh D M; Ye Z; Vekrellis K; Zhang J; Podlasky M  
B; Rosner M R;  
Safavi A; Hersh L B; Selkoe D J  
CS Department of Neurology and Program in Neuroscience, Harvard  
Medical  
School and Center for Neurologic Diseases, Brigham and Women's  
Hospital,  
Boston, Massachusetts 02115-5716, USA.  
NC AG12749 (NIA)  
DA 02243 (NIDA)  
DO 07062 (NIDA)  
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1998 Dec 4) 273  
(49) 32730-8.  
Journal code: HIV. ISSN: 0021-9258.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199903  
EW 19990303  
AB Excessive cerebral accumulation of the 42-residue amyloid  
beta-protein  
(Abeta) is an early and invariant step in the pathogenesis of  
Alzheimer's  
disease. Many studies have examined the cellular production of  
Abeta from  
its membrane-bound precursor, including the role of the  
\*\*\*presenilin\*\*\*  
proteins therein, but almost nothing is known about how Abeta is  
degraded  
and cleared following its secretion. We previously screened  
neuronal and  
nonneuronal cell lines for the production of proteases capable of  
degrading naturally secreted Abeta under biologically relevant  
conditions

and concentrations. The major such protease identified was a metalloprotease released particularly by a \*\*\*microglial\*\*\* cell line.

BV-2. We have now purified and characterized the protease and find that it is indistinguishable from insulin-degrading enzyme (IDE), a thiol metalloendopeptidase that degrades small peptides such as insulin, glucagon, and atrial natriuretic peptide. Degradation of both endogenous and synthetic Abeta at picomolar to nanomolar concentrations was completely inhibited by the competitive IDE substrate, insulin, and by two other IDE inhibitors. Immunodepletion of conditioned medium with an IDE antibody removed its Abeta-degrading activity. IDE was present in BV-2 cytosol, as expected, but was also released into the medium by intact, healthy cells. To confirm the extracellular occurrence of IDE in vivo, we identified intact IDE in human cerebrospinal fluid of both normal and Alzheimer subjects. In addition to its ability to degrade Abeta, IDE activity was unexpectedly found to be associated with a time-dependent oligomerization of synthetic Abeta at physiological levels in the conditioned media of cultured cells; this process, which may be initiated by IDE-generated proteolytic fragments of Abeta, was prevented by three different IDE inhibitors. We conclude that a principal protease capable of down-regulating the levels of secreted Abeta extracellularly is IDE.

L49 ANSWER 2 OF 7 MEDLINE DUPLICATE 2  
AN 1998441028 MEDLINE  
DN 98441028  
TI [Alzheimer disease. Epidemiology, genetics and physiopathological hypotheses].  
MA Maladie d'Alzheimer. Epidemiologie, genetique et hypotheses physiopathologiques.  
AU Blain H; Jeandel C  
CS Service de Medecine B, CHU Nancy-Brabois, Vandoeuvre.  
SO PRESSE MEDICALE (1998 Apr 18) 27 (15) 725-30. Ref: 99  
Journal code: PMT. ISSN: 0755-4982.  
CY France  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, ACADEMIC)  
LA French  
FS Priority Journals; Cancer Journals  
EM 199901  
EW 19990104  
AB RISK FACTORS: Aging is the chief risk factor for Alzheimer's disease (AD).  
Other risk factors are aluminum in drinking water, diabetes

mellitus, head trauma. Protective factors are: higher education, cigarette smoking, nonsteroidal anti-inflammatory drugs and estrogen use. GENETIC FACTORS: Mutations of \*\*\*presenilins\*\*\* 1 and 2 and of the APP gene in families with early-onset AD. Apolipoprotein E polymorphism in late-onset familial Alzheimer disease and sporadic AD. PATHOGENIC HYPOTHESES: Amyloid deposits in senile plaques and therefore dementia could be due to an overproduction of Abeta (Down's syndrome) or due to the primary (APP mutation) or secondary (role of diabetes, mellitus, apoE polymorphism: protective effect of estrogen) abnormal neurotoxic feature of Abeta. The hyperphosphorylation of tau (a protein which plays a pivotal role in the axonal transport), perhaps regulated by the apoE polymorphism could lead to neurofibrillar degeneration. Neurotoxic mediators produced by the activated \*\*\*microglia\*\*\* (perhaps activated by neuronal damage) and oxidative stress could also be involved in the neurodegeneration.

L49 ANSWER 3 OF 7 MEDLINE  
AN 1998210340 MEDLINE  
DN 98210340  
TI Molecular physiopathology of Alzheimer's disease.  
AU Yamada T  
CS Department of Neurology, Faculty of Medicine, Kyushu University, Fukuoka.  
SO FUKUOKA IGAKU ZASSHI. FUKUOKA ACTA MEDICA, (1998 Feb) 89 (2) 29-33. Ref: 5  
Journal code: F8R. ISSN: 0016-254X.  
CY Japan  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, TUTORIAL)  
LA Japanese  
EM 199808  
EW 19980803  
L49 ANSWER 4 OF 7 MEDLINE DUPLICATE 3  
AN 1998330346 MEDLINE  
DN 98330346  
TI Lack of specific association of \*\*\*presenilin\*\*\* 1 (PS-1) protein with plaques and tangles in Alzheimer's disease.  
AU Xia M Q; Bereznikova O; Kim T W; Xia W M; Liao A; Tanzi R E; Selkoe D; Hyman B T  
CS Alzheimer's Research Unit, Department of Neurology, Massachusetts General Hospital-East, Charlestown 02129, USA.  
NC AG05134 (NIA)

AG08487 (NIA)  
AG14744 (NIA)  
SO JOURNAL OF THE NEUROLOGICAL SCIENCES, (1998 Jun 1) 158 (1) 15-23.  
Journal code: JBJ. ISSN: 0022-510X.  
CY Netherlands  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199812  
EW 19981203  
AB Missense mutations in the \*\*\*presenilin\*\*\* -1 (PS-1) gene are causally related to the majority of familial early-onset Alzheimer's disease (FAD).  
PS-1 immunohistochemical expression in normal human brain and in brains with Alzheimer's disease (AD) has so far been controversial. Here, we report a study of PS-1 expression in brains, cell lines and peripheral blood mononuclear cells using a panel of well characterized PS-1-specific antibodies. These antibodies were characterized by immunofluorescent staining of PS-1 transfectants followed by flow cytometric analysis.  
In human brain, widespread neuronal staining was observed. PS-1 immunoreactivity was primarily confined to neuronal cell bodies and proximal dendrites. Weaker staining of \*\*\*microglia\*\*\* was also detected, in accord with the finding of PS-1 immunoreactivity in monocytes. PS-1 expression is not particularly associated with neurons either containing or spared from neurofibrillary tangles, nor with senile plaques. The level of PS-1 expression does not differ between normal and AD brains. Immunoprecipitation from AD, FAD and control brains revealed only a 32 kDa N-terminal fragment and an 18-20 kDa C-terminal fragment.  
Little or no full length PS-1 was detected. The enriched presence of PS-1 in neurons implies an important role in neuronal function, however, the lack of apparent association of its expression with AD pathology signifies the need for a better understanding of its pathophysiological role.

L49 ANSWER 5 OF 7 MEDLINE DUPLICATE 4  
AN 1998348622 MEDLINE  
DN 98348622  
TI Amyotrophic lateral sclerosis and Alzheimer disease. Lessons from model systems.  
AU Price D L; Wong P C; Borchelt D R; Pardo C A; Thinakaran G;

- Doan A P; Lee  
M K; Martin L J; Sisodia S S  
CS Department of Pathology, Johns Hopkins, University School of Medicine,  
Baltimore, Maryland 21205-2196, USA.  
NC NS 20471 (NINDS)  
AG 05146 (NIA)  
NS 10580 (NINDS)  
+  
SO REVUE NEUROLOGIQUE, (1997 Sep) 153 (8-9) 484-95. Ref: 143  
Journal code: SU9. ISSN: 0035-3787.  
CY France  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, ACADEMIC)  
LA English  
FS Priority Journals  
EM 199810  
EW 19981003  
AB The human neurodegenerative diseases, including motor neuron disease and Alzheimer's disease (AD), are characterized by a selective involvement of certain regions of the brain/spinal cord and selected populations of neurons. Sporadic amyotrophic lateral sclerosis (ALS) is an age-associated disease with cytoskeletal abnormalities and death of motor neurons; familial ALS (FALS), an autosomal dominant disease linked to mutations in superoxide dismutase 1 (SOD1), is manifested by inclusions and degeneration of motor neurons. Autosomal dominant familial AD (FAD), linked to mutations in **presenilin\*\*\*** (PS1 and PS2) genes or the amyloid precursor protein (APP) gene, shows brain abnormalities (e.g., neurofibrillary tangles, deposits of  $\beta$ -amyloid A., and death of subsets of neurons) similar to those that occur in sporadic AD, the risk of which is enhanced by the presence of one or two copies of apolipoprotein E4 (apoE4) alleles. To examine the mechanisms of these diseases, investigators have used a variety of animal models, including experimentally produced, spontaneously occurring, or genetically engineered models of disease. Studies of models of degeneration of motor neurons (axotomy) and cytoskeletal abnormalities seen in motor neuron disease (i.e., axonopathy induced by  $\beta$ -iminodipionitrile (IDPN), hereditary canine spinal muscular atrophy (HCSMA), and neurofilament NF transgenic Tg mice) have demonstrated that NF-filled swellings of axons are related to
- alterations in the biology of NF transport. Tg mice with SOD1 mutations, which develop the clinical features of FALS, show selective degeneration of motor neurons, which is attributed to the acquisition of toxic properties by mutant SOD1. Models of AD include: aged monkeys that show both cognitive/memory deficits and cellular abnormalities (amyloid deposition/cytoskeletal abnormalities of neurons) in cortex and hippocampus; and Tg mice that express mutant human FAD-linked genes (i.e., APP and PS1) and show increased levels of A $\beta$ 42, amyloid deposits, dystrophic neurites, and local responses of astrocytes and **microglia\*\*\***. This review discusses the behavioral/neuropathological features of AD, the results of investigations of mechanisms of disease in model systems, and potential utility of some of these models for testing new therapies.
- L49 ANSWER 6 OF 7 MEDLINE DUPLICATE 5  
AN 97343985 MEDLINE  
DN 97343985  
TI Immunoreactivity of **presenilin\*\*\*** -1 in human, rat and mouse brain.  
AU Kim K S; Wegiel J; Sapienza V; Chen J; Hong H; Wisniewski H M  
CS New York State Institute for Basic Research in Developmental Disabilities,  
Staten Island 10314, USA.  
NC P01-AG111531 (NIA)  
SO BRAIN RESEARCH, (1997 May 16) 757 (1) 159-63.  
Journal code: B5L. ISSN: 0006-8993.  
CY Netherlands  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199710  
AB Monoclonal antibodies (mAbs) D3G6 and C8A5, specific for amino acid residues 160-168 of S182 protein, immunolabeled neurons, ependymal and choroid plexus cells, and myocytes in brain sections from normal subjects and people with Alzheimer disease or Down syndrome and in rats and mice. Oligodendroglia, **microglia\*\*\***, and the majority of astrocytes were negative. S182 protein or a fragment of the protein detected with these mAbs is not a constituent of amyloid-beta deposits or tangles.
- L49 ANSWER 7 OF 7 CAPLUS COPYRIGHT 1999 ACS  
AN 1998:109324 CAPLUS  
DN 128:163691
- T1 Central role of oxyradicals in the mechanism of amyloid  $\beta$ -peptide cytotoxicity  
AU Mattson, Mark P.  
CS Sanders-Brown Res. Cent. on Aging and Dep. Anatomy & Neurobiol., Univ. Kentucky, Lexington, KY, 40636-0230, USA  
SO Alzheimer's Dis. Rev. (1997), 2(1/2), 1-14  
CODEN: ADREFN  
URL: <http://www.coa.uky.edu/ADReview/Mattson.htm>  
PB Sanders-Brown Center on Aging, University of Kentucky  
DT Journal; General Review; (online computer file)  
LA English  
AB A review and discussion with many refs. Overwhelming evidence indicates that cells in Alzheimer's disease brain are subjected to abnormally high levels of oxidative stress, and that amyloids are a focus of cellular and mol. oxidn. Recent studies suggest that amyloid  $\beta$ -peptide (Ab) plays a major role in promoting oxidative stress in neurons and glial cells, and that such oxidative stress can account for many of the metabolic and neurodegenerative alterations obsd. in AD brain. Ab induces membrane lipid peroxidn. in neurons which leads to impairment of ion-motive ATPases, and glutamate and glucose transporters. These actions of Ab lead to membrane depolarization and energy failure which, in turn, promote excitotoxic and apoptotic degenerative depolarization and energy failure which, in turn, promote excitotoxic and apoptotic degenerative cascades involving calcium overload. Membrane oxidn., as induced by Ab, also disrupts coupling of metabotropic receptors to their GTP-binding proteins, which may account for the well-known cholinergic signaling deficits and assoc. cognitive impairment in AD. 4-Hydroxynonenal, an aldehydic prodn. of membrane lipid peroxidn., is implicated as a mediator of Ab-induced disruption of cellular ion and energy homeostasis, and neuronal apoptosis. Oxidative stress induced by Ab in **microglia\*\*\*** and astrocytes likely contributes to the inflammatory process in AD brain. Moreover, Ab-mediated oxidative damage to vascular endothelial cells may contribute to the impaired glucose transport and compromised barrier function of the cerebral vessels in AD. Finally, the possible mechanistic links between mutations in **presenilin\*\*\*** genes, oxidative stress, and



Thus, the development of AD-like pathology is substantially enhanced when a PS1 mutation, which causes a modest increase in A beta42(43), is introduced into Tg2576-derived mice. Remarkably, both doubly and singly transgenic mice showed reduced spontaneous alternation performance in a "Y" maze before substantial A beta deposition was apparent. This suggests that some aspects of the behavioral phenotype in these mice may be related to an event that precedes plaque formation.

L53 ANSWER 4 OF 7 CAPLUS COPYRIGHT 1999 ACS  
 AN 1998:1559 CAPLUS  
 DN 128:73898  
 TI Transgenic animals expressing perlecan and amyloid genes at high levels and methods of identifying compounds for the treatment of amyloidoses  
 IN Snow, Alan; Fukuchi, Ken-ichiro; Hassell, John  
 PA University of Washington, USA  
 SO PCT Int. Appl., 146 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1  
 PATENT NO. KIND DATE APPLICATION NO.  
 DATE

PI WO 9746664 A1 19971211 WO 97-US9875  
 19970606  
 W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE,  
 ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR,  
 LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU,  
 SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, AM, AZ,  
 BY, KG, KZ, MD, RU, TJ, TM  
 RW: GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB,  
 GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN,  
 ML, MR, NE, SN, TD, TG  
 AU 9736402 A1 19980105 AU 97-36402 19970606  
 PRAI US 96-17830 19960606  
 WO 97-US9875 19970606  
 AB Transgenic animals expressing a foreign gene for a perlecan, or genes for perlecan and an amyloid are constructed for use in the testing of compds. that can alter the rate or extent of amyloid deposition.  
 Over-expression

of perlecan and amyloid proteins results in animals showing symptoms closer to amyloidoses than found in animals only over-expressing an amyloid gene, esp. Alzheimer's disease. Over-expression of a gene encoding domains 1-5 of mouse perlecan and the 695-amino acid isoform of beta-amyloid in P19 cells led to an up-regulation of beta-amyloid synthesis and secretion. P19 cells induced to form neurons degenerated when the perlecan gene was overexpressed.

L53 ANSWER 5 OF 7 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997:468096 BIOSIS  
 DN PREV199799767299  
 TI Expression of Alzheimer's disease related genes in three human astrocytic cell lines.  
 AU Shepherd, C. E.; Calvert, E. L.; Cambray-Deakin, M. A.; Pearson, R. C. A.  
 CS Dep. Biomedical Sci., Univ. Sheffield, S10 2TN UK  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 266.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience, Part 1  
 New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295.  
 DT Conference; Abstract; Conference  
 LA English

L53 ANSWER 6 OF 7 MEDLINE DUPLICATE 2  
 AN 97347186 MEDLINE  
 DN 97347186  
 TI Immunohistochemical analysis of \*\*\*presenilin\*\*\* \*\*2\*\*\* expression in the mouse brain: distribution pattern and co-localization with \*\*\*presenilin\*\*\* 1 protein.  
 AU Blanchard V; Czech C; Bonici B; Clavel N; Gohin M; Dalet K; Revah F;  
 Pradier L; Imperato A; Moussouli S  
 CS Rhone-Poulenc Rorer, Centre de Recherche de Vitry-Alfortville, Vitry-sur-Seine, France.  
 SO BRAIN RESEARCH, (1997 May 30) 758 (1-2) 209-17.  
 Journal code: B5L. ISSN: 0006-8993.  
 CY Netherlands  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 199711  
 EW 19971103  
 AB Missense mutations of \*\*\*presenilin\*\*\* 1 (PS-1) and \*\*\*presenilin\*\*\* \*\*2\*\*\* (PS-2) genes cause the majority of early-onset familial forms of Alzheimer's disease (AD). We previously characterized the distribution of the PS-1 protein in the mouse brain by immunohistochemistry

using an antibody directed against an epitope located in the large hydrophilic loop [Moussouli, S., Czech, C., Pradier, L., Blanchard, V., Bonici, B., Gohin, M., Imperato, A. and Revah, F., Immunohistochemical analysis of \*\*\*presenilin\*\*\* 1 expression in the mouse brain, FEBS Lett., 383 (1996) 219-222]. Similarly, we now report the distribution pattern of PS-2 protein in the mouse brain. For these experiments we used a polyclonal antibody raised against a synthetic peptide corresponding to the amino-acid sequence 7-24 of the predicted human PS-2 protein. The specificity of the antibody was evidenced by its ability to recognize PS-2 protein in immunoprecipitation studies and by antigen-peptide competition. In the mouse brain, PS-2 protein was present in numerous cerebral structures, but its distribution in these structures did not correlate with their susceptibility to AD pathology. In all examined structures of the gray matter, PS-2 protein was concentrated in neuronal cell bodies but it was not detected in the \*\*\*glial\*\*\* cells of the white matter. The regional distribution pattern of PS-2 protein was almost identical to that of PS-1 protein. Moreover, PS-2 protein co-localized with PS-1 in a large number of neuronal cell bodies. In terms of subcellular localization, PS-2 immunostaining was present almost exclusively in neuronal cell bodies while PS-1 immunostaining was also present in dendrites. This could be explained by the different epitopes of the antibodies and the known proteolytic processing of both \*\*\*presenilins\*\*\* in vivo [Tanzi, R.E., Kovacs, D.M., Kim, T.-W., Moir, R.D., Guenette, S.Y. and Wasco, W., The \*\*\*presenilin\*\*\* genes and their role in early-onset familial Alzheimer's disease, Alzheimer's disease Rev., 1 (1996) 91-98]. Within neuronal cell bodies, the immunostaining of PS-2 protein, as well as that of PS-1 protein, had a reticular and granular appearance. This suggests in agreement with previous observations on PS-1 and PS-2 in COS and H4 cells [Kovacs, D.M., Faustet, H.J., Page, K.J., Kim, T.-W., Moir, R.D., Merriam, D.E., Hollister, R.D., Hallmark, O.G., Mancini, R., Felsenstein, K.M., Hyman, B.T., Tanzi, R.E., Wasco, W., Alzheimer-associated \*\*\*presenilins\*\*\* 1 and 2: neuronal expression in brain and localization to intracellular membranes in mammalian cells, Nature Med., 2 (1996) 224-229] that these proteins are situated in intracytoplasmic organelles, possibly the

endoplasmic reticulum and the Golgi complex.

L53 ANSWER 7 OF 7 MEDLINE DUPLICATE 3

AN 97081125 MEDLINE

DN 97081125

TI Expression of \*\*\*presenilin\*\*\* 1 and 2 (PS1 and PS2) in human and

murine tissues.

AU Lee M K; Slunt H H; Martin L J; Thinakaran G; Kim G; Gandy S E; Seeger M;

Koo E; Price D L; Sisodia S S

CS Department of Pathology, The Johns Hopkins University School of Medicine,

Baltimore, Maryland 21205, USA.

NC AG05146 (NIA)

NS20471 (NINDS)

SO JOURNAL OF NEUROSCIENCE, (1996 Dec 1) 16 (23)

7513-25.

Journal code: JDF. ISSN: 0270-6474.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199703

EW 19970302

AB Mutations in genes encoding related proteins, termed

\*\*\*presenilin\*\*\*

1 (PS1) and \*\*\*presenilin\*\*\* \*\*2\*\*\* (PS2), are linked to the

majority of cases with early-onset familial Alzheimer's disease

(FAD). To

clarify potential function(s) of \*\*\*presenilins\*\*\* and relationships

of \*\*\*presenilin\*\*\* expression to pathogenesis of AD, we

examined the

expression of PS1 and PS2 mRNA and PS1 protein in human and mouse.

Semi-quantitative PCR of reverse-transcribed RNA (RT-PCR)

revealed that PS1 and PS2 mRNA are expressed ubiquitously and

at

comparable levels in most human and mouse tissues, including adult brain.

However, PS1 mRNA is expressed at significantly higher levels in developing brain. In situ hybridization studies of mouse embryos revealed

widespread expression of PS1 mRNA with a neural expression

pattern that,

in part, overlaps that reported for mRNA encoding specific Notch

homologs.

In situ hybridization analysis in adult mouse brain revealed that PS1 and

PS2 mRNAs are enriched in neurons of the hippocampal formation

and

entorhinal cortex. Although PS1 and PS2 mRNA are expressed

most

prominently in neurons, lower but significant levels of PS1 and PS2

transcripts are also detected in white matter \*\*\*glia\*\*\* cells. Moreover, cultured neurons and astrocytes express PS1 and PS2 mRNAs. Using

PS1-specific antibodies in immunoblot analysis, we demonstrate that PS1

accumulates as approximately 28 kDa N-terminal and

approximately 18 kDa

C-terminal fragments in brain. Immunocytochemical studies of

mouse brain

reveal that PS1 protein accumulates in a variety of neuronal

populations

with enrichment in somatodendritic and neuropil compartments.

=> s 11 and neuron?/ab,bi

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

L54 500 L1 AND NEURON?/AB,BI

=> s 154 and presenilin-2/ab,bi

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

L55 135 L54 AND PRESENILIN-2/AB,BI

=> s 155 and (cell death or apoptosis)/ab,bi

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

'AB' IS NOT A VALID FIELD CODE

L56 43 L55 AND (CELL DEATH OR APOPTOSIS)/AB,BI

=> dup rem 156

PROCESSING COMPLETED FOR L56

L57 23 DUP REM L56 (20 DUPLICATES REMOVED)

=> d 1- bib ab

YOU HAVE REQUESTED DATA FROM 23 ANSWERS - CONTINUE? Y(N)?y

L57 ANSWER 1 OF 23 MEDLINE DUPLICATE

1

AN 1999145560 MEDLINE

DN 99145560

TI Phosphorylation of \*\*\*presenilin\*\*\* - \*\*2\*\*\* regulates its cleavage

by caspases and retards progression of \*\*\*apoptosis\*\*\*

AU Walter J; Schindzielorz A; Grunberg J; Haass C

CS Central Institute of Mental Health, Department of Molecular

Biology, J5.

68159 Mannheim, Germany.

SO PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1999 Feb 16) 96 (4) 1391-6.

Journal code: PV3. ISSN: 0027-8424.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals; Cancer Journals

EM 199905

EW 19990504

AB Mutations within the \*\*\*Presenilin\*\*\* - \*\*2\*\*\* (PS-2)

gene are

associated with early onset familial Alzheimer's disease. The gene

encodes

a polytopic transmembrane protein that undergoes endoproteolytic processing resulting in the generation of N-terminal and C-terminal

fragments (CTFs). PS-2 is also cleaved by proteases of the caspase

family

during apoptotic \*\*\*cell\*\*\* \*\*\*death\*\*\* . CTFs of PS-2

were shown

to inhibit \*\*\*apoptosis\*\*\* , suggesting an important role in the

regulation of programmed \*\*\*cell\*\*\* \*\*\*death\*\*\* . Recently,

we

found that the CTF of PS-2 is phosphorylated in vivo. We mapped

the in

vivo phosphorylation sites of PS-2 to serine residues 327 and 330,

which

are localized immediately adjacent to the cleavage sites of caspases

after

aspartate residues 326 and 329. Phosphorylation of PS-2 inhibits its cleavage by caspase-3. This effect can be mimicked by substitutions

of

serines 327 and 330 by aspartate or glutamate. In addition, the

uncleavable form of PS-2 CTF was found to enhance its

antiapoptotic

properties, leading to a slower progression of \*\*\*apoptosis\*\*\* .

These

results demonstrate that PS-2 cleavage as well as its function in

\*\*\*apoptosis\*\*\* can be regulated by protein phosphorylation.

Alterations

in the phosphorylation of PS-2 may therefore promote the

pathogenesis of

AD by affecting the susceptibility of \*\*\*neurons\*\*\* to apoptotic

stimuli.

L57 ANSWER 2 OF 23 MEDLINE DUPLICATE

2

AN 1999098950 MEDLINE

DN 99098950

TI Contrasting role of \*\*\*presenilin\*\*\* -1 and \*\*\*presenilin\*\*\*

- \*\*2\*\*\* in \*\*\*neuronal\*\*\* differentiation in vitro.

AU Hong C S; Caromile L; Nomata Y; Mort H; Bredeisen D E; Koo

E H

CS Department of Neurosciences, University of California, San

Diego, La  
Jolla, California 92093, USA.  
NC NS01812 (NINDS)  
AG12282 (NIA)  
SO JOURNAL OF NEUROSCIENCE, (1999 Jan 15) 19 (2) 637-43.  
Journal code: JDF. ISSN: 0270-6474.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199904  
EW 19990403  
AB \*\*\*Presenilin\*\*\* -1 (PS1) and \*\*\*presenilin\*\*\* . \*\*\*2\*\*\*  
(PS2),  
the major genes of familial Alzheimer's disease, are homologous to  
sel-12,  
a Caenorhabditis elegans gene involved in cell fate decision during  
development. Recently, wild-type and mutant \*\*\*presenilins\*\*\*  
have  
been associated also with apoptotic \*\*\*cell\*\*\* \*\*\*death\*\*\*.  
By  
using stable transfection of antisense cDNAs, we studied the  
functions of  
PS1 and PS2 during \*\*\*neuronal\*\*\* differentiation in the  
NTera2 human  
teratocarcinoma (NT2) cell line. Expression of antisense PS1  
resulted in a  
failure of the clones to differentiate into \*\*\*neurons\*\*\* after  
retinoic acid induction, whereas cells transfected with antisense  
PS2  
differentiated normally. Concomitantly, antisense PS1 clones were  
associated with increased \*\*\*apoptosis\*\*\* both under basal  
conditions  
and during the early period of \*\*\*neuronal\*\*\* differentiation  
after  
retinoic acid treatment. Overexpression of bel-2 in antisense PS1  
clones  
reduced \*\*\*cell\*\*\* \*\*\*death\*\*\* and resulted in a recovery  
of  
\*\*\*neuronal\*\*\* differentiation. These studies suggest that PS1  
plays a  
role in differentiation and \*\*\*cell\*\*\* \*\*\*death\*\*\* and that  
PS1  
and PS2 have differing physiological roles in this experimental  
paradigm.  
L57 ANSWER 3 OF 23 EMBASE COPYRIGHT 1999 ELSEVIER  
SCI.B.V.DUPLICATE 3  
AN 1999088629 EMBASE  
TI The \*\*\*presenilins\*\*\*  
AU Mattson M.P.; Guo Q.  
CS Dr. M.P. Mattson, 211 Sanders-Brown Building, University of  
Kentucky,  
Lexington, KY 40536-0230, United States.  
nmattson@iging.coa.uky.edu  
SO Neuroscientist, (1999) 5/2 (112-124).

Refs: 61  
ISSN: 1073-8584 CODEN: NROSFJ  
CY United States  
DT Journal; General Review  
FS 008 Neurology and Neurosurgery  
LA English  
SL English  
AB \*\*\*Presenilin\*\*\* -1 and \*\*\*presenilin\*\*\* - \*\*\*2\*\*\* are  
highly  
homologous genes located on chromosomes 14 and 1, respectively,  
that have  
recently been linked to some cases of early-onset autosomal  
dominant  
inherited forms of Alzheimer's disease (AD). \*\*\*Presenilins\*\*\*  
are  
integral membrane proteins localized in the endoplasmic reticulum  
of  
\*\*\*neurons\*\*\* throughout the nervous system. Studies of  
\*\*\*presenilin\*\*\* -1 knockout mice, and of invertebrate  
homologues of  
\*\*\*presenilins\*\*\* and their interacting proteins, suggest major  
roles  
for \*\*\*presenilins\*\*\* in normal development.  
\*\*\*Presenilin\*\*\* -1  
mutant knockin mice do not exhibit developmental abnormalities,  
which  
indicates that the pathogenic mechanism of \*\*\*presenilin\*\*\*  
mutations  
involves gain of an adverse property of the mutant protein.  
Expression of  
\*\*\*presenilin\*\*\* mutations in cultured \*\*\*neurons\*\*\* and  
transgenic  
mice results in increased sensitivity to \*\*\*apoptosis\*\*\* induced  
by  
trophic factor withdrawal and exposure to oxidative and metabolic  
insults,  
and also alters gene expression. The pathogenic mechanism of  
\*\*\*presenilin\*\*\* mutations may involve perturbed endoplasmic  
reticulum  
calcium homeostasis resulting in enhanced oxidative stress, altered  
proteolytic processing of the amyloid precursor protein (APP), and  
increased \*\*\*neuronal\*\*\* vulnerability to excitotoxicity.  
Studies of  
\*\*\*presenilins\*\*\* are rapidly increasing our understanding the  
molecular  
and cellular underpinnings of AD and are also elucidating novel  
roles of  
the endoplasmic reticulum in \*\*\*neuronal\*\*\* plasticity and  
\*\*\*cell\*\*\* \*\*\*death\*\*\*.  
L57 ANSWER 4 OF 23 EMBASE COPYRIGHT 1999 ELSEVIER  
SCI. B.V.  
AN 1999049702 EMBASE  
TI Alzheimer's disease and stroke.  
AU Dineley K.; Denner L.  
CS L. Denner, Texas Biotechnology Corporation, 7000 Fannin,  
Houston, TX

77030, United States. ldenner@tbc.com  
SO Drugs, (1999) 2/1 (7-8).  
ISSN: 1369-7056 CODEN: IDRUFN  
CY United Kingdom  
DT Journal; Conference Article  
FS 005 General Pathology and Pathological Anatomy  
008 Neurology and Neurosurgery  
037 Drug Literature Index  
LA English  
SL English  
AB This report focuses on the two most common neurological  
diseases in man:  
Alzheimer's disease (AD) and stroke. One common feature of both  
of these  
diseases is the death of cells, particularly \*\*\*neurons\*\*\*. Since  
a  
typical mechanism of \*\*\*cell\*\*\* \*\*\*death\*\*\* is  
\*\*\*apoptosis\*\*\*,  
, this will be an additional focal point in this report.  
L57 ANSWER 5 OF 23 MEDLINE  
AN 1998401687 MEDLINE  
DN 98401687  
TI Is \*\*\*apoptosis\*\*\* key in Alzheimer's disease? [news] [see  
comments].  
CM Comment in: Science 1998 Nov 13;282(5392):1268-9  
AU Barnaga M  
SO SCIENCE, (1998 Aug 28) 281 (5381) 1303-4.  
Journal code: U7. ISSN: 0036-8075.  
CY United States  
DT News Announcement  
LA English  
FS Cancer Journals; Priority Journals  
EM 199811  
L57 ANSWER 6 OF 23 MEDLINE  
4  
AN 1998442695 MEDLINE  
DN 98442695  
TI Calsenilin: a calcium-binding protein that interacts with the  
\*\*\*presenilin\*\*\* and regulates the levels of a  
\*\*\*presenilin\*\*\*  
fragment [see comments].  
CM Comment in: Nat Med 1998 Oct;4(10):1127-8  
AU Buxbaum J D; Choi E K; Luo Y; Lilliehook C; Crowley A C;  
Merriam D E;  
Wasco W  
CS Department of Psychiatry, Mount Sinai School of Medicine, New  
York, New  
York 10029, USA. bxbaj01@doc.mssm.edu  
SO NATURE MEDICINE, (1998 Oct) 4 (10) 1177-81.  
Journal code: CG5. ISSN: 1078-8956.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199901

EW 19990104  
AB Most early-onset familial Alzheimer disease (AD) cases are caused by mutations in the highly related genes \*\*\*presenilin\*\*\* 1 (PS1) and \*\*\*presenilin\*\*\* 2\*\*\* (PS2). \*\*\*Presenilin\*\*\* mutations produce increases in beta-amyloid (A $\beta$ ) formation and \*\*\*apoptosis\*\*\* in many experimental systems. A cDNA (ALG-3) encoding the last 103 amino acids of PS2 has been identified as a potent inhibitor of \*\*\*apoptosis\*\*\*. Using this PS2 domain in the yeast two-hybrid system, we have identified a \*\*\*neural\*\*\* protein that binds calcium and \*\*\*presenilin\*\*\*, which we call calsenilin. Calsenilin interacts with both PS1 and PS2 in cultured cells, and can regulate the levels of a proteolytic product of PS2. Thus, calsenilin may mediate the effects of wild-type and mutant \*\*\*presenilins\*\*\* on \*\*\*apoptosis\*\*\* and on A $\beta$  formation. Further characterization of calsenilin may lead to an understanding of the normal role of the \*\*\*presenilins\*\*\* and of the role of the \*\*\*presenilins\*\*\* in Alzheimer disease.

L57 ANSWER 7 OF 23 MEDLINE DUPLICATE  
5 AN 199902590 MEDLINE  
DN 9902590  
TI Familial Alzheimer's disease: oxidative stress, beta-amyloid, \*\*\*presenilins\*\*\*, and \*\*\*cell\*\*\* death\*\*\*  
AU Velez-Pardo C; Jimenez Del Rio M; Lopera F  
CS Department of Neurology, University Hospital, Medellin, Colombia.  
SO GENERAL PHARMACOLOGY, (1998 Nov) 31 (5) 675-81.  
Ref: 108  
Journal code: FLK. ISSN: 0306-3623.  
CY ENGLAND: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, TUTORIAL)  
LA English  
FS Priority Journals  
EM 199902  
EW 19990204  
AB 1. The basic etiology of Alzheimer's disease remains unknown, although four genes have so far been involved: beta-amyloid precursor protein, \*\*\*presenilin\*\*\* -1, \*\*\*presenilin\*\*\* - 2\*\*\* and apolipoprotein E genes. 2. The largest familial Alzheimer's disease (FAD) kindred so far

reported belong to a point mutation in codon 280 that results in a glutamic acid-to-alanine substitution in \*\*\*presenilin\*\*\* -1 characterized in Antioquia, Colombia. 3. A hypothetical unified molecular mechanism model of \*\*\*cell\*\*\* death\*\*\* in FAD mediated by \*\*\*presenilin\*\*\* -1, beta-amyloid, and oxidative stress is proposed as an attempt to explain the mechanisms of \*\*\*neural\*\*\* loss in this neurodegenerative disorder.

L57 ANSWER 8 OF 23 MEDLINE  
AN 1998439039 MEDLINE  
DN 98439039  
TI \*\*\*Presenilins\*\*\* -in search of functionality.  
AU Kattan E H; Allsop D; Christie G; Davis J; Gray C; Mansfield F; Ward R V  
CS Neurosciences Research, SmithKline Beecham Pharmaceuticals, New Frontiers  
Science Park, Harlow, Essex, UK.  
SO BIOCHEMICAL SOCIETY TRANSACTIONS, (1998 Aug) 26 (3) 491-6. Ref: 48  
Journal code: E48. ISSN: 0300-5127.  
CY ENGLAND: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, TUTORIAL)  
LA English  
FS Priority Journals  
EM 199902  
EW 19990204  
AB The discovery of the PS proteins, the complexities of their biochemistry and their potential involvement in signalling pathways and in \*\*\*apoptosis\*\*\* have galvanized research into AD. To date, the aspect of the functionality of the PSs most relevant to the pathology of AD is the effect of PS FAD mutants to increase the proportion of A $\beta$  42 produced from cells. This, coupled to the observation that gamma-secretase cleavage is considerably reduced in \*\*\*neurons\*\*\* derived from PS-1 knockout mice, argues strongly that PS plays a very direct role in the proteolytic processing of APP.

L57 ANSWER 9 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1998:224560 BIOSIS  
DN PREV199800224560  
TI Regulation of \*\*\*apoptosis\*\*\* by \*\*\*presenilin\*\*\* 1.  
AU Wolozin, Benjamin (1); Alexander, P.; Palacino, J.  
CS (1) Dep. Pharmacol., Loyola Univ. Medical Cent., Build. 102, Room 3634,  
2160 South First Ave., Maywood, IL 60153 USA

SO Neurobiology of Aging, (Jan-Feb., 1998) Vol. 19, No. 1 SUPPL., pp. S23-S27.  
ISSN: 0197-4580.  
DT Article  
LA English  
AB Familial Alzheimer's disease is transmitted as an autosomal dominant disorder and, in 5-10% of the cases, is caused by mutations in the coding regions of two homologous genes, \*\*\*Presenilin\*\*\* 1 and 2 (PS1 and PS2). Previously, we have shown that PS2, a homolog of PS1, regulates \*\*\*apoptosis\*\*\* induced in \*\*\*neurons\*\*\* by trophic withdrawal or A $\beta$ , and in T-cells by Fas ligand. We now report that PS1 also regulates \*\*\*apoptosis\*\*\*. Both wild-type and the H15Y mutant form of PS1 enhance Fas-mediated \*\*\*apoptosis\*\*\* in Jurkat cells. We also observed that wild-type and the H15Y mutant form of PS1 differentially regulate Kinase, an important enzyme regulating \*\*\*apoptosis\*\*\*.

L57 ANSWER 10 OF 23 MEDLINE DUPLICATE  
6 AN 1998082804 MEDLINE  
DN 98082804  
TI \*\*\*Presenilins\*\*\*, the endoplasmic reticulum, and \*\*\*neural\*\*\* apoptosis\*\*\* in Alzheimer's disease.  
AU Mattson M P; Guo Q; Funkawa K; Pedersen W A  
CS Department of Anatomy and Neurobiology, University of Kentucky, Lexington 40536-0230, USA.  
NC AG14554 (NIA)  
AG05144 (NIA)  
AG05119 (NIA)  
+  
SO JOURNAL OF NEUROCHEMISTRY, (1998 Jan) 70 (1) 1-14.  
Ref: 105  
Journal code: JAV. ISSN: 0022-3042.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, TUTORIAL)  
LA English  
FS Priority Journals  
EM 199804  
EW 19980401  
AB Many cases of autosomal dominant inherited forms of early-onset Alzheimer's disease are caused by mutations in the genes encoding \*\*\*presenilin\*\*\* -1 (PS-1; chromosome 14) and \*\*\*presenilin\*\*\* - 2\*\*\* (PS-2; chromosome 1). PSs are expressed in



\*\*\*neurons\*\*\* throughout the brain wherein they appear to be localized primarily to the endoplasmic reticulum (ER) of cell bodies and dendrites. PS-1 and PS-2 show high homology and are predicted to have eight transmembrane domains with the C terminus, N terminus, and a loop domain all on the cytosolic side of the membrane; an enzymatic cleavage of PSs occurs at a site near the loop domain. The normal function of PSs is unknown, but data suggest roles in membrane trafficking, amyloid precursor protein processing, and regulation of ER calcium homeostasis. Homology of PSs to the C. elegans gene sel-12, which is involved in Notch signaling, and phenotypic similarities of PS-1 and Notch knockout mice suggest a developmental role for PSs in the nervous system. When expressed in cultured cells and transgenic mice, mutant PSs promote increased production of a long form of amyloid beta-peptide (A beta1-42) that may possess enhanced amyloidogenic and neurotoxic properties. PS mutations sensitize cultured neural cells to \*\*\*apoptosis\*\*\* induced by trophic factor withdrawal, metabolic insults, and amyloid beta-peptide. The mechanism responsible for the proapoptotic action of mutant PSs may involve perturbed calcium release from ER stores and increased levels of oxidative stress. Recent studies of \*\*\*apoptosis\*\*\* in many different cell types suggest that ER signaling can modulate \*\*\*apoptosis\*\*\*. The evolving picture of PS roles in \*\*\*neural\*\*\* plasticity and Alzheimer's disease is bringing to the forefront the ER, an organelle increasingly recognized as a key regulator of \*\*\*neural\*\*\* plasticity and survival.

L57 ANSWER 11 OF 23 MEDLINE DUPLICATE  
 7 AN 1998346881 MEDLINE  
 DN 98346881  
 TI Localization and possible functions of \*\*\*presenilins\*\*\* in brain.  
 AU McGeer P L; Kawamata T; McGeer E G  
 CS Kinsmen Laboratory of Neurological Research, University of British Columbia, Vancouver, Canada.  
 SO REVIEWS IN THE NEUROSCIENCES, (1998) 9 (1) 1-15.  
 Ref: 83

Journal code: BYT. ISSN: 0334-1763.  
 CY ENGLAND: United Kingdom  
 DT Journal: Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LA English  
 FS Priority Journals  
 EM 199811  
 EW 19981104  
 AB \*\*\*Presenilin\*\*\* -1 (PS-1) is localized to chromosome 14 and \*\*\*presenilin\*\*\* -2\*\*\* (PS-2) to chromosome 1. Mutations in these genes, primarily in PS-1, account for an estimated 60% of early onset familial Alzheimer's disease cases (FAD), while FAD cases account for about 10% of all Alzheimer's disease (AD) cases. The mutations are minor but are 100% penetrant, suggesting that the proteins have acquired a toxic gain in function. The proteins have multiple transmembrane domains and have been reported to be localized to the Golgi apparatus, endoplasmic reticulum, nuclear membranes and cell surface membranes. They are thought to have functions associated with vesicular trafficking, Notch signaling and \*\*\*apoptosis\*\*\*. PS mutants show relative increases in the amount of A beta42/43 compared with A beta40 in plasma, fibroblasts and brain, observations which have been taken as a possible mechanism of their role in AD. In brain, the mRNAs for these two genes are localized primarily in \*\*\*neurons\*\*\*, with the strongest in situ hybridization signals being observed in the hippocampus, cerebellum and cerebral cortex. In AD, signals detected in the hippocampus are weaker than those in normals, while signals in the cerebellum are comparable. Immunohistochemical localization of the proteins is also primarily in \*\*\*neurons\*\*\*, and, at least for PS-1, is reduced in AD affected areas. PS-1 is localized to granular structures which are most abundant in cell bodies and dendrites. The functions of the \*\*\*presenilins\*\*\* are not yet known, but available evidence points to pyramidal \*\*\*neurons\*\*\* as the most logical site for pathological change in AD.

L57 ANSWER 12 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997:471351 BIOSIS

DN PREV19979770554  
 TI Effects of inducible expression on \*\*\*presenilins\*\*\* on cell proliferation and \*\*\*cell\*\*\* \*\*\*death\*\*\*  
 AU Kang, David E. (1); Kammescheidt, Anja; Koo, Edward H.  
 CS (1) Dep. Neurosci., Univ. Calif. San Diego, La Jolla, CA 92093 USA  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 824.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience, Part 1  
 New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295.  
 DT Conference; Abstract; Conference  
 LA English

L57 ANSWER 13 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997:471344 BIOSIS  
 DN PREV19979770547  
 TI Dissecting functional domains of \*\*\*presenilins\*\*\*  
 AU Hong, Chang-Sook; Koo, Edward H.  
 CS Dep. Neurosci., Univ. Calif. San Diego, La Jolla, CA 92093 USA  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 823.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience, Part 1  
 New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295.  
 DT Conference; Abstract; Conference  
 LA English

L57 ANSWER 14 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1997:471341 BIOSIS  
 DN PREV19979770544  
 TI Characterization of \*\*\*presenilin\*\*\* overexpressing cerebellar \*\*\*neural\*\*\* cells.  
 AU Weggen, S.; Diehlmann, A.; Ida, N.; Czech, C.; Weidemann, A.; Masters, C.; Wiestler, O. D.; Beyreuther, K.; Bayer, T. A.  
 CS Dep. Neuropathol., Univ. Bonn Med. Cent., Sigmund-Freud-Str. 25, 53105 Bonn Germany  
 SO Society for Neuroscience Abstracts, (1997) Vol. 23, No. 1-2, pp. 822.  
 Meeting Info.: 27th Annual Meeting of the Society for Neuroscience, Part 1  
 New Orleans, Louisiana, USA October 25-30, 1997  
 ISSN: 0190-5295.  
 DT Conference; Abstract; Conference  
 LA English

L57 ANSWER 15 OF 23 MEDLINE DUPLICATE  
 8 AN 1998067216 MEDLINE  
 DN 98067216  
 TI Cell and molecular neurobiology of \*\*\*presenilins\*\*\* : a role for the endoplasmic reticulum in the pathogenesis of Alzheimer's disease?

AU Mattson M P; Guo Q  
CS Sanders-Brown Research Center on Aging and Department of Anatomy and Neurobiology, University of Kentucky, Lexington, USA...  
MMattson@aging.coa.uky.edu  
NC NS30583 (NINDS)  
AG10836 (NIA)  
AG05144 (NIA)  
+  
SO JOURNAL OF NEUROSCIENCE RESEARCH, (1997 Nov 15)  
50 (4) 505-13. Ref: 71  
Journal code: KAC. ISSN: 0360-4012.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
DT General Review; (REVIEW)  
(LA English  
FS Priority Journals  
EM 199803  
EW 19980305  
AB Mutations in genes encoding \*\*\*presenilin\*\*\* -1 (PS-1) and \*\*\*presenilin\*\*\* -2 (PS-2) cause many cases of autosomal dominant inherited forms of early-onset Alzheimer's disease (AD). PSs are expressed in \*\*\*neurons\*\*\* throughout the nervous system, with differences in abundance among cell populations. PS-1 and PS-2 each have six to eight transmembrane domains and are localized mainly in the endoplasmic reticulum (ER). PSs may interact with cytoskeletal proteins and beta-amyloid precursor protein (APP) in ways consistent with roles in membrane trafficking and APP processing. Expression of mutant PSs in cultured cells and transgenic mice results in increased production of an amyloidogenic-cytotoxic form of amyloid beta-peptide (Abeta). Neural cells expressing mutant PSs exhibit increased sensitivity to \*\*\*apoptosis\*\*\* induced by trophic factor withdrawal and Abeta. The proapoptotic action of mutant PSs involves perturbed calcium release from ER stores and increased levels of oxidative stress. PS mutations may also suppress neurotransmitter synthesis in cholinergic \*\*\*neurons\*\*\*, suggesting a role in regulation of \*\*\*neuronal\*\*\* phenotype. Homology of PSs with the C. elegans gene sel-12 and phenotypic similarities of PS-1 and Notch knockout mice suggest a developmental role for PSs in somitogenesis. Collectively, the emerging data suggest intriguing roles of PSs in \*\*\*neuronal\*\*\* plasticity and \*\*\*cell\*\*\* \*\*\*death\*\*\*

and highlight the importance of the ER as a regulatory site involved in the pathogenesis of \*\*\*neuronal\*\*\* degeneration in AD.  
L57 ANSWER 16 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
AN 1997:395321 BIOSIS  
DN PREV199799694524  
TI Superoxide free radical and intracellular calcium mediate A-beta-1-42 induced endothelial toxicity.  
AU Suo, Zhiming (1); Fang, Chunhong; Crawford, Fiona; Mullan, Mike  
CS (1) Roskamp Lab., Dep. Psychiatry, 3515 E. Fletcher Ave., Univ. South Fla., Tampa, FL 33613 USA  
SO Brain Research, (1997) Vol. 762, No. 1-2, pp. 144-152.  
ISSN: 0006-8993.  
DT Article  
LA English  
AB The 39-42 amino acid residue amyloid beta peptide (A-beta), the major protein component in senile plaques and cerebrovascular amyloidosis in the brain in Alzheimer's disease (AD), has been shown to be neurotoxic in vitro. Accumulating data from several areas suggest that dysfunction and damage may also play a significant role in the AD process. For instance, we have recently demonstrated enhanced vasoconstriction and resistance to relaxation in intact rat aorta treated with A-beta (Thomas et al., beta-Amyloid-mediated vasoactivity and vascular endothelial damage, Nature, 380 (1996) 168-171). Significant vessel damage occurred after thirty minutes of exposure, but could be prevented with superoxide dismutase. To further investigate the role of A beta toxicity on endothelial cells, we have applied AP peptides to cultures of human aortic endothelial cells (HAEC). Our results show that both A beta-1-42 and A beta-25-35 are toxic to HAEC in a time- and dose-dependent manner, and that this toxicity can be partially prevented by the calcium channel blocker, verapamil, and the antioxidant, superoxide dismutase. The common form of A beta, A beta-1-40, which has been shown to be neurotoxic, is much less toxic to HAEC. AG toxicity to HAEC occurs within 30 min of treatment with relatively lower doses than those usually observed in primary cultured \*\*\*neurons\*\*\* and vascular smooth muscle cells. It was recently reported that a variety of mutations in the

beta-amyloid protein precursor gene and the \*\*\*presenilin\*\*\* -1 and -2 genes linked to early-onset familial AD cause an increase in the plasma concentration of A beta-1-42 in mutation carriers (Scheuner et al., Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vitro by the \*\*\*presenilin\*\*\* -1 and 2 and APP mutations linked to familial Alzheimer's disease, Nature Med., 2 (1996) 864-8701. Human aortic endothelial cells are more sensitive to A beta-1-42 than A beta-1-40, via a pathway involving an excess of superoxide free radicals and influx of extracellular calcium. Finally, we have evidence that both apoptotic and necrotic processes are activated by the AP peptides in these endothelial cells.  
L57 ANSWER 17 OF 23 MEDLINE  
AN 97134506 MEDLINE  
DN 97134506  
TI Dissecting how presenilins function--and malfunction [news].  
AU Marx J  
SO SCIENCE, (1996 Dec 13) 274 (5294) 1838-40.  
Journal code: UJ7. ISSN: 0036-8075.  
CY United States  
DT News Announcement  
LA English  
FS Priority Journals; Cancer Journals  
EM 199703

L57 ANSWER 18 OF 23 MEDLINE  
9  
AN 97094374 MEDLINE  
DN 97094374  
TI Participation of \*\*\*presenilin\*\*\* \*\*\*2\*\*\* in \*\*\*apoptosis\*\*\* : enhanced basal activity conferred by an Alzheimer mutation.  
AU Wolozin B; Iwasaki K; Vito P; Ganjei J K; Lacan a E; Sunderland T; Zhao B; Kusiak J W; Wasco W; D'Adamio L  
CS Unit on Alzheimer Biology, Laboratory of Clinical Science, National Institute of Mental Health, Building 10, Room 3D41, 9000 Rockville Pike, Bethesda, MD 20892, USA.. Idadamio@atlas.nimh.nih.gov  
SO SCIENCE, (1996 Dec 6) 274 (5293) 1710-3.  
Journal code: UJ7. ISSN: 0036-8075.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals; Cancer Journals  
EM 199703  
AB Overexpression of the familial Alzheimer's disease gene

\*\*\*Presenilin\*\*\*  
 \*\*\*2\*\*\* (PS2) in nerve growth factor-differentiated PC12 cells increased  
 \*\*\*apoptosis\*\*\* induced by trophic factor withdrawal or beta-amyloid.  
 Transfection of antisense PS2 conferred protection against \*\*\*apoptosis\*\*\* induced by trophic withdrawal in nerve growth factor-differentiated or amyloid precursor protein-expressing PC12 cells.  
 The apoptotic \*\*\*cell\*\*\* \*\*\*death\*\*\* induced by PS2 protein was sensitive to pertussis toxin, suggesting that heterotrimeric GTP-binding proteins are involved. A PS2 mutation associated with familial Alzheimer's disease was found to generate a molecule with enhanced basal apoptotic activity. This gain of function might accelerate the process of neurodegeneration that occurs in Alzheimer's disease, leading to the earlier age of onset characteristic of familial Alzheimer's disease.  
 L57 ANSWER 19 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1996:547081 BIOSIS  
 DN PREV199699269437  
 TI Alzheimer's PS-1 L286V mutation increases \*\*\*neuronal\*\*\* vulnerability to A-beta toxicity and trophic factor withdrawal-induced \*\*\*apoptosis\*\*\*  
 AU Guo, O. (1); Sopher, B. L.; Furukawa, K.; Robinson, N.; Martin, G. M.; Mattson, M. P.  
 CS (1) Sanders-Brown Res. Cent. Aging, Univ. Kentucky, Lexington, KY 40536 USA  
 SO Society for Neuroscience Abstracts, (1996) Vol. 22, No. 1-3, pp. 1664.  
 Meeting Info.: 26th Annual Meeting of the Society for Neuroscience  
 Washington, D.C., USA November 16-21, 1996  
 ISSN: 0190-5295.  
 DT Conference  
 LA English  
 L57 ANSWER 20 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1996:547073 BIOSIS  
 DN PREV199699269429  
 TI Genetic dissection of \*\*\*presenilin\*\*\* functions in a \*\*\*neuronal\*\*\* precursor cell line.  
 AU Hong, Chang-Sook; Koo, Edward H.  
 CS Harvard Medical Sch., Cent. Neurol. Dis., Brigham Women's Hosp., Boston, MA 02115 USA  
 SO Society for Neuroscience Abstracts, (1996) Vol. 22, No. 1-3, pp. 1662.  
 Meeting Info.: 26th Annual Meeting of the Society for

Neuroscience  
 Washington, D.C., USA November 16-21, 1996  
 ISSN: 0190-5295.  
 DT Conference  
 LA English  
 L57 ANSWER 21 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1996:552862 BIOSIS  
 DN PREV199699275218  
 TI Characterization and analysis of \*\*\*presenilin\*\*\* \*\*\*2\*\*\* in mammalian cells: Effects of expression on cell viability.  
 AU Crowley, A. C. (1); Merriam, D. E.; Kovacs, D. M.; Kim, T.-W.; Wasco, W.  
 CS (1) Genetics Aging Unit, Dep. Neurology, Mass. General Hosp.-East, Boston, MA 02129 USA  
 SO Society for Neuroscience Abstracts, (1996) Vol. 22, No. 1-3, pp. 1437.  
 Meeting Info.: 26th Annual Meeting of the Society for Neuroscience  
 Washington, D.C., USA November 16-21, 1996  
 ISSN: 0190-5295.  
 DT Conference  
 LA English  
 L57 ANSWER 22 OF 23 BIOSIS COPYRIGHT 1999 BIOSIS  
 AN 1996:552860 BIOSIS  
 DN PREV199699275216  
 TI Developmentally regulated expression of Alzheimer-related \*\*\*presenilin\*\*\* genes (PS-1 and PS-2) matches notch in mouse brain.  
 AU Berezovskaja, O. (1); Page, K.; Xia, M. Q.; Berezovskii, V.; Wasco, W.; Tanzi, R.; Hyman, B. T.  
 CS (1) Dep. Neurol., Mass. General Hosp., Boston, MA USA  
 SO Society for Neuroscience Abstracts, (1996) Vol. 22, No. 1-3, pp. 1436.  
 Meeting Info.: 26th Annual Meeting of the Society for Neuroscience  
 Washington, D.C., USA November 16-21, 1996  
 ISSN: 0190-5295.  
 DT Conference  
 LA English  
 L57 ANSWER 23 OF 23 CAPLUS COPYRIGHT 1999 ACS  
 AN 1997:227701 CAPLUS  
 DN 126:304309  
 TI Alzheimer's Disease: melting pot or mosaic?  
 AU Blass, John P.  
 CS Burke Medical Research Institute, Cornell University Medical College, White Plains, NY, 10605, USA  
 SO Alzheimer's Dis. Rev. [Electronic Publication] (1996), 1(1/2), 17-20  
 CODEN: ADREFN  
 URL: <http://www.coa.uky.edu/ADReview/blass.htm>

PB Sanders-Brown Center on Aging, University of Kentucky  
 DT Journal, General Review; (online computer file)  
 LA English  
 AB A brief review with 18 refs. Alzheimer's Disease (AD), like the proverbial elephant, can be described in a no. of ways, all of which are accurate and all of which are incomplete. AD can be described, correctly, as: a loss of synapses; a premature loss of \*\*\*neurons\*\*\* in a selectively vulnerable pattern, often assocd. with \*\*\*apoptosis\*\*\* and other mechanisms of \*\*\*cell\*\*\* \*\*\*death\*\*\* which involve free radicals; a disorder of free radical metab. ("oxidative stress"); a cerebrometabolic disease involving impaired glucose/energy metab.; a cytoskeletal disease; a form of cerebral amyloidosis; a disorder of signal transduction; a disorder of cerebral calcium homeostasis; a membrane disorder; and a disorder of neurotransmission, with prominent impairment of cholinergic function and more variable but typical involvement of other neurotransmitter systems. Mol. genetic studies to date suggest that the most important trait predisposing to the common, late onset form of AD is possession of the 4 allele of the ApoE gene. Studies in progress suggest the possibility that a genetic abnormality in a component of the Krebs tricarboxylic acid cycle (the  $\alpha$ -ketoglutarate dehydrogenase complex) is also be important factor in the common, late onset form of AD. In the rarer, early onset familial forms of AD (FAD), the most common genetic abnormalities appear to be the \*\*\*presenilin\*\*\* -1 or \*\*\*presenilin\*\*\* -2 genes, which seems likely from the predicted amino acid sequences to lead to abnormalities in signal transduction or cellular calcium homeostasis. Abnormalities in the gene for the amyloid precursor protein were the first mutations assocd. with AD, but in fact have proven to be rare even in FAD. Based on currently available data, any one of the mechanisms listed above could be proposed to be the central step in the pathophysiol. of AD, with other mechanisms acting through their effects on that "mainstream" abnormality. An alternative hypothesis is that a complex mosaic of abnormalities leads to the pattern of brain scarring which characterizes AD. Different parts of

the mosaic may have or more or less important roles, depending on genetic  
endowment and environmental factors. Different parts of the  
mosaic may interact with each other. For instance, the abnormality in  
glucose/energy metab. in AD which the authors and others have been studying may  
well influence the progression of the disease by diminishing the ability  
of nerve cells to adapt to challenges ("stressors") created by other  
mechanisms which are part of AD. Precedents for this "mosaic  
hypothesis" include other complex degenerative diseases which are better  
understood than AD, such as atherosclerosis or clotting disorders.

=> d his

(FILE 'HOME' ENTERED AT 15:56:10 ON 11 APR 1999)

FILE 'MEDLINE' ENTERED AT 15:56:17 ON 11 APR 1999

L1 407 S PRESENILIN7/AB,BI  
L2 0 S L1 AND ADVANCED GLYCATION END  
PRODUCT/AB,BI  
L3 0 S L1 AND RAGE/AB,BI  
L4 71 S ADVANCED GLYCATION END PRODUCT/AB,BI  
L5 478 S L1 OR L4  
L6 114 S L5 AND NEURON7/AB,BI  
L7 111 S L1 AND NEURON7/AB,BI  
L8 3 S L4 AND NEURON7/AB,BI  
L9 4 S (L4)(3A)(RECEPTOR#)/AB,BI

FILE 'MEDLINE, EMBASE, BIOSIS, WPIDS, CAPLUS'  
ENTERED AT 16:02:33 ON 11  
APR 1999

L10 0 S L1 AND L9  
L11 10 S MUTANT PRESENILIN-2/AB,BI  
L12 4 DUP REM L11 (6 DUPLICATES REMOVED)  
E STERN DAVID/AU  
L13 162 S E3  
L14 0 S L13 AND PRESENILIN7/AB,BI  
E YAN SHI DU/AU  
L15 80 S E2-E3  
L16 0 S L15 AND PRESENILIN7/AB,BI  
E WOLOZIN, BENJAMIN/AU  
E WOLOZIN, B/AU  
E WOLOZIN B/AU  
L17 201 S E3-E10  
L18 23 S L17 AND PRESENILIN/AB,BI  
L19 12 DUP REM L18 (11 DUPLICATES REMOVED)  
E STERN D/AU  
L20 527 S E3  
L21 1 S L20 AND PRESENILIN/AB,BI  
L22 56 S L9  
L23 5 S L22 AND NEURON7/AB,BI

L24 5 DUP REM L23 (0 DUPLICATES REMOVED)  
L25 1 S L9 AND GLIAL/AB,BI  
L26 1 S L9 AND MICROGLIAL/AB,BI  
L27 1 S L9 AND ASTROCYTE#/AB,BI  
L28 15 S L9 AND ENDOTHELIAL/AB,BI  
L29 15 DUP REM L28 (3 DUPLICATES REMOVED)  
L30 6 S L9 AND MONONUCLEAR/AB,BI  
L31 6 DUP REM L30 (0 DUPLICATES REMOVED)  
L32 4 S L9 AND TUMOR#/AB,BI  
L33 4 DUP REM L32 (0 DUPLICATES REMOVED)  
L34 2 S L9 AND PC12/AB,BI  
L35 2 DUP REM L34 (0 DUPLICATES REMOVED)  
L36 88 S L1 AND PC12/AB,BI  
L37 0 S L36 AND L9  
L38 37 S L1 AND TUMOR#/AB,BI  
L39 31 DUP REM L38 (6 DUPLICATES REMOVED)  
L40 8 S L1 AND MONONUCLEAR/AB,BI  
L41 2 DUP REM L40 (6 DUPLICATES REMOVED)  
L42 10 S L1 AND ENDOTHELIAL/AB,BI  
L43 4 DUP REM L42 (6 DUPLICATES REMOVED)  
L44 61 S L1 AND ASTROCYTE#/AB,BI  
L45 26 DUP REM L44 (35 DUPLICATES REMOVED)  
L46 27 S L44 AND PRESENILIN-2/AB,BI  
L47 13 DUP REM L46 (14 DUPLICATES REMOVED)  
L48 19 S L1 AND MICROGLIA#/AB,BI  
L49 7 DUP REM L48 (12 DUPLICATES REMOVED)  
L50 41 S L1 AND GLIA#/AB,BI  
L51 23 DUP REM L50 (18 DUPLICATES REMOVED)  
L52 16 S L50 AND PRESENILIN-2/AB,BI  
L53 7 DUP REM L52 (9 DUPLICATES REMOVED)  
L54 500 S L1 AND NEURON7/AB,BI  
L55 135 S L54 AND PRESENILIN-2/AB,BI  
L56 43 S L55 AND (CELL DEATH OR APOPTOSIS)/AB,BI  
L57 23 DUP REM L56 (20 DUPLICATES REMOVED)

=>

--Logging off of STN--

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	ENTRY	SINCE FILE	TOTAL
FULL ESTIMATED COST	443.85	448.19	
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)			
SINCE FILE	TOTAL	ENTRY	SESSION
CA SUBSCRIBER PRICE	-18.21	-18.21	-18.21
STN INTERNATIONAL LOGOFF AT 16:33:10 ON 11 APR 1999			